arXiv:math/0412012v2 [math.OA] 4 Dec 2004

COMPACT QUANTUM ERGODIC SYSTEMS

REIJI TOMATSU

Abstract. We develop theory of multiplicity maps for compact quantum groups, as an application, we obtain a complete classification of right coideal C^*-algebras of $C(SU_q(2))$ for $q \in [-1,1] \setminus \{0\}$. They are labeled with Dynkin diagrams, but classification results for positive and negative cases of q are different. Many of the coideals are quantum spheres or quotient spaces by quantum subgroups, but we do have other ones in our classification list.

1. Introduction

The aim of this paper is studying compact quantum ergodic systems and classifying right coideals of $C(SU_q(2))$ for $q \in [-1,1] \setminus \{0\}$. Compact quantum groups have been introduced by S. L. Woronowicz [Wo2]. They are considered as the continuous function algebra on a non commutative space of a compact quantum group. The continuous function algebra on a compact quantum group is denoted by $C(G)$ and we have the coproduct $\delta : C(G) \longrightarrow C(G) \otimes C(G)$. In this paper, a quantum group discovered by V. G. Drinfel'd and M. Jimbo is called a quantum universal enveloping algebra because it is considered as a deformation of the universal enveloping algebra of a Lie algebra [D], [J]. A C^*-subalgebra $A \subset C(G)$ is called a right coideal C^*-algebra or often simply a right coideal if it is invariant by the right action of G, that is, we have $\delta(A) \subset A \otimes C(G)$. Let A be a unital C^*-algebra and G be a compact quantum group. A triple $\{A, G, \alpha\}$ is a compact quantum covariant system if a unital faithful $*$-homomorphism $\alpha : A \longrightarrow A \otimes C(G)$ satisfies $(\mathrm{id} \otimes \delta) \circ \alpha = (\alpha \otimes \mathrm{id}) \circ \alpha$. Moreover if it satisfies $A^\alpha := \{a \in A \mid \alpha(a) = a \otimes 1\} = \mathbb{C}1_A$, then it is called a compact quantum ergodic system. When we consider the classical compact group case, the invariant state on a C^*-algebra A must be a faithful tracial state and multiplicities of irreducible representations are bounded by dimensions of their representation spaces [H-L-S]. However in the quantum case the invariant state has no longer tracial property and the multiplicities are bounded by quantum dimensions which are larger than or equal to usual dimensions. Keeping in mind these phenomena, one shall notice there are rather differences between classical compact group case and compact quantum group case. It is still a fascinating problem to construct an example of a compact quantum ergodic system which has a multiplicity of an irreducible representation strictly larger than its dimension. In [Wa1], he has developed the theory of multiplicity maps and it has worked very well in the classification program of ergodic systems of $SU(2)$ [Wa2]. Since multiplicity maps are defined via equivariant K-theory, one easily has its quantum version by using the work due to, for example, [BS1, B, V]. The most nontrivial important problem is whether there exists a common eigenvector of multiplicity maps or not. In compact group case its existence is guaranteed by the tracial invariant state. For that problem we show that such a vector exists if A has a (not necessarily faithful) tracial state and $G = SU_q(2)$ in Theorem 4.10. If we consider a right coideal $A \subset C(G)$, then we get a compact quantum ergodic system $\{A, \delta, C(G)\}$. Fortunately we can show there always exists a desired common eigenvector in Theorem 4.21 in this case. With this result we can proceed to the classification of right coideals of $C(SU_q(2))$. In the classical case we know the continuous function algebra on the homogeneous space $H \setminus G$ by a closed subgroup $H \subset G$ gives a right coideal and its converse holds via the Gelfand-Naimark theorem on abelian C^*-algebras. However in the quantum case, such correspondence does not hold in general. For example, in [Po1] he has constructed one-parameter family of, so called, quantum spheres $C(S^2_{q,\lambda})$ which are right coideals and most of them are not obtained by taking quotient by subgroups. Therefore it is interesting to investigate the other non-quotient type right coideals. The aim of the latter half of this paper is giving the complete classification of right coideals of $C(SU_q(2))$. Most important information of a right coideal is its spectral pattern, that is, multiplicities of irreducible representations. We will see this information is remembered by equivariant K-theory and we classify right coideals into the several types by connected graphs of norm 2 as is used in the classification of ergodic systems of $SU(2)$. They are labeled by closed subgroups of $SU(2)$ or $SU_{-1}(2)$ as McKay diagrams with respect to the fundamental two-dimensional representation. When we work on the negative q case, it is needed to treat the graphs with a single loop at a vertex. Then we carry out case-by-case study of them. In that procedure one has to be careful of the essential effect of $|q| \neq 1$. In fact, right coideals of some types such as the regular polyhedrons do not appear in those case. We state the main result on this classification.

(1) The case $0 < q < 1$: A right coideal must be one of type 1, $SU(2)$, $\mathbb{T}_n$, $\mathbb{T}$ and D^*_∞. When it is of type $\mathbb{T}$, then it is one of series of the quantum spheres. Otherwise it is uniquely determined by the type.

(2) The case $-1 < q < 0$: A right coideal must be one of type 1, $SU(2)$, $\mathbb{T}_n$, $\mathbb{T}$, D^*_∞ and D_1. When it is of type $\mathbb{T}$, then it is one of series of the quantum spheres. Otherwise it is uniquely determined by the type.

(3) The case $q = -1$: If a right coideal A is not of type $\mathbb{T}_n$ (odd $n \geq 3$) or D_n (odd $n \geq 1$), there exists a closed subgroup H in $SO_{-1}(3)$ such that A is $C(H \setminus SO_{-1}(3))$. If a right coideal A is of type $\mathbb{T}_n$ (odd $n \geq 3$), A is conjugated to $C(\mathbb{T}_n \setminus SU_{-1}(2))$ or $C^*(\eta^{\frac{n}{2}}, \widehat{\eta}^{\frac{n}{2}})$. If a right coideal A is of type D_1, then A is conjugated to $C(D_1 \setminus SU_{-1}(2))$. If a right coideal A is of type D_n (odd $n \geq 3$), A is conjugated to $C(D_n \setminus SU_{-1}(2))$ or $C^*(\eta^{\frac{n}{2}})$. Here conjugation is given by the left action β^L_z of the maximal torus for some $z \in \mathbb{T}$.

In each of the above cases, uniqueness means conjugation by the left action of the maximal torus $\mathbb{T}$. We find right coideals which are of type D_∞^* in the case $0 < q < 1$, of type D_∞^* and D_1 in the case $-1 < q < 0$ and $C^*(\eta^{\frac{n}{2}}, \widehat{\eta}^{\frac{n}{2}})$, $C^*(\eta^{\frac{n}{2}})$ in the case $q = -1$. They are not the quantum spheres nor the quotient spaces. In the case $q = -1$, a right coideal which is not of type $\mathbb{T}_n$ (odd $n \geq 3$) is $SU_{-1}(2)$-isomorphic to each other in the same type, however, $C(\mathbb{T}_n \setminus SU_{-1}(2))$ is not $SU_{-1}(2)$-isomorphic to $C^*(\eta^{\frac{n}{2}}, \widehat{\eta}^{\frac{n}{2}})$.

We briefly explain the contents of study of each section. In section 2 we collect required facts on compact quantum groups and their actions. In section 3 we study general compact quantum ergodic systems, especially hom-spaces of them. In section 4 we consider quantum version of the theory of multiplicity maps. In section 5 we give a summary for representation theory of $SU_q(2)$. In sections 6, 7 and 8 we carry out classification of right coideals of $C(SU_q(2))$ for $0 < q < 1$, $-1 < q < 0$ and $q = -1$, respectively. In Appendix we list all the connected graphs of norm 2. If one has only interest in classification of right coideals, he or she can skip results in sections 1, 2, 3 and 4 except for Corollary 4.21.

Acknowledgements. The author is grateful to Yasuyuki Kawahigashi, Shigeru Yamagami and Junichi Shiraishi for permanent encouragement and fruitful discussions. This work was partially supported by Research Fellowship for Young Scientists of the Japan Society for the Promotion of Science.

2. PRELIMINARIES

We collect basic notions on compact quantum groups and their actions. Our standard references are [Wo2] for theory of compact quantum groups and [BS2] for their actions. In this paper we only treat minimal tensor products for C^*-algebras and use the notation simply $\otimes$. Although it is not essential, separability of compact quantum groups is always assumed.

Definition 2.1. [Wo2, p.853] Let A be a unital C^*-algebra and $\delta : A \longrightarrow A \otimes A$ be a $*$-homomorphism. If they satisfy the following conditions, the pair (A, δ) is called a compact quantum group.

(1) The map δ satisfies coassociativity condition:

$$(\delta \otimes \mathrm{id}_A) \circ \delta = (\mathrm{id}_A \otimes \delta) \circ \delta.$$

(2) The vector spaces $\delta(A)(\mathbb{C} \otimes A)$ and $\delta(A)(A \otimes \mathbb{C})$ are dense in $A \otimes A$.

Let (A, δ) be a compact quantum group. Then there exists the unique state h which satisfies the following invariance condition:

$$(h \otimes \mathrm{id}_A) \circ \delta(a) = (\mathrm{id}_A \otimes h) \circ \delta(a) = h(a)1_A, \text{for all } a \in A.$$

This state is called the *Haar state*. We always consider a compact quantum group whose Haar state is faithful. A compact quantum group (A, δ) is often regarded as the continuous function algebra on a non-commutative space G and we write $G = (A, \delta)$ and $A = C(G)$.

Let H be a Hilbert space and v be a unitary in $M(\mathbb{K}(H) \otimes A)$. A unitary v is called a *unitary representation* of G if it satisfies the following equality:

$$(\mathrm{id}_H \otimes \delta)(v) = v_{12} v_{13}.$$

Let $\{v_i\}_{i \in I}$ be a family of unitary representations on Hilbert spaces $\{H_{v_i}\}_{i \in I}$, respectively. The direct sum representation $\prod_{i \in I} v_i$ is defined as the direct sum of unitary operators via the natural inclusion $\prod_{i \in I} M(\mathbb{K}(H_{v_i}) \otimes C(G)) \subset M(\mathbb{K}(\bigoplus_{i \in I} H_v) \otimes C(G))$. Let v, w be unitary representations on Hilbert spaces H_v, H_w respectively. The tensor product representation $v \dot{\otimes} w$ is defined as a unitary operator $v_{13} w_{23}$ which is an element of $M(\mathbb{K}(H_v \otimes H_w) \otimes C(G))$. An operator T in $\mathbb{B}(H_v, H_w)$ is called an *intertwiner* of v and w if it satisfies $(T \otimes 1)v = w(T \otimes 1)$. The set of intertwiners between v and w is written as $\mathrm{Hom}(v, w)$ and it is called a *hom-space*. For one unitary representation v its hom-space $\mathrm{Hom}(v, v)$ becomes a C^*-subalgebra of $\mathbb{B}(H_v)$. If the space is trivial (that is, $\mathrm{Hom}(v, v) = \mathbb{C}1_{H_v}$), we say that the unitary representation v is *irreducible*. Let v be a unitary representation of a compact quantum group G and v is a direct sum of finite dimensional irreducible representations ([Wo2, p.864]). In particular any irreducible unitary representation is finite dimensional.

In the set of all irreducible unitary representations of G, we define unitary equivalence relation naturally and denote its quotient set by $\widehat{G}$. We can select one unitary representation $w(\pi) \in \mathbb{B}(H_\pi) \otimes C(G)$ for each equivalence class $\pi \in \widehat{G}$. Let $\{\xi_i\}_{i \in I_\pi}$ be an orthonormal basis for H_π. We can identify $w(\pi)$ with the matrix $\{w(\pi)_{i,j}\}_{i,j \in I_\pi}$ where $w(\pi)_{i,j}$ are elements of $C(G)$. From the definition of unitary representation we obtain the following equality:

$$\delta(w(\pi)_{i,j}) = \sum_{k \in I_\pi} w(\pi)_{i,k} \otimes w(\pi)_{k,j}, \text{ for all } i, j \in I_\pi.$$

Let us define the *smooth function algebra* $A(G) = \mathrm{span}\{w(\pi)_{i,j} \mid i, j \in I_\pi,\ \pi \in \widehat{G}\}$. It is in fact a unital $*$-subalgebra dense in $C(G)$ and the set $\{w(\pi)_{i,j} \mid i, j \in I_\pi, \pi \in \widehat{G}\}$ is a linear basis for $A(G)$. Introducing the maps $\varepsilon : A(G) \longrightarrow \mathbb{C}$ and $\kappa : A(G) \longrightarrow A(G)$ by

$$\varepsilon(w(\pi)_{i,j}) = \delta_{i,j},\ \kappa(w(\pi)_{i,j}) = w(\pi)^*_{j,i}$$

for each $\pi \in \widehat{G}$ and $i, j \in I_\pi$ gives the algebra $A(G)$ a *Hopf* $*$-*algebra* structure:

(1) The map $\delta : A(G) \longrightarrow A(G) \otimes A(G)$ satisfies coassociativity:

$$(\delta \otimes \mathrm{id}) \circ \delta = (\mathrm{id} \otimes \delta) \circ \delta.$$

(2) The map (called the *counit*) $\varepsilon : A(G) \longrightarrow \mathbb{C}$ is a unital $*$-homomorphism and satisfies

$$(\varepsilon \otimes \mathrm{id}) \circ \delta = (\mathrm{id} \otimes \varepsilon) \circ \delta = \mathrm{id}.$$

(3) The map (called the *antipode*) $\kappa : A(G) \longrightarrow A(G)$ is a linear anti-multiplicative map and satisfies

$$m(\kappa \otimes \mathrm{id}) \circ \delta = m(\mathrm{id} \otimes \kappa) \circ \delta = \varepsilon$$

and

$$\kappa(\kappa(a^*)^*) = a \text{ for all } a \in A(G).$$

Let $A(G)^*$ be an algebraic dual space of $A(G)$. For $a \in A(G)$ and $\theta, \omega_1, \omega_2 \in A(G)^*$, we define their convolution products and involution:

$$\theta * a = (\mathrm{id} \otimes \theta)(\delta(a)),$$
$$a * \theta = (\theta \otimes \mathrm{id})(\delta(a)),$$
$$\omega_1 * \omega_2 = (\omega_1 \otimes \omega_2) \circ \delta,$$
$$\theta^*(a) = \overline{\theta(\kappa(a)^*)}.$$

With these operations the dual space $A(G)^*$ becomes a unital $*$-algebra (its unit is the counit) and acts on $A(G)$ from the left and right. On the smooth function algebra $A(G)$ there exists a unique family of characters $\{f_z\}_{z\in\mathbb{C}}$which are called *Woronowicz characters* with following properties:

(1) $f_z(1) = 1$ for all $z \in \mathbb{C}$.
(2) For any element $a \in A(G)$, the mapping $z \in \mathbb{C} \mapsto f_z(a) \in \mathbb{C}$ is an entire holomorphic function.
(3) $f_{z_1} * f_{z_2} = f_{z_1+z_2}$ for all $z_1, z_2 \in \mathbb{C}$.
(4) $f_0 = \varepsilon$.
(5) $f_z(\kappa(a)) = f_{-z}(a)$ for all $z \in \mathbb{C}, a \in A(G)$.
(6) $f_z(a^*) = \overline{f_{-\bar{z}}(a)}$ for all $z \in \mathbb{C}, a \in A(G)$.

We can define one-parameter automorphism groups $\{\sigma^h_t\}_{t\in\mathbb{R}}$ and $\{\tau_t\}_{t\in\mathbb{R}}$ and $*$-anti-multiplicative linear map R by

$$\sigma^h_t(a) = f_{it} * a * f_{it} \text{ for all } a \in A(G),$$
$$\tau_t(a) = f_{-it} * a * f_{it} \text{ for all } a \in A(G),$$
$$R(a) = f_{\frac{1}{2}} * \kappa(x) * f_{-\frac{1}{2}} \text{ for all } a \in A(G).$$

They are called the *modular automorphism group*, the *scaling automorphism group* and the *unitary antipode* respectively. They are norm continuously extendable to the continuous function algebra $C(G)$. The Haar state h is a σ^h-KMS state: for any $a, b \in A(G)$, we have

$$h(ab) = h(\sigma^h_i(b)a).$$

Relations of these maps are as follows.

$$\sigma^h_s \circ \tau_t = \tau_t \circ \sigma^h_s \text{ for all } s, t \in \mathbb{R},$$
$$R \circ \sigma^h_s = \sigma^h_s \circ R,\ R \circ \tau_t = \tau_t \circ R,$$
$$\kappa = R \circ \tau_{-\frac{i}{2}} = \tau_{-\frac{i}{2}} \circ R.$$

Let v be a unitary representation on a finite dimensional Hilbert space H_v and $j_v : H_v \longrightarrow \overline{H_v}$ be a conjugate unitary map where $\overline{H_v}$ is the conjugate Hilbert space. The transpose map $t_v : \mathbb{B}(H_v) \longrightarrow \mathbb{B}(\overline{H_v})$ is defined by $t_v(x) = j_v x^* j_v^{-1}$. We define the *contragradient representation* (which is not necessarily a unitary representation) v^c and the conjugate unitary representation $\bar{v}$ of a finite dimensional unitary representation v by

$$v^c = (t_v \otimes \kappa)(v),\ \ \bar{v} = (t_v \otimes R)(v).$$

The *F-matrix* of a finite dimensional unitary representation $v \in \mathbb{B}(H_v) \otimes C(G)$ is defined by

$$F_v = (\mathrm{id} \otimes f_1)(v)\ \ \in \mathbb{B}(H_v).$$

This matrix is strictly positive definite. It is known that $F_v^z = (\mathrm{id} \otimes f_z)(v)$ for any $z \in \mathbb{C}$ and $\mathrm{Tr}(F_v) = \mathrm{Tr}(F_v^{-1})$. For a unitary irreducible representation v, its second contragradient (not necessarily unitary) representation v^{cc} is also irreducible and equivalent to v. In fact we have $\mathrm{Hom}(v, v^{cc}) = \mathbb{C}F_v$. The value $\mathrm{Tr}(F_v)$ is called the *quantum dimension* and denoted by D_v. Since we have $F_{\bar{v}} = t_v(F_v^{-1})$, the quantum dimension of v is equal to that of $\bar{v}$. We write d_v for the usual dimension of the vector space H_v. The quantum dimension is larger than or equal to the usual dimension. When we fix a selection of $\{w(\pi)\}_{\pi \in \widehat{G}}$, we write H_π, F_π, D_π and d_π for $H_{w(\pi)}$, $F_{w(\pi)}$, $D_{w(\pi)}$ and $d_{w(\pi)}$, respectively.

Using F-matrices, we have the following equalities about the Haar state:

$$h(w(\pi)_{i,j} w(\rho)^*_{r,s}) = D_\pi^{-1} (F_\pi)_{s,j} \delta_{\pi,\rho} \delta_{i,r}$$

$$h(w(\pi)^*_{i,j} w(\rho)_{r,s}) = D_\pi^{-1} (F_\pi^{-1})_{r,i} \delta_{\pi,\rho} \delta_{j,s}$$

for any $\pi, \rho \in \widehat{G}$, $i, j \in I_\pi$ and $r, s \in I_\rho$.

For the tensor product $H \otimes K$ of two Hilbert spaces H and K, we define its conjugate unitary map by $j_{H \otimes K} : H \otimes K \longrightarrow \overline{K} \otimes \overline{H}$, $j_{H \otimes K}(\xi \otimes \eta) = j_K \eta \otimes j_H \xi$. Then for two finite dimensional unitary representation v and w, we obtain $\overline{v \dot{\otimes} w} = \bar{w} \dot{\otimes} \bar{v}$.

Let $L^2(G)$ be the GNS-representation space of $C(G)$ associated to the Haar state h. Its cyclic vector is denoted by $\hat{1}_h$. We define the *modular conjugation* J and $\hat{J}$ by

$$\begin{aligned} J\, x \hat{1}_h &= \sigma^h_{\frac{i}{2}}(x)^* \hat{1}_h\,, \\ \hat{J}\, x \hat{1}_h &= R(x)^* \hat{1}_h \end{aligned}$$

where x is an element of $A(G)$. We also define a unitary $U = J\hat{J} = \hat{J}J$. When we consider $L^\infty(G)$ which is the σ-weak closure of $C(G)$ in $\mathbb{B}(L^2(G))$, it becomes a *von Neumann algebraic compact quantum group.* Precisely speaking, the coproduct δ and the Haar state h extend to $L^\infty(G)$ as a normal $*$-homomorphism and the invariant faithful normal state respectively. By the invariance of h, we define the following two unitaries on $L^2(G) \otimes L^2(G)$: for all $x, y \in C(G)$,

$$V_\ell^*(x\hat{1}_h \otimes y\hat{1}_h) = \delta(y)(x\hat{1}_h \otimes \hat{1}_h)\,,$$
$$V(x\hat{1}_h \otimes y\hat{1}_h) = \delta(x)(\hat{1}_h \otimes y\hat{1}_h)\,.$$

They satisfy the following *pentagonal equality* and therefore are called *multiplicative unitaries*:

$$V_{12}V_{13}V_{23} = V_{23}V_{12}\,.$$

We also use other unitaries:

$$\begin{aligned} \widetilde{V} &= \Sigma(1 \otimes U)V(1 \otimes U)\Sigma\,, \\ W &= (U \otimes 1)V(U \otimes 1)\,, \end{aligned}$$

where $\Sigma : L^2(G) \otimes L^2(G) \longrightarrow L^2(G) \otimes L^2(G)$ is a flipping unitary $\xi \otimes \eta \mapsto \eta \otimes \xi$. The unitary $\widetilde{V}$ satisfies the above pentagonal equality and W satisfies $W_{23}W_{13}W_{12} = W_{12}W_{23}$. It is easy to see $V_\ell = \Sigma(\hat{J} \otimes \hat{J})V^*(\hat{J} \otimes \hat{J})\Sigma = (1 \otimes U)\Sigma V \Sigma(1 \otimes U)$. We define a left G-action on H_π by $\xi_i^\pi \mapsto \sum_{j \in I_\pi} w(\pi)_{i,j} \otimes \xi_j^\pi$ and denote it by H_π^ℓ. Let Θ_π be a unitary map $H_\pi^\ell \otimes H_\pi \longrightarrow L^2(G)$ defined by $\Theta_\pi(\xi_i \otimes \xi_j) = \sqrt{D_\pi(F_\pi)_{i,i}}w(\pi)_{i,j}$. It intertwines the left and right G-actions. Then we have the *Peter-Weyl* decomposition $\Theta : \oplus_{\pi \in \widehat{G}} H_\pi^\ell \otimes H_\pi \longrightarrow L^2(G)$ with $\Theta = \oplus_{\pi \in \widehat{G}} \Theta_\pi$. The *left, right reduced group C^*-algebras* are defined by

$$C_\ell^*(G) = \overline{\operatorname{span}}\{(\omega \otimes \mathrm{id})(V_\ell) \mid \omega \in \mathbb{B}(L^2(G))_*\}\,,$$
$$C_r^*(G) = \overline{\operatorname{span}}\{(\mathrm{id} \otimes \omega)(V) \mid \omega \in \mathbb{B}(L^2(G))_*\}\,,$$

where the closure is taken with respect to the operator norm of $\mathbb{B}(L^2(G))$. Note that $C_r^*(G)$ is contained in the commutant algebra of $C_\ell^*(G) = \hat{J}C_r^*(G)\hat{J}$. There is a distinguished projection $p_0 = (\mathrm{id} \otimes h)(V)$ in $C_r^*(G)$. It is also included in $C_\ell^*(G)$ and hence it is a central projection of $C_r^*(G)$. Moreover it is a minimal projection of $\mathbb{K}(L^2(G))$, in fact, we have $p_0 L^2(G) = \mathbb{C}\hat{1}_h$. We now have a map $\rho : \mathbb{B}(L^2(G))_* \longrightarrow C_r^*(G)$ by $\rho(\omega) = (\mathrm{id} \otimes \omega)(V)$. Let us define a normal functional $\theta(\pi)_{i,j}(x) = D_\pi(F_\pi)_{j,j}^{-1} h(xw(\pi)_{i,j}^*)$ for $x \in \mathbb{B}(L^2(G))$. Then the linear space $\operatorname{span}\{\rho(\theta(\pi)_{i,j}) \mid i, j \in I_\pi, \pi \in \widehat{G}\}$ is dense in $C_r^*(G)$ and moreover we have $\rho(\theta(\pi)_{i,j})\rho(\theta(\sigma)_{p,q}) = \delta_{\pi,\sigma}\delta_{j,p}\rho(\theta(\pi)_{i,q})$. Hence we obtain a $*$-isomorphism:

$$C_r^*(G) \longrightarrow \bigoplus_{\pi \in \widehat{G}} \mathrm{End}(H_\pi)$$

defined by $\rho(\theta(\pi)_{i,j}) \mapsto E_{i,j}^\pi$ which is a matrix unit. With this identification, the action of $E_{i,j}^\pi$ on $L^2(G)$ is given by

$$E_{i,j}^\pi w(\sigma)_{p,q}\hat{1}_h = \delta_{\pi,\sigma}\delta_{j,q}w(\pi)_{p,i}\hat{1}_h\,.$$

Using this fact and the Peter-Weyl decomposition, we see that the C^*-algebras $C_\ell^*(G)$ and $C_r^*(G)$ act on the first and the second tensor components of the Hilbert space $\oplus_{\pi \in \widehat{G}} H_\pi^\ell \otimes H_\pi$ respectively. This is just the differential representation of $\mathbb{B}(L^2(G))_* \subset A(G)^*$. Notice that $E_{i,j}^\pi$ is an operator of rank n and hence

$C_\ell^*(G)$ and $C_r^*(G)$ are contained in $\mathbb{K}(L^2(G))$. The projection $*$-homomorphism $C_r^*(G) \longrightarrow \mathrm{End}(H_\pi)$ is denoted by pr_π.

The *dual coproduct* is defined by $\hat{\delta} : C_r^*(G) \longrightarrow M(C_r^*(G) \otimes C_r^*(G))$ by

$$\hat{\delta}(x) = V^*(1 \otimes x)V \quad \text{for all } x \in C_r^*(G)$$

and it follows that linear subspaces $\hat{\delta}(C_r^*(G))(\mathbb{C} \otimes C_r^*(G))$ and $\hat{\delta}(C_r^*(G))(C_r^*(G) \otimes \mathbb{C})$ are dense in $C_r^*(G) \otimes C_r^*(G)$. On $C_\ell^*(G)$, we define $\widehat{\delta}_\ell(x) = \Sigma V_\ell^*(x \otimes 1)V_\ell \Sigma$.

Let $R(G)$ be a $\mathbb{Z}$-module $\bigoplus_{\pi \in \widehat{G}} \mathbb{Z}\pi$. For two $\pi, \sigma \in \widehat{G}$, we define their product via irreducible decomposition of the tensor product representation $w(\pi) \dot{\otimes} w(\sigma)$. More precisely, let $N_\tau^{\pi\,\sigma}$ be the dimension of the *hom-space* $\mathrm{Hom}(w(\tau), w(\pi)\dot{\otimes} w(\sigma))$, which does not depend on the choice of representatives of $\{w(\pi)\}_{\pi \in \widehat{G}}$. Then we have

$$\pi \cdot \sigma = \sum_{\tau \in \widehat{G}} N_\tau^{\pi\,\sigma} \tau.$$

The module $R(G)$ has an involution, that is, $\overline{\pi}$ is an equivalence class of $\overline{w(\pi)}$, which also does not depend on the choice of representatives of $\{w(\pi)\}_{\pi \in \widehat{G}}$. By these operations, $R(G)$ has an involutive $\mathbb{Z}$-ring structure and it is called the *representation ring* of G. We define the positive cone $R(G)_+ = \bigoplus_{\pi \in \widehat{G}} \mathbb{Z}_{\geq 0}\pi$.

Since a hom-space $\mathrm{Hom}(w(\tau), w(\pi)\dot{\otimes} w(\sigma))$ is naturally isomorphic to a hom-space $\mathrm{Hom}(w(\pi), w(\tau)\dot{\otimes}\overline{w(\sigma)})$ or $\mathrm{Hom}(w(\sigma), \overline{w(\pi)}\dot{\otimes} w(\tau))$, we have symmetry of $N_\tau^{\pi\,\sigma} = N_\pi^{\tau\,\bar{\sigma}} = N_\sigma^{\bar{\pi}\,\tau}$.

From now on, we use the letter A not for a continuous function algebra on a quantum group but for an arbitrary C^*-algebra. We give a brief summary of compact quantum group actions. The following definition is standard.

Definition 2.2. Let A be a (not necessarily unital) C^*-algebra and $(C(G), \delta)$ be a compact quantum group and $\alpha : A \longrightarrow A \otimes C(G)$ be a $*$-homomorphism. The triple $\{A, G, \alpha\}$ is called compact quantum covariant system (or simply covariant system) if the following statements hold:

(1) The map α is injective and satisfies $(\alpha \otimes \mathrm{id}) \circ \alpha = (\mathrm{id} \otimes \delta) \circ \alpha$.
(2) The vector space $\alpha(A)(\mathbb{C} \otimes C(G))$ is dense in $A \otimes C(G)$.

If we consider a compact quantum covariant system $\{A, G, \alpha\}$ where A is unital, we always assume that α is a unital $*$-homomorphism. Let $\{A, G, \alpha\}$ be a compact quantum covariant system. We can select representatives $\{w(\pi)\}_{\pi \in \widehat{G}}$ of $\widehat{G}$ whose F-matrices are diagonal and fix them in this section. We give comodule structure $\Gamma_\pi : H_\pi \longrightarrow H_\pi \otimes A(G)$ to a representation space H_π by fixing an orthonormal basis $\{\xi_j^\pi\}_{j \in I_\pi}$: $\Gamma_\pi(\xi_j^\pi) = \sum_{k \in I_\pi} \xi_k^\pi \otimes w(\pi)_{k,j}$ for any $j \in I_\pi$. For $\pi \in \widehat{G}$, define the functional θ_π on $C(G)$ by $\theta_\pi(x) = D_\pi h(x(\sum_{i \in I_\pi} (F_\pi^{-1})_{i,i} w(\pi)_{i,i}))$

for any $x \in C(G)$. As in the case of a compact group [S, Theorem 2], A is completely decomposable. Define the linear map $P_\pi = (\mathrm{id} \otimes \theta_\pi) \circ \alpha$ on A. From simple calculations, we have $P_\pi P_\rho = \delta_{\pi,\rho} P_\pi$. Put A_π for the range space of the projection P_π. Then the linear subspace $\mathcal{A} = \bigoplus_{\pi \in \widehat{G}} A_\pi$ is norm dense in A. Let $\mathrm{Hom}_G(H_\pi, A)$ be a set of intertwiners, that is, $\mathrm{Hom}_G(H_\pi, A) \ni S$ if and only if $\alpha \circ S = (S \otimes \mathrm{id}) \circ \Gamma_\pi$. Then we have $A_\pi = \overline{\mathrm{span}}\{S\xi \mid \xi \in H_\pi, S \in \mathrm{Hom}_G(H_\pi, A)\}$. The dimension of $\mathrm{Hom}_G(H_\pi, A)$ is called the *multiplicity* of π. Especially for the trivial representation π_0, the closed linear subspace A_{π_0} coincides with the fixed point algebra $A^\alpha = \{x \in A \mid \alpha(x) = x \otimes 1\}$. Let $m(\pi)$ be the multiplicity of π. We often use a formal symbol $\oplus_{\pi \in \widehat{G}} m(\pi)\pi$ which is called the *spectral pattern* of $\{A, G, \alpha\}$ (or simply A).

Next we recall the notion of the *crossed product*. Let $\{A, G, \alpha\}$ be a covariant system. Consider the Hilbert A-module $A \otimes L^2(G)$. In a C^*-algebra $\mathbb{K}(A \otimes L^2(G)) = A \otimes \mathbb{K}(L^2(G))$ define the crossed product C^*-algebra:

$$A \rtimes_\alpha G = \overline{\alpha(A)(\mathbb{C} \otimes C_r^*(G))}\,,$$

where the closure of linear space is taken with respect to the operator norm. It is also characterized as a fixed point algebra of $\{A \otimes \mathbb{K}(L^2(G)), G, \widetilde{\alpha}\}$: $A \rtimes_\alpha G = (A \otimes \mathbb{K}(L^2(G)))^{\widetilde{\alpha}}$, where $\widetilde{\alpha}(a \otimes k) = W_{23}\alpha(a)(1 \otimes k \otimes 1)W_{23}^*$ for all $a \in A$ and $k \in \mathbb{K}(L^2(G))$. Note that $\alpha(A)$, $C_r{}^*(G)$ are C^*-subalgebras of $M(A \rtimes_\alpha G)$ naturally. When we consider $A = C(G)$ and $\alpha = \delta$, we have a natural isomorphism:

$$C(G) \rtimes_\delta G \longrightarrow \mathbb{K}(L^2(G))$$

defined by $\delta(x)(1 \otimes y) \mapsto xy$. The *dual coaction* $\hat{\alpha} : A \rtimes_\alpha G \longrightarrow M(A \rtimes_\alpha G \otimes C_r^*(G))$ is defined by

$$\hat{\alpha}(x) = \widetilde{V}_{23}(x \otimes 1)\widetilde{V}_{23}^*$$

which satisfies the density condition that a linear space $\hat{\alpha}(A \rtimes_\alpha G)(\mathbb{C} \otimes C_r^*(G))$ is dense in $A \rtimes_\alpha G \otimes C_r^*(G)$. We define a $*$-homomorphism $\hat{\alpha}_\pi : A \rtimes_\alpha G \longrightarrow A \rtimes_\alpha G \otimes \mathrm{End}(H_\pi)$ by the composition $(\mathrm{id} \otimes \mathrm{pr}_\pi) \circ \hat{\alpha}$. Now we introduce the famous duality theorem. Let $\{A, G, \alpha\}$ be a covariant system. We consider the $*$-homomorphism $\hat{\alpha}^o = (\mathrm{id} \otimes \mathrm{Ad}\, U) \circ \hat{\alpha} : A \rtimes_\alpha G \longrightarrow M(A \rtimes_\alpha G \otimes C_\ell^*(G))$. It satisfies $(\mathrm{id} \otimes \widehat{\delta_\ell}^{\mathrm{op}}) \circ \hat{\alpha}^o = (\hat{\alpha}^o \otimes \mathrm{id}) \circ \hat{\alpha}^o$, that is, $\hat{\alpha}^o$ is a coaction of the opposite discrete quantum group $C_\ell^*(G)$. Define the crossed product $A \rtimes_\alpha G \rtimes_{\hat{\alpha}^o} (\widehat{G})'^{\mathrm{op}}$ by the norm closure of the linear space $\hat{\alpha}^o(A \rtimes_\alpha G)(\mathbb{C} \otimes \mathbb{C} \otimes C(G))$. Its dual action $\widehat{\hat{\alpha}^o}$ is given by $\widehat{\hat{\alpha}^o}(x) = V_{34}(x \otimes 1)V_{34}^*$. The duality theorem says there exists an isomorphism between $A \rtimes_\alpha G \rtimes_{\alpha^o} (\widehat{G})'^{\mathrm{op}}$ and $A \otimes \mathbb{K}(L^2(G))$ such that $\hat{\alpha}^o\big(\alpha(a)(1 \otimes x)\big)(1 \otimes 1 \otimes y)$ is mapped to $\alpha(a)(1 \otimes x)(1 \otimes UyU)$ for all $a \in A$, $x \in C_r^*(G)$ and $y \in C(G)$. This isomorphism conjugates $\widehat{\hat{\alpha}^o}$ and $\widetilde{\alpha}$.

Let $G = (C(G), \delta_G)$ and $H = (C(H), \delta_H)$ be compact quantum groups. We say H is a *quantum subgroup* (or sometimes simply called a subgroup) if there exists a surjective $*$-homomorphism $r : C(G) \longrightarrow C(H)$ such that r satisfies $\delta_H \circ r = (r \otimes r) \circ \delta_G$. This surjection is often called the restriction homomorphism.

In general a choice of r is not unique. The *(left) quotient space* is defined by

$$C(H \setminus G) = \{x \in C(G) \mid (r \otimes \mathrm{id}) \circ \delta_G(x) = 1 \otimes x\}\,.$$

Actually it depends on r and we shall denote it by ${}^{H}C(G)$, however, we abuse the notation of $H \setminus G$. We have to be careful of r when we consider a quotient space.

3. Ergodic systems

If a compact quantum covariant system $\{A, G, \alpha\}$ with a unital C^*-algebra A satisfies $A^\alpha = \mathbb{C}1$, then we call it a *compact quantum ergodic system* (or simply an ergodic system). We investigate the ergodic system in this section. The faithful conditional expectation $\varphi = (\mathrm{id} \otimes h) \circ \alpha$ onto $A^\alpha = \mathbb{C}1$ is the unique invariant state in this case, where invariance means: $(\varphi \otimes \mathrm{id})(\alpha(x)) = \varphi(x)1$ for any $x \in A$. Note that φ is a trace if G is a compact group [H-L-S, Theorem 4.1]. In [B, Theorem 17], it has been proved that the dimension of A_π is less than or equal to D_π^2 for any $\pi \in \widehat{G}$. We will make a sharper estimate of these dimensions. In order to do this, we have to explain the existence of the modular automorphism group with respect to the state φ. By [B, Proposition 18], there exists the unital multiplicative linear map $\Theta : \mathcal{A} \longrightarrow \mathcal{A}$ such that $\varphi(ab) = \varphi(\Theta(b)a)$ for any $a, b \in \mathcal{A}$. Let M be the von Neumann algebra $\pi_\varphi(A)''$ associated to the state φ via GNS-representation $\{H_\varphi, \pi_\varphi, \xi_\varphi\}$ and the extension of φ to M is also denoted by φ. We often identify A with $\pi_\varphi(A)$. Let $p \in M$ be a projection with $\varphi(p) = 0$. Then for any $a \in \mathcal{A}$ we have $\varphi(a^*pa) = \varphi(\Theta(a)a^*p)$. This is in fact equal to 0 by the Cauchy-Schwarz inequality. From this we see $p\mathcal{A}\xi_\varphi = 0$ and $p = 0$, because the linear subspace $\mathcal{A}\xi_\varphi$ is dense in H_φ. Hence φ is a faithful normal state on M and there exists the modular automorphism group $\{\sigma^\varphi\}_{t\in\mathbb{R}}$ on M. Since φ is an invariant state on A for the action of G, the action α extends to the action on M. An action of a compact quantum group on a von Neumann algebra is defined similarly to the case of a C^*-algebra by using the σ-weak topology. Note the following useful equality about the modular automorphism group and the scaling automorphism group:

$$\alpha \circ \sigma_t^\varphi = (\sigma_t^\varphi \otimes \tau_{-t}) \circ \alpha \text{ for all } t \in \mathbb{R}.$$

For its proof, readers are referred to [E, Théorème 2.9]. The spectral subspace M_π is the σ-weak closure of A_π, however, they coincide because of the finite dimensionality of A_π.

Lemma 3.1. *Let S be an intertwiner of H_π and A. For any $t \in \mathbb{R}$, we define the map $S_t : H_\pi \longrightarrow A$ by $S_t\xi_j^\pi = (F_\pi)_{j,j}^{-it}\sigma_t^\varphi(S\xi_j^\pi)$ for any $j \in I_\pi$. Then S_t is also an intertwiner of H_π and A for any $t \in \mathbb{R}$, in particular we have $\sigma_t^\varphi(A_\pi) = A_\pi$ for any $t \in \mathbb{R}$ and we see any element of A_π is analytic for $\{\sigma_t^\varphi\}_{t\in\mathbb{R}}$.*

Proof. We shall show that $\alpha(S_t\xi_j^\pi) = \sum_{k\in I_\pi} S_t\xi_k^\pi \otimes w(\pi)_{k,j}$ for any $j \in I_\pi$ and $t \in \mathbb{R}$. In fact, we have

$$\begin{aligned}\alpha(S_t\xi_j^\pi) =&(F_\pi)_{j,j}^{-it}\alpha(\sigma_t^\varphi(S\xi_j^\pi))\\ =&(F_\pi)_{j,j}^{-it}(\sigma_t^\varphi \otimes \tau_{-t})(\alpha(S\xi_j^\pi)))\\ =&(F_\pi)_{j,j}^{-it}\sum_{k\in I_\pi}\sigma_t^\varphi(S\xi_k^\pi)\otimes \tau_{-t}(w(\pi)_{k,j})\\ =&(F_\pi)_{j,j}^{-it}\sum_{k\in I_\pi}\sigma_t^\varphi(S\xi_k^\pi)\otimes (F_\pi)_{k,k}^{-it}(F_\pi)_{j,j}^{it}w(\pi)_{k,j}\\ =&\sum_{k\in I_\pi} S_t\xi_k^\pi \otimes w(\pi)_{k,j}.\end{aligned}$$

□

From this lemma we see that $\sigma_t^\varphi(A) = A$ for $t \in \mathbb{R}$ and the following proposition holds.

Proposition 3.2. *Let $\{A, G, \alpha\}$ be an ergodic system and φ be the invariant state on A. Then there exists the modular automorphism group $\{\sigma_t^\varphi\}_{t\in\mathbb{R}}$ on A.*

Definition 3.3. Let $\{A, G, \alpha\}$ be an ergodic system and π be an element of $\widehat{G}$. A vector $\xi = (\xi_j)_{j\in I_\pi} \in \bigoplus_{j\in I_\pi} A_\pi$ is called a π-eigenvector if $\alpha(\xi_j) = \sum_{k\in I_\pi}\xi_k \otimes w(\pi)_{k,j}$ for any $j \in I_\pi$. The set of π-eigenvectors is called the π-eigenvector space and denoted by $X_{w(\pi)}$.

Let $\{A, G, \alpha\}$ and $\{\xi^\lambda\}_{\lambda\in\Lambda}$ be a covariant system and a set of its π_λ-eigenvectors, respectively. A C^*-subalgebra generated by ξ_r^λ for all $r \in I_{\pi_\lambda}$ and $\lambda \in \Lambda$ is denoted by $C^*(\{\xi^\lambda\}_{\lambda\in\Lambda})$. The quantum group G acts on it invariantly. We say that it is a G-invariant C^*-subalgebra generated by $\{\xi^\lambda\}_{\lambda\in\Lambda}$. We make a π-eigenvector space to a Hilbert space by defining its inner product with

$$(\xi \mid \eta) = \sum_{k\in I_\pi}\xi_k\eta_k^*, \text{ for all } \xi, \eta \in X_{w(\pi)}.$$

In fact, the above right hand side is an element of the fixed point algebra and therefore a scalar by the ergodicity of the action.

We remark eigenvector spaces do not essentially depend on the selection of unitaries $\{w(\pi)\}_{\pi\in\widehat{G}}$. In fact, let us take two selections from π: $w(\pi)$, $w'(\pi)$ and let $X_{w(\pi)}$ and $X_{w'(\pi)}$ be the π-eigenvector spaces corresponding $w(\pi)$ and $w'(\pi)$, respectively. Then there exists a unitary $u : H_{w(\pi)} \longrightarrow H_{w'(\pi)}$ satisfying $uw(\pi) = w'(\pi)u$. The unitary u gives a unitary isomorphism between $X_{w(\pi)}$ and $X_{w'(\pi)}$ by sending $\xi \in X_{w(\pi)}$ to $\bar{u}\xi \in X_{w(\pi)}$, where $(\bar{u}\xi)_k = \sum_{\ell\in I_\pi}\overline{u_{k,\ell}}\xi_\ell$.

Definition 3.4. Let $\{A, G, \alpha\}$ be an ergodic system and π be an element of $\widehat{G}$. We define the following two operations on eigenvector spaces.

(1) For any $t \in \mathbb{R}$, we define the linear map $U_t^{w(\pi)} : X_{w(\pi)} \longrightarrow X_{w(\pi)}$ by

$$(U_t^{w(\pi)}\xi)_k = (F_\pi)_{k,k}^{-it}\sigma_t^\varphi(\xi_k) \quad \text{for all} \quad \xi \in X_\pi, k \in I_\pi.$$

(2) We define the conjugate linear map $T_{w(\pi)} : X_{w(\pi)} \longrightarrow X_{\overline{w(\pi)}}$.

$$(T_{w(\pi)}\xi)_k = (F_\pi)_{k,k}^{-\frac{1}{2}}\xi_k^* \quad \text{for all} \quad \xi \in X_\pi, k \in I_\pi.$$

The well-definedness of the above first operation has been already checked in Lemma 3.1. For the second one we shall justify the equality $\alpha((T_{w(\pi)}\xi)_k) = \sum_{\ell \in I_\pi}(T_{w(\pi)}\xi)_\ell \otimes \overline{w(\pi)}_{\ell,k}$ for any $\xi \in X_{w(\pi)}$ and $k \in I_\pi$. In fact,

$$\begin{aligned}
\alpha((T_{w(\pi)}\xi)_k) =& (F_\pi)_{k,k}^{-\frac{1}{2}}\alpha(\xi_k^*) \\
=& (F_\pi)_{k,k}^{-\frac{1}{2}}\sum_{\ell \in I_\pi}\xi_\ell^* \otimes w(\pi)_{\ell,k}^* \\
=& (F_\pi)_{k,k}^{-\frac{1}{2}}\sum_{\ell \in I_\pi}\xi_\ell^* \otimes (F_\pi)_{k,k}^{\frac{1}{2}}(F_\pi)_{\ell,\ell}^{-\frac{1}{2}}\overline{w(\pi)}_{\ell,k} \\
=& \sum_{\ell \in I_\pi}(T_{w(\pi)}\xi)_\ell \otimes \overline{w(\pi)}_{\ell,k}.
\end{aligned}$$

Proposition 3.5. *Let $\{A, G, \alpha\}$ be an ergodic system and π be an element of $\widehat{G}$. Then the map* $\mathrm{Hom}_G(H_\pi, A) \ni S \longrightarrow (S(\xi_k^\pi))_{k \in I_\pi} \in X_{w(\pi)}$ *is a linear isomorphism.*

Proof. For $\xi \in X_{w(\pi)}$, define the element $S \in \mathrm{Hom}_G(H_\pi, A)$ by $S(\xi_k^\pi) = \xi_k$ for any $k \in I_\pi$. It is easy to see that this map is well-defined and gives the desired inverse map. □

From this proposition we see the spectral subspace A_π is spanned by the entries of π-eigenvectors.

The next lemma has already appeared in the proof of [B, Proposition 18], however, we give a proof for readers' convenience.

Lemma 3.6. *Let ξ and η be π-eigenvectors of an ergodic system $\{A, G, \alpha\}$. Then the following equalities hold*

(1) $\varphi(\xi_k\eta_\ell^*) = D_\pi^{-1}(F_\pi)_{k,k}\delta_{k,\ell}(\xi \mid \eta)$, *for all* $k, \ell \in I_\pi$,
(2) $\varphi(\xi_k^*\eta_\ell) = 0$, *if* $k \neq \ell \in I_\pi$, *and* $\varphi(\xi_k^*\eta_k) = \varphi(\xi_\ell^*\eta_\ell)$, *for all* $k, \ell \in I_\pi$.

Proof. (1)It is a straightforward calculation

$$\begin{aligned}\varphi(\xi_k\eta_\ell^*) &= \sum_{r,s\in I_\pi} \xi_r\eta_s^* h(w(\pi)_{r,k}w(\pi)_{s,\ell}^*)\\ &= \sum_{r,s\in I_\pi} \xi_r\eta_s^* D_\pi^{-1}(F_\pi)_{k,k}\delta_{r,s}\delta_{k,\ell}\\ &= D_\pi^{-1}(F_\pi)_{k,k}\delta_{k,\ell}\sum_{r\in I_\pi}\xi_r\eta_r^*\\ &= D_\pi^{-1}(F_\pi)_{k,k}\delta_{k,\ell}(\xi \mid \eta).\end{aligned}$$

(2) Let $Z = (\varphi(\xi_k^*\eta_\ell))_{k,\ell\in I_\pi}$ be a matrix on the representation space H_π. Then we have

$$\begin{aligned}(w(\pi)^*Zw(\pi))_{i,j} &= \sum_{k,\ell\in I_\pi} w(\pi)_{k,i}^* Z_{k,\ell} w(\pi)_{\ell,j}\\ &= \sum_{k,\ell\in I_\pi} w(\pi)_{k,i}^*\varphi(\xi_k^*\eta_\ell)w(\pi)_{\ell,j}\\ &= \sum_{k,\ell\in I_\pi} (\varphi\otimes \mathrm{id})((\xi_k^*\otimes w(\pi)_{k,i}^*)(\eta_\ell\otimes w(\pi)_{\ell,j}))\\ &= (\varphi\otimes\mathrm{id})\alpha(\xi_i^*\eta_j)\\ &= \varphi(\xi_i^*\eta_j)\\ &= Z_{i,j}.\end{aligned}$$

Therefore the operator Z commutes with the irreducible unitary representation $w(\pi)$, so we have $Z \in \mathbb{C}$. □

Lemma 3.7. *Let $\{A, G, \alpha\}$ be an ergodic system and π be an element of $\widehat{G}$. Then it follows $T_{w(\pi)}^* T_{w(\pi)} = U_i^{w(\pi)}$.*

Proof. Take π-eigenvectors ξ and η. Then we have

$$\begin{aligned}(T_{w(\pi)}\xi \mid T_{w(\pi)}\eta) &= \sum_{k\in I_\pi}(T_{w(\pi)}\xi)_k(T_{w(\pi)}\eta)_k^*\\ &= \sum_{k\in I_\pi}(F_\pi)_{k,k}^{-\frac{1}{2}}(F_\pi)_{k,k}^{-\frac{1}{2}}\xi_k^*\eta_k\\ &= \sum_{k\in I_\pi}(F_\pi)_{k,k}^{-1}\varphi(\xi_k^*\eta_k)\\ &= \sum_{k\in I_\pi}(F_\pi)_{k,k}^{-1}\varphi(\sigma_i^\varphi(\eta_k)\xi_k^*)\\ &= \sum_{k\in I_\pi}(F_\pi)_{k,k}^{-1}\varphi((F_\pi)_{k,k}^{-1}(U_i^{w(\pi)}\eta)_k\xi_k^*)\\ &= \sum_{k\in I_\pi}(F_\pi)_{k,k}^{-2}\cdot D_\pi^{-1}(F_\pi)_{k,k}(U_i^{w(\pi)}\eta \mid \xi)\end{aligned}$$

$$=(U_i^{w(\pi)}\eta \mid \xi),$$

where we use the result of the previous lemma and $\sum_{k\in I_\pi}(F_\pi^{-1})_{k,k} = D_\pi$. Hence we obtain the desired equality. $\square$

Let $T_{w(\pi)} = J_{w(\pi)}|T_{w(\pi)}|$ be the polar decomposition of the conjugate linear map $T_{w(\pi)} : X_{w(\pi)} \longrightarrow X_{\overline{w(\pi)}}$. Since it is easy to see $T_{\overline{w(\pi)}} \circ T_{w(\pi)} = \mathrm{id}_{X_{w(\pi)}}$, the map $J_{w(\pi)}$ is a conjugate unitary map and satisfies

$$J_{\overline{w(\pi)}} = J_{w(\pi)}^{-1}, \text{ and } J_{\overline{w(\pi)}}|T_{\overline{w(\pi)}}|J_{w(\pi)} = \left|T_{w(\pi)}\right|^{-1}.$$

In particular we obtain $\mathrm{Sp}\left(U_{\frac{i}{2}}^{w(\pi)}\right) = \mathrm{Sp}\left(U_{\frac{i}{2}}^{\overline{w(\pi)}}\right)^{-1}$. With these preparations we show the following result about bounds of multiplicities.

Theorem 3.8. *Let $\{A, G, \alpha\}$ be an ergodic system and π be an element of $\widehat{G}$. Then we have*

$$\dim \mathrm{Hom}_G(H_\pi, A) \leq D_\pi.$$

Proof. The proof is essentially due to [B, Theorem 17] or [Wa1, Theorems 1, 2]. As we have proved, in Proposition 3.5, the hom-space $\mathrm{Hom}_G(H_\pi, A)$ is naturally isomorphic to $X_{w(\pi)}$. Hence it suffices to show $d := \dim(X_{w(\pi)}) \leq D_\pi$. The unitary $\mathbb{R}$-action $\{U_t^{w(\pi)}\}_{t\in\mathbb{R}}$ enables us to take an orthonormal basis of $X_{w(\pi)}$ $\{\xi^p\}_{1\leq p\leq d}$ as $U_t^{w(\pi)}\xi^p = \lambda_p^{it}\xi^p$ for all $t \in \mathbb{R}$ where λ_p is a positive real number. Then we define an operator entry matrix M by

$$M = \begin{pmatrix} \xi^1 \\ \xi^2 \\ \vdots \\ \xi^d \end{pmatrix} \in \mathbb{M}_{d,d_\pi}(A_\pi),$$

where each ξ^p is treated as a row vector. Adding 0 entries, we embed the matrix M to a larger square matrix (still denoted by M) of size n ($\geq d, d_\pi$), if necessary. By orthonormality, we have $MM^* = 1_d$. Let $k, \tilde{F}$ be positive matrices $\mathrm{diag}(\lambda_1^{-1}, \lambda_2^{-1}, \dots, \lambda_d^{-1}, 1, \dots, 1)$ and $\mathrm{diag}(F_{w(\pi)}, 1, \dots, 1)$ in $\mathbb{M}_n(\mathbb{C})_+$, respectively.

Then we have

$$\begin{aligned}\sum_{p=1}^{d}\lambda_p^{-1} &= \mathrm{Tr}_k(1_d)\\ &=(\varphi\otimes\mathrm{Tr}_k)(MM^*)\\ &=(\varphi\otimes\mathrm{Tr}_k)(M^*(\sigma_{-i}^{\varphi}\otimes\mathrm{Ad}(k))(M))\\ &=(\varphi\otimes\mathrm{Tr}_k)(M^*M\tilde{F}k^{-1})\\ &=(\varphi\otimes\mathrm{Tr})(M^*M\tilde{F})\\ &\leq(\varphi\otimes\mathrm{Tr})(1\otimes\tilde{F}1_{d_\pi})\\ &=D_\pi,\end{aligned}$$

where we use $(\sigma_t^{\varphi}\otimes\mathrm{Ad}(k^{it}))(M)=M\tilde{F}^{it}k^{-it}$ for $t\in\mathbb{R}$. Hence we obtain $\mathrm{Tr}_{X_{w(\pi)}}(U_i^{w(\pi)})\leq D_\pi$ and similarly $\mathrm{Tr}_{X_{\overline{w(\pi)}}}(U_i^{\overline{w(\pi)}})\leq D_{\overline{\pi}}=D_\pi$. Since we have already known $\mathrm{Sp}\left(U_i^{w(\pi)}\right)=\mathrm{Sp}\left(U_i^{\overline{w(\pi)}}\right)^{-1}$, it follows $2d\leq\mathrm{Tr}_{X_{w(\pi)}}(U_i^{w(\pi)})+\mathrm{Tr}_{X_{\overline{w(\pi)}}}(U_i^{\overline{w(\pi)}})$. Therefore we obtain $2d\leq 2D_\pi$. □

From this result, the dimension of the π-eigenspace A_π is less than or equal to $d_\pi D_\pi$. We can derive the quantum version of [Wa1, Theorem 2].

Theorem 3.9. *Let $\{A,G,\alpha\}$ be a compact quantum covariant system and π be an element of $\widehat{G}$. If p and q are minimal projections in A^α, then we have:*

$$\dim\mathrm{Hom}_G(H_\pi,pAq)\leq D_\pi.$$

Proof. This proof is also essentially due to [B, Theorem 17] or [Wa1, Theorems 1, 2], however, we cannot use the trace property as in the case of a compact group, we have to prove the finite dimensionality of $\mathrm{Hom}_G(H_\pi,pAq)$ at first. If the linear space $pA^\alpha q$ is not 0, we can take non-zero norm 1 element x from $pA^\alpha q$. Then we see $xx^*\in pA^\alpha p=\mathbb{C}p$ and $x^*x\in qA^\alpha q=\mathbb{C}q$ and have $xx^*=p$ and $x^*x=q$. It follows $(pAq)_\pi=(pAp)_\pi x$. Therefore the assertion of this theorem follows from applying the previous theorem to pAp. Next we assume $pA^\alpha q=0$. Because $p\perp q$, we may assume that A is unital and $1=p+q$ by considering $(p+q)A(p+q)$. Let $\varphi(x)=\varphi_p(pxp)+\varphi_q(qxq)$ where φ_p and φ_q are the invariant states on pAp and qAq respectively. Since the conditional expectation with respect to φ $E_\alpha:A\longrightarrow A^\alpha$ is given by $E_\alpha(x)=pxp+qxq$, φ is the invariant positive functional on A. Let $d\leq\infty$ be the dimension of the linear space $\mathrm{Hom}_G(H_\pi,A)$ and fix a natural number $0\leq d'\leq d$. Note that we can take π-eigenvectors $\{\xi^j\}_{1\leq j\leq d'}$ from $(pAq)_\pi=pA_\pi q$, which satisfy $\varphi((\xi_r^j)^*\xi_s^k)=\delta_{j,k}\delta_{r,s}$ for $1\leq j,k\leq d'$ and $r,s\in I_\pi$. In fact, if we have $\{\xi^j\}_{1\leq j\leq k-1}$ as before, we can take a π-eigenvector ξ^k which satisfies $\varphi((\xi_1^j)^*\xi_1^k)=\delta_{j,k}$ for $1\leq j\leq p-1$. Then modifying Lemma 3.6 to the case of pAq, we achieve to take desired vectors. Let

M be the matrix

$$M = \begin{pmatrix} T_{w(\pi)}\xi^1 \\ T_{w(\pi)}\xi^2 \\ \vdots \\ T_{w(\pi)}\xi^{d'} \end{pmatrix}.$$

Adding 0-entries, we embed M to a larger square matrix (still denoted by M) of size n ($\geq d', d_\pi$). Note that $T_{w(\pi)}\xi^j$ is a $\bar{\pi}$-eigenvector for qAp: $T_{w(\pi)}\xi^j = \left((F_\pi)_{r,r}^{-\frac{1}{2}}\xi_r^{j*}\right)_{r\in I_\pi}$. By the computation in the proof of Lemma 3.7, it follows $MM^* = D_\pi q \otimes 1_{d'}$. Hence we have $M^*M \leq D_\pi p \otimes 1_{d_\pi}$ in $pAp \otimes \mathbb{B}(H_\pi)$. Then it yields:

$$\begin{aligned} D_\pi^2 \geq& (\varphi \otimes \mathrm{Tr}_{F_\pi})(M^*M) \\ =& \sum_{r\in I_\pi} \left((F_\pi)_{r,r}(F_\pi^{-\frac{1}{2}})_{r,r}(F_\pi^{-\frac{1}{2}})_{r,r} \sum_{1\leq j\leq d'} \varphi(\xi_r^j \xi_r^{j*})\right) \\ =& \sum_{1\leq j\leq d'} (\xi^j \mid \xi^j), \end{aligned}$$

where $(\xi^j \mid \xi^j)$ means the inner product of π-eigenvectors for pAq. Let us define another inner product $(\cdot \mid \cdot)_2$ of π-eigenvectors for pAq by $(\xi \mid \eta)_2 = \varphi(\eta_r^*\xi_r)$ for any π-eigenvectors ξ, η. Note that this value does not depend on the choice of r by Lemma 3.6. Let W be a linear space spanned by $\{\xi^j\}_{1\leq j\leq d'}$. There are two inner products $(\cdot \mid \cdot)$ and $(\cdot \mid \cdot)_2$ on W. Let A_W be the matrix which satisfies $(A_W\xi \mid \eta)_2 = D_\pi^{-1}(\xi \mid \eta)$. Then from the above inequality we have $\mathrm{Tr}_W(A_W) \leq D_\pi$, where Tr_W is the non-normalized trace associated to W. Similarly considering $T_{w(\pi)}(W)$ for W, we also obtain $\mathrm{Tr}_{T_{w(\pi)}(W)}(A_{T_{w(\pi)}(W)}) \leq D_\pi$. Moreover, we see $A_W = T_{w(\pi)}^* T_{w(\pi)}$, where the involution $*$ comes from the conjugate linear map $T_{w(\pi)} : W \longrightarrow T_{w(\pi)}(W)$ between $(\cdot \mid \cdot)_2$-inner product spaces. In fact, we have

$$\begin{aligned} (T_{w(\pi)}\xi \mid T_{w(\pi)}\eta)_2 =& \varphi({(T\eta)_1}^*(T\xi)_1) \\ =& (F_\pi^{-1})_{1,1}\varphi(\eta_1\xi_1^*) \\ =& (F_\pi^{-1})_{1,1}D_\pi^{-1}(F_\pi)_{1,1}(\eta \mid \xi) \\ =& D_\pi^{-1}(\eta \mid \xi). \end{aligned}$$

Let us denote $T_{w(\pi)}(W)$ by $\overline{W}$. Let $T_{w(\pi)} = J_W|T_{w(\pi)}|$ and $T_{\overline{w(\pi)}} = J_{\overline{W}}|T_{\overline{w(\pi)}}|$ be the polar decomposition of the conjugate linear maps $T_{w(\pi)}$ and $T_{\overline{w(\pi)}}$ between W and $T_{w(\pi)}(W)$. The equality $T_{w(\pi)} \circ T_{\overline{w(\pi)}} = \mathrm{id}_{\overline{W}}$ yields $J_W A_W^{\frac{1}{2}} J_{\overline{W}} = A_{\overline{W}}^{-\frac{1}{2}}$. Then we have $J_W A_W J_W^* = A_{\overline{W}}^{-1}$ because J_W and $J_{\overline{W}}$ are conjugate unitary. Hence we obtain $\mathrm{Tr}(A_W) = \mathrm{Tr}(A_{\overline{W}}^{-1})$ and it follows $2d' \leq 2\,\mathrm{Tr}(A_W) \leq 2D_\pi$. □

Finally we state the modified version of Theorem 3.8 and Theorem 3.9.

Proposition 3.10. *Let $\{A, G, \alpha\}$ be an ergodic system and π be an element of $\widehat{G}$. Assume that A has a tracial state. Then we have:*

$$\dim \operatorname{Hom}(H_\pi, A) \leq d_\pi.$$

Proof. We have proved finite dimensionality of $X_{w(\pi)}$. Let d be its dimension and $\{\xi^p\}_{1\leq p\leq d}$ be an orthonormal basis of $X_{w(\pi)}$ for the inner product $(\cdot \mid \cdot)$. Take a tracial state τ on A. Define the matrix M by

$$M = \begin{pmatrix} \xi^1 \\ \xi^2 \\ \vdots \\ \xi^d \end{pmatrix}.$$

As the proof of Theorem 3.8, we look at M in a bigger square matrix algebra. Then we have $MM^* = 1_d$ and it yields

$$\begin{aligned} d_\pi \geq& (\tau \otimes \mathrm{Tr})(M^*M) \\ =&(\tau \otimes \mathrm{Tr})(MM^*) \\ =&d. \end{aligned}$$

□

Proposition 3.11. *Let $\{A, G, \alpha\}$ be a compact quantum covariant system and π be an element of $\widehat{G}$. Assume that A has a tracial state. If p and q are minimal projections in A^α and they are not annihilated by a trace on A, then we have*

$$\dim \operatorname{Hom}(H_\pi, pAq) \leq d_\pi.$$

Proof. Let $d(\leq D_\pi)$ be the dimension of $\dim \operatorname{Hom}(H_\pi, pAq)$. Let $\{\xi^p\}_{1\leq p\leq d}$ be an orthonormal π-eigenvector for pAq where we define the inner product by

$$(\xi \mid \eta)p = \sum_{r\in I_\pi} \xi_r \eta_r^*$$

for all π-eigenvector ξ and η. Take a tracial state τ on A with $\tau(p)\tau(q) \neq 0$. Define the matrix M by

$$M = \begin{pmatrix} \xi^1 \\ \xi^2 \\ \vdots \\ \xi^d \end{pmatrix}.$$

and embed it into the larger square matrix as in previous theorems. Then we have $MM^* = p \otimes 1_d$. This yields

$$\begin{aligned} d\,\tau(p) =&(\tau \otimes \mathrm{Tr})(MM^*) \\ =&(\tau \otimes \mathrm{Tr})(M^*M) \\ \leq&(\tau \otimes \mathrm{Tr})(q \otimes 1_{d_\pi}) \\ =&\tau(q)\, d_\pi. \end{aligned}$$

Hence we have $d \leq d_\pi \frac{\tau(q)}{\tau(p)}$. Considering $qAp = (pAq)^*$ and $d = \dim \mathrm{Hom}(H_\pi, qAp)$, we also have $d \leq d_{\overline{\pi}} \frac{\tau(p)}{\tau(q)} = d_\pi \frac{\tau(p)}{\tau(q)}$. Finally we obtain $d \leq d_\pi \min\left(\frac{\tau(q)}{\tau(p)}, \frac{\tau(p)}{\tau(q)}\right) \leq d_\pi$.

$\square$

Remark 3.12. Assume that the linear space pAq is not 0. Then $\tau(p) \neq 0$ if and only if $\tau(q) \neq 0$. Indeed, take $\pi \in \widehat{G}$ as $pA_\pi q \neq 0$ and assume $\tau(q) = 0$, then we obtain $\tau(p) = 0$ by the proof of the above proposition. The converse assertion holds by applying the involution $*$ from pAq to qAp.

4. Equivariant K-theory

We follow [Wa1] and use equivariant K-theory to obtain a multiplicity map. We briefly recall the notions. Readers are referred to, for example, [BS1] and [V] for the theory of equivariant K-theory. Let $\{A, G, \alpha\}$ be a compact quantum covariant system with A unital. Let E be a finitely generated projective Hilbert A-module and $\delta_E : E \longrightarrow E \otimes C(G)$ be a linear map. The pair $\{E, \delta_E\}$ is called a G-equivariant A-module if they satisfy the following

1. $(\mathrm{id} \otimes \delta) \circ \delta_E = (\delta_E \otimes \mathrm{id}) \circ \delta_E$,
2. $\delta_E(ea) = \delta_E(e)\alpha(a)$ for all $e \in E$ and $a \in a$,
3. $\langle \delta_E(e) \mid \delta_E(e') \rangle = \alpha(\langle e \mid e' \rangle)$ for all $e, e' \in E$,
4. the linear subspace $\delta_E(E)(\mathbb{C} \otimes C(G))$ is dense in $E \otimes C(G)$.

Actually the second equality follows from the third one. Inner products of Hilbert modules are conjugate-linear for the first variable. Two G-equivariant A-modules are equivalent if there exists a bijective linear map intertwining of the actions of G and A. The set of these equivalence classes becomes an abelian semigroup by the direct sum and its Grothendieck group is denoted by $K_0^G(A)$ and this is called an *equivariant K-group*. For a G-equivariant A-module $\{E, \delta_E\}$ we define its crossed product Hilbert $A \rtimes_\alpha G$-module by $E \rtimes_\alpha G = E \otimes_A (A \rtimes_\alpha G)$. This module becomes a (left) $C_r^*(G)$-module by an appropriate way ([V, Lemme 5.2]). Here we take a non-rigorous picture for its action. Let x be an element of $C_r^*(G)$ and we assume its dual coproduct has an expansion $\hat{\delta}(x) = \sum x_{(0)} \otimes x_{(1)}$. It is considered that $C_r^*(G)$ acts on E through the differential representation with respect to δ_E. Hence we have an action $x \cdot (e \otimes_A y) = \sum x_{(0)}(e) \otimes_A x_{(1)} y$. This action is compatible with the right action of A, because in $A \rtimes_\alpha G$ we have $z(\alpha(y)) = \sum (z_{(0)} \cdot y) z_{(1)}$ for $z \in C_r^*(G)$ and $y \in A$ where $C_r^*(G)$ acts on A with the differential representation about α. We next recall the Green-Julg isomorphism. If E_π is a tensor product module $H_\pi \otimes A$ with a finite dimensional irreducible G-module H_π, then we have $E_\pi \otimes_A (A \rtimes_\alpha G) = H_\pi \otimes (A \rtimes_\alpha G)$. We apply $p_0 \in C_r^*(G)$ to this space. Since this action is obtained by the dual coproduct, we have $p_0(E_\pi \otimes_A A \rtimes_\alpha G) = p_0(H_\pi \otimes C_r^*(G)) \otimes_{C_r^*(G)} A \rtimes_\alpha G \cong p_{\overline{\pi}} C_r^*(G) \otimes_{C_r^*(G)} A \rtimes_\alpha G = (1 \otimes p_{\overline{\pi}}) A \rtimes_\alpha G$. In general case E is a direct summand of a direct sum of E_π. Hence the module $p_0(E \rtimes_\alpha G)$ corresponds to a projection in $M_n(A \rtimes_\alpha G)$ for some n. We also notice that $A \rtimes_\alpha G$ is stably unital. In fact we have an approximate unit of $\mathbb{K}(L^2(G))$ which consists of projections in $C^*_r(G)$. In this way we obtain the Green-Julg isomorphism (see [V, Théorème 5.10] for its proof of KK-version):

$$\Phi_1 : K_0^G(A) \longrightarrow K_0(A \rtimes_\alpha G)$$

defined by $[E] \mapsto [p_0(E \rtimes_\alpha G)]$. The inverse map of Φ_1 is given by

$$\Phi_2 : K_0(A \rtimes_\alpha G) \longrightarrow K_0^G(A)$$

which sends $[q]$ to $[q(A \otimes L^2(G))^n]$, where $q \in M_n(A \rtimes_\alpha G)$ is a projection and $A \otimes L^2(G)$ is a G-equivariant A-module with $\delta_{A\otimes L^2(G)}(a \otimes \xi) = (1 \otimes W(\xi \otimes 1))\alpha(a)_{13}$. We define the usual $R(G)$-module structures on $K_0^G(A)$ and $K_0(A \rtimes_\alpha G)$. Let π be an element of $\widehat{G}$ and E be a G-equivariant A-module and $q \in M_n(A \rtimes_\alpha G)$ be a projection. We define the action of π

$$\begin{aligned} \pi \cdot [E] &= [H_\pi \otimes E]\,, \\ \pi \cdot [q] &= [(\mathrm{id}_{M_n(\mathbb{C})} \otimes \hat{\alpha}_\pi)(q)]\,, \end{aligned}$$

where we use the isomorphism $K_0(A \rtimes_\alpha G) \cong K_0(A \rtimes_\alpha G \otimes \mathrm{End}(H_\pi))$. We observe the Green-Julg isomorphism is an $R(G)$-module map in the compact group case.

Lemma 4.1. *Let B be a C^*-algebra and E be a Hilbert B-module which has a $C^*_r(G)$-action $\phi : C^*_r(G) \longrightarrow \mathbb{B}(E)$. Then for any $\pi \in \widehat{G}$ there is an isomorphism as Hilbert B-modules between $p_0(H_\pi \otimes E)$ and $p_0(E \otimes H_\pi)$.*

Proof. First we observe the difference of $\hat{\delta}(p_0)$ and its opposite $\hat{\delta}^{op}(p_0)$. Using $p_0 = (\mathrm{id} \otimes h)(V)$ and the pentagonal identity, we obtain $\hat{\delta}(p_0) = (\mathrm{id} \otimes \mathrm{id} \otimes h)(V_{13}V_{23})$ and $\hat{\delta}^{op}(p_0) = (\mathrm{id} \otimes \mathrm{id} \otimes h)(V_{23}V_{13})$. Since V has the expansion $\sum_{\pi \in \widehat{G}, i,j \in I_\pi} E^\pi_{i,j} \otimes w(\pi)_{i,j}$, we have the following equalities:

$$\begin{aligned} \hat{\delta}(p_0) &= \sum_{\pi \in \widehat{G}, i,j \in I_\pi, k,\ell \in I_{\overline{\pi}}} h(w(\pi)_{i,j} w(\overline{\pi})_{k,\ell}) E^\pi_{i,j} \otimes E^{\overline{\pi}}_{k,\ell}\,, \\ \hat{\delta}^{op}(p_0) &= \sum_{\pi \in \widehat{G}, i,j \in I_\pi, k,\ell \in I_{\overline{\pi}}} h(w(\overline{\pi})_{k,\ell} w(\pi)_{i,j}) E^\pi_{i,j} \otimes E^{\overline{\pi}}_{k,\ell}\,. \end{aligned}$$

We use $\sigma^h_{-i}(w(\overline{\pi})_{k,\ell}) = (F_{\overline{\pi}})_{k,k}(F_{\overline{\pi}})_{\ell,\ell} w(\overline{\pi})_{k,\ell}$ in order to get $h(w(\overline{\pi})_{k,\ell} w(\pi)_{i,j}) = (F_{\overline{\pi}})_{k,k}(F_{\overline{\pi}})_{\ell,\ell} h(w(\pi)_{i,j} w(\overline{\pi})_{k,\ell})$. Define a densely defined unbounded positive operator $F = \sum_{\pi \in \widehat{G}} F_\pi$. With this operator we get the identity

$$\hat{\delta}^{op}(p_0) = (1 \otimes F)\hat{\delta}(p_0)(1 \otimes F)\,.$$

Then we define the map $\chi : H_\pi \otimes E \longrightarrow E \otimes H_\pi$ by $\chi(\xi \otimes e) = e \otimes F_\pi^{-1}\xi$. This is a bijective adjointable map for Hilbert B-modules. Then we get

$$\begin{aligned}\chi \circ (\mathrm{id} \otimes \phi)(\hat{\delta}(p_0))(\xi \otimes e) &= (1 \otimes F_\pi^{-1})(\phi \otimes \mathrm{id})(\hat{\delta}^{op}(p_0))(e \otimes \xi) \\ &= (\phi \otimes \mathrm{id})(\hat{\delta}(p_0))(e \otimes F_\pi \xi) .\end{aligned}$$

Hence we see χ maps $p_0(H_\pi \otimes E)$ onto $p_0(E \otimes H_\pi)$. □

Lemma 4.2. *The Green-Julg isomorphism Φ_1 is an $R(G)$-module map.*

Proof. Take $\pi \in \widehat{G}$ and a G-equivariant A-module E. We shall prove $\Phi_1(\pi \cdot [E]) = (\hat{\alpha}_\pi)_*(\Phi_1([E]))$. For the left side we have

$$[p_0((H_\pi \otimes E) \otimes_A A \rtimes_\alpha G)] = [p_0(H_\pi \otimes E \rtimes_\alpha G)] .$$

For the right one we have

$$\begin{aligned}(\hat{\alpha}_\pi)_*([p_0(E \rtimes_\alpha G)]) &= [p_0(E \rtimes_\alpha G) \otimes_{\hat{\alpha}_\pi} \big(A \rtimes_\alpha G \otimes \mathrm{End}(H_\pi)\big)] \\ &= [p_0(E \rtimes_\alpha G \otimes \mathrm{End}(H_\pi))] .\end{aligned}$$

Moving it to $K_0(A \rtimes_\alpha G)$ by tensoring $A \rtimes_\alpha G \otimes H_\pi$ over $A \rtimes_\alpha G \otimes \mathrm{End}(H_\pi)$, we get: $(\hat{\alpha}_\pi)_*(\Phi_1([E])) = [p_0(E \rtimes_\alpha G \otimes H_\pi)]$. Hence we obtain the desired equality by applying the previous lemma to $E \rtimes_\alpha G$ and $B = A \rtimes_\alpha G$. □

We consider a compact quantum ergodic system $\{A, G, \alpha\}$. Then there exists an index set I and Hilbert spaces H_i for each $i \in I$ such that the crossed product $A \rtimes_\alpha G$ is isomorphic to $\bigoplus_{i \in I} \mathbb{K}(H_i)$, where $\{H_i\}_{i \in I}$ are Hilbert spaces (see [B, Theorem 19], [Wa1, Corollary 2] for its proof). Let us fix minimal projections $e_i \in \mathbb{K}(H_i)$ for all $i \in I$. Hence we have $K_0^G(A) \cong \bigoplus_{i \in I} \mathbb{Z}[e_i]$ by the Green-Julg theorem. Using the isomorphism $\Phi_2 : K_0(A \rtimes_\alpha G) \longrightarrow K_0^G(A)$, we can easily check that in $K_0^G(A)$, $[e_i]$ becomes a G-equivariant A-module $[e_i(A \otimes L^2(G))]$. Now we consider the $R(G)$-module structure of $K_0^G(A)$. Let π be an element of $\widehat{G}$. We define the (not necessarily finite size) matrix $\mathbb{M}(\pi) = (\mathbb{M}(\pi)_{i,j})_{i,j \in I}$ by

$$\pi \cdot [e_j] = \sum_{i \in I} \mathbb{M}(\pi)_{i,j} [e_i].$$

This equality holds in $K_0^G(A) \cong K_0(A \rtimes_\alpha G)$. If we treat $[e_i] = [e_i(A \otimes L^2(G))]$ in $K_0^G(A)$, we get the equality $\mathbb{M}(\pi)_{i,j} = \dim \mathrm{Hom}_{(G,A)}(H_\pi \otimes e_j(A \otimes L^2(G)), e_i(A \otimes L^2(G)))$.

Lemma 4.3. *We have an isomorphism between linear spaces* $\mathrm{Hom}_{(G,A)}(H_\pi \otimes e_j(A \otimes L^2(G)), e_i(A \otimes L^2(G)))$ *and* $\mathrm{Hom}_G(H_\pi, e_i(A \otimes \mathbb{K}(L^2(G)))e_j)$.

Proof. We may assume projections e_i and e_j are majored by $1 \otimes p$ where p is a projection in $C_r^*(G) \subset \mathbb{K}(L^2(G))$. Let H_0 be a closed subspace $pL^2(G)$. It is finite dimensional and G-invariant, that is, $W(H \otimes \mathbb{C}) \subset H \otimes C(G)$. Let $\{\eta_p\}_{p \in J}$ be an orthonormal basis of H_0. With this basis we give a matrix representation $(w_{p,q})_{p,q \in J}$ of W by $W(\eta_p \otimes 1) = \sum_{q \in J} \eta_q \otimes w_{q,p}$. From now we write Γ_0 and Γ_1

for $C(G)$-comodule maps $H_\pi \otimes A \otimes H_0 \longrightarrow H_\pi \otimes A \otimes H_0 \otimes C(G)$ and $A \otimes H_0 \longrightarrow A \otimes H_0 \otimes C(G)$, respectively. Take an intertwiner S from $\mathrm{Hom}_{(G,A)}(H_\pi \otimes A \otimes H_0, A \otimes H_0)$. Then we can choose $a_{k,qp} \in A$ for all $k \in I_\pi$ and $p, q \in J$ such that they satisfy $S(\xi_k \otimes a \otimes \eta_p) = \sum_{q \in J} a_{k,qp} a \otimes \eta_q$ for all $a \in A$ by A-linearity. Since S is a G-homomorphism, we have $\Gamma_1 S(\xi_k \otimes 1 \otimes \eta_p) = (S \otimes 1)(\Gamma_0(\xi_k \otimes 1 \otimes \eta_p))$. The the left hand side is equal to

$$\sum_{q \in J} \Gamma_1(a_{k,qp} \otimes \eta_q) = \sum_{q,r \in J} (1 \otimes \eta_r \otimes w_{rq}) \alpha(a_{k,qp})_{13} \,.$$

The right hand side is equal to

$$\sum_{\ell \in I_\pi, s \in J} (S \otimes 1)(\xi_\ell \otimes 1 \otimes \eta_s \otimes w(\pi)_{\ell k} w_{sp}) = \sum_{\ell \in I_\pi, r, s \in J} a_{\ell,rs} \otimes \eta_r \otimes w(\pi)_{\ell k} w_{sp} \,.$$

Hence we get

$$\sum_{q \in J} (1 \otimes w_{rq}) \alpha(a_{k,qp}) = \sum_{\ell \in I_\pi, s \in J} a_{\ell,rs} \otimes w(\pi)_{\ell k} w_{sp} \,. \tag{4.1}$$

Next take an intertwiner T from $\mathrm{Hom}_G(H_\pi, A \otimes \mathbb{B}(H_0))$. Then we can choose $b_{k,qp} \in A$ for all $k \in I_\pi$ and $p, q \in J$ such that they satisfy $T\xi_k = \sum_{q,p \in J} b_{k,qp} \otimes \eta_q \odot \eta_p$. Since T is a G-homomorphism, we have $\tilde{\alpha} \circ T(\xi_k) = (T \otimes 1) \circ w(\pi)(\xi_k \otimes 1)$ for all $k \in I_\pi$. The left hand side is equal to

$$\begin{aligned} \sum_{q,t \in J} \tilde{\alpha}(b_{k,qt} \otimes \eta_q \odot \eta_t) &= \sum_{q,t \in J} W_{23} \alpha(b_{k,qt})_{13} (1 \otimes 1 \otimes \eta_q \odot \eta_t) W_{23}^* \\ &= \sum_{q,r,s,t \in J} (1 \otimes \eta_r \odot \eta_s \otimes 1)(1 \otimes 1 \otimes w_{rq}) \alpha(b_{k,qt})_{13} (1 \otimes 1 \otimes w_{st}^*) \,. \end{aligned}$$

The right hand side is equal to

$$\sum_{\ell \in I_\pi} T\xi_\ell \otimes w(\pi)_{\ell k} = \sum_{\ell, r, s} b_{\ell,rs} \otimes \eta_r \odot \eta_s \otimes w(\pi)_{\ell k} \,.$$

Hence we obtain the following equality

$$\sum_{q,t \in J} (1 \otimes w_{rq}) \alpha(b_{k,qt})(1 \otimes w_{st}^*) = \sum_{\ell} b_{\ell,rs} \otimes w(\pi)_{\ell k} \,.$$

Multiplying w_{sp} and summing up with s, we get

$$\sum_{q \in J} (1 \otimes w_{rq}) \alpha(b_{k,qp}) = \sum_{\ell \in I_\pi, s \in J} b_{\ell,rs} \otimes w(\pi)_{\ell k} w_{sp} \,. \tag{4.2}$$

Now we construct maps $\nu : \mathrm{Hom}_{(G,A)}(H_\pi \otimes A \otimes H_0, A \otimes H_0) \longrightarrow \mathrm{Hom}_G(H_\pi, A \otimes \mathbb{B}(H_0))$ and $\nu^{-1} : \mathrm{Hom}_G(H_\pi, A \otimes \mathbb{B}(H_0)) \longrightarrow \mathrm{Hom}_{(G,A)}(H_\pi \otimes A \otimes H_0, A \otimes H_0)$ by the following equalities. For all $k \in I_\pi$ and $p \in J$,

$$\nu(S)(\xi_k) = \sum_{q,p \in J} a_{k,qp} \otimes \eta_q \odot \eta_p \,,$$

$$\nu^{-1}(T)(\xi_k \otimes a \otimes \eta_p) = \sum_{q \in J} b_{k,qp} a \otimes \eta_q \,.$$

They are actually linear operators in each hom-space because equalities (4.1) and (4.2) have the same form. Since clearly we have $\nu \circ \nu^{-1} = \mathrm{id}$ and $\nu^{-1} \circ \nu = \mathrm{id}$, they give an isomorphism between $\mathrm{Hom}_{(G,A)}(H_\pi \otimes A \otimes H_0, A \otimes H_0)$ and $\mathrm{Hom}_G(H_\pi, A \otimes \mathbb{B}(H_0))$. Then we also easily check that ν maps the subspace $\mathrm{Hom}_{(G,A)}(H_\pi \otimes e_j(A \otimes L^2(G)), e_i(A \otimes L^2(G))) \subset \mathrm{Hom}_{(G,A)}(H_\pi \otimes A \otimes H_0, A \otimes H_0)$ to the subspace $\mathrm{Hom}_G(H_\pi, e_i(A \otimes \mathbb{K}(L^2(G)))e_j) \subset \mathrm{Hom}_G(H_\pi, A \otimes \mathbb{B}(H_0))$. ☐

Therefore, as in the classical case we can derive the following result by A. Wassermann ([Wa1, p.304]).

Corollary 4.4. *Let $\{A, G, \alpha\}$ be a compact quantum ergodic system and $\mathbb{M} : R(G) \longrightarrow M_{|I|}(\mathbb{Z})$ be the multiplicity map. For all i, j in the index set I, we have*

$$\mathbb{M}(\pi)_{i,j} = \dim \mathrm{Hom}(H_\pi, e_i(A \otimes \mathbb{K}(L^2(G)))e_j).$$

It yields the identity $\mathbb{M}(\overline{\pi})_{i,j} = \mathbb{M}(\pi)_{j,i}$ for all $\pi \in \widehat{G}$ and $i, j \in I$ and therefore the multiplicity map $\mathbb{M} : R(G) \longrightarrow M_{|I|}(\mathbb{Z})$ is a $*$-homomorphism. As we have seen, some properties of multiplicity maps also hold in the quantum case. However we have to be much careful about the existence of its common eigenvector (see, [Wa1, Theorem17] for classical case). In order to study this problem we need several lemmas.

Lemma 4.5. *For all $i, j \in I$, the reduced G-space $e_i(A \otimes \mathbb{K}(L^2(G)))e_j$ is not* 0.

Proof. Let M be the σ-weak closure of A in $\mathbb{B}(H_\varphi)$ where H_φ is the GNS Hilbert space of the invariant state φ. Define the relation $\sim$ in the index set I as $i \sim j$ if and only if $e_i(A \otimes \mathbb{K}(L^2(G)))e_j \neq 0$. We show this is an equivalence relation. A non-trivial part is transitivity of $\sim$. Assume $i \sim j$ and $j \sim k$. Then there exists $\pi, \rho \in \widehat{G}$ which satisfy $\mathbb{M}(\pi)_{i,j} > 0$ and $\mathbb{M}(\rho)_{j,k} > 0$. Then we have $\mathbb{M}(\pi \cdot \rho)_{i,k} \geq \mathbb{M}(\pi)_{i,j}\, \mathbb{M}(\rho)_{j,k} > 0$ and the left hand side is equal to $\sum_{\tau \in \widehat{G}} N_\tau^{\pi\,\rho}\, \mathbb{M}(\tau)_{i,k}$. Hence there exists $\tau \in \widehat{G}$ such that $\mathbb{M}(\tau)_{i,k} > 0$. Therefore the reduced space $e_i(A \otimes \mathbb{K}(L^2(G)))e_k$ is not 0, that is, $i \sim k$. Let $I = \bigsqcup_{\lambda \in \Lambda} I_\lambda$ be the decomposition with respect to the equivalence relation $\sim$. Let z_i be the central projection in $Z(M)$ with $z(e_i) = z_i \otimes 1$ for all $i \in I$ where $z(e_i)$ means the central support projection of e_i in $M \otimes \mathbb{B}(L^2(G))$. Since for $\lambda \neq \mu \in \Lambda$ we have $e_i(A \otimes \mathbb{K}(L^2(G)))e_j = 0$ for all $i \in I_\lambda$ and $j \in I_\mu$, we see $z_i\, z_j = 0$ for all $i \in I_\lambda$ and $j \in I_\mu$. We claim $1 = \bigvee_{i \in I} z_i = \sum_{\lambda \in \Lambda} \bigvee_{i \in I_\lambda} z_i$. This follows because $A \rtimes_\alpha G \cong \bigoplus_{i \in I} \mathbb{K}(H_i)$ and the sum of all diagonal atoms is equal to 1 in strong operator topology. By definition of a central support projection we obtain $Mz_i \otimes \mathbb{B}(L^2(G)) = M \otimes \mathbb{B}(L^2(G))\, e_i\, M \otimes \mathbb{B}(L^2(G))$ for all $i \in I$. Then we see $\tilde{\alpha}(Mz_i \otimes \mathbb{B}(L^2(G))) \subset (M \otimes \mathbb{B}(L^2(G))e_i M \otimes \mathbb{B}(L^2(G))) \otimes L^\infty(G) = Mz_i \otimes \mathbb{B}(L^2(G)) \otimes L^\infty(G)$ and hence $\tilde{\alpha}(z_i \otimes 1) \leq z_i \otimes 1 \otimes 1$ for all $i \in I$. Let z_λ be the projection $\bigvee_{i \in I_\lambda} z_i$. Then a family $\{z_\lambda\}_{\lambda \in \Lambda}$ is a partition of unity and satisfies $\tilde{\alpha}(z_\lambda \otimes 1) \leq z_\lambda \otimes 1 \otimes 1$ for all $\lambda \in \Lambda$. Therefore we obtain $\tilde{\alpha}(z_\lambda \otimes 1) = z_\lambda \otimes 1 \otimes 1$ for all $\lambda \in \Lambda$. Since the left hand side is equal to $W_{23}\alpha(z_\lambda)_{13}W_{23}^*$, we have $\alpha(z_\lambda) = z_\lambda \otimes 1$ for all $\lambda \in \Lambda$.

Ergodicity of α yields $z_\lambda = 1$ or 0 for all $\lambda \in \Lambda$. Hence Λ is a singleton and it completes the proof. $\square$

From this lemma we have the following irreducibility criterion which has been studied in [Wa1, Section 10].

Lemma 4.6. *Let π be an element of $\widehat{G}$. If for any $\rho \in \widehat{G}$ there exists $k \in \mathbb{Z}_{\geq 0}$ such that ρ is contained in π^k, then for any $i, j \in I$ there exists $\ell \in \mathbb{Z}_{\geq 0}$ with $(\mathbb{M}(\pi)^\ell)_{i,j} > 0$.*

Proof. Since G-space $e_i(A \otimes \mathbb{K}(L^2(G)))e_j$ is not 0, there exists an element ρ in I such that $\mathbb{M}(\rho)_{i,j} = \dim \mathrm{Hom}(H_\rho, e_i(A \otimes \mathbb{K}(L^2(G)))e_j) > 0$. By assumption there exists $\ell \in \mathbb{N}$ such that π^ℓ contains ρ. Then we have $(\mathbb{M}(\pi)^\ell)_{i,j} = \mathbb{M}(\pi^\ell)_{i,j} \geq \mathbb{M}(\rho)_{i,j} > 0$. $\square$

In that case we want to show the existence of a Perron-Frobenius eigenvector, however, we cannot use the dual weight $\hat{\varphi}$ as in the compact group case because it is not necessarily a trace. We will give an eigenvector under some assumptions later.

Lemma 4.7. *Let $m : R(G) \longrightarrow \mathbb{Z}$ be a $*$-homomorphism such that it satisfies $m(\pi_0) = 1$ and $m(\widehat{G}) \subset \mathbb{Z}_{>0}$. Assume $\mathbb{M}(\pi)_{i,j} \leq m(\pi)$ for all $\pi \in \widehat{G}$ and $i, j \in I$. Then the matrix $\mathbb{M}(\rho)$ is a bounded operator on $l^2(I)$ and its norm is less than or equal to $m(\rho)^2$ for all $\rho \in R(G)_+$.*

Proof. Take ρ from $R(G)_+$ and we have $\sum_{j \in I} \mathbb{M}(\rho)_{i,j}^2 \leq m(\rho)^2$. In fact we can directly check it by the following calculation

$$\begin{aligned}
\sum_{j \in I} \mathbb{M}(\rho)_{i,j}^2 =& (\mathbb{M}(\rho)\mathbb{M}(\overline{\rho}))_{i,i} \\
=& \mathbb{M}(\rho \cdot \overline{\rho})_{i,i} \\
=& \sum_{\sigma \in \widehat{G}} N_\sigma^{\rho\bar{\rho}} \mathbb{M}(\sigma)_{i,i} \\
\leq& \sum_{\sigma \in \widehat{G}} N_\sigma^{\rho\bar{\rho}} m(\sigma) \\
=& m(\rho) m(\bar{\rho}) \\
=& m(\rho)^2 \,.
\end{aligned}$$

Take a finitely supported vector ξ in $l^2(\mathbb{Z})$. Let $C(i, \rho)$ be a finite set $\{j \in I \mid \mathbb{M}(\rho)_{i,j} \neq 0\}$. From the above inequality we particularly obtain $|C(i, \rho)| \leq m(\rho)^2$.

We assume $\rho = \overline{\rho} \in R(G)_+$. Then we have the desired inequality

$$\begin{aligned}
||\mathbb{M}(\rho)\xi||^2 &= \sum_{i\in I}\Big|\sum_{j\in I}\mathbb{M}(\rho)_{i,j}\xi_j\Big|^2 \\
&= \sum_{i\in I}\Big|\sum_{j\in C(i,\rho)}\mathbb{M}(\rho)_{i,j}\xi_j\Big|^2 \\
&= \sum_{i\in I}\sum_{j\in C(i,\rho)}\mathbb{M}(\rho)_{i,j}^2 \cdot \sum_{j\in C(i,\rho)}|\xi_j|^2 \\
&\leq m(\rho)^2\sum_{i\in I}\sum_{j\in C(i,\rho)}|\xi_j|^2 \\
&= m(\rho)^2\sum_{j\in I}|C(j,\rho)||\xi_j|^2 \\
&\leq m(\rho)^4\sum_{j\in I}|\xi_j|^2 \,.
\end{aligned}$$

For general $\rho \in R(G)_+$ we apply the above result to a positive self-conjugate $\overline{\rho}\rho$ and get

$$\begin{aligned}
||\mathbb{M}(\rho)||^2 &= ||\mathbb{M}(\overline{\rho}\rho)|| \\
&\leq m(\overline{\rho}\rho)^2 \\
&= m(\rho)^4 \,.
\end{aligned}$$

□

Lemma 4.8. *If a self-conjugate element $\pi \in \widehat{G}$ has the property in Lemma 4.6, then $\mathbb{M}(\pi)$ has an eigenvector with the eigenvalue $||\mathbb{M}(\pi)||(\leq D_\pi^2)$. Moreover if A has a tracial state, then $||\mathbb{M}(\pi)||(\leq d_\pi^2)$.*

Proof. Since we know the matrix $\mathbb{M}(\pi)$ is irreducible and bounded, there exists an eigenvector with the eigenvalue $||\mathbb{M}(\pi)||$. For all $i, j \in I$ we have proved $e_i(A\otimes\mathbb{K}(L^2(G)))e_j \neq 0$. By Lemma 3.9 we have $\mathbb{M}(\pi)_{i,j} \leq D_\pi$. Hence we obtain the first assertion by Lemma 4.7. If A has a tracial state τ, then the tracial weight $\tilde{\tau} = \tau \otimes \mathrm{Tr}$ is semifinite on $A \rtimes_\alpha G$ and $\tilde{\tau}(e_i) > 0$ for all $i \in I$. In fact if $\tilde{\tau}(e_i)$ is equal to 0 for some $i \in I$, we have $\tilde{\tau}(e_j) = 0$ for all $j \in I$ by Remark 3.12 and Lemma 4.5. It shows $\tilde{\tau} = 0$ and this is contradiction. Now we apply Lemma 4.7 with $m(\pi) = d_\pi$ and obtain the second assertion. □

Now we consider the special case $G = SU_q(2)$. We refer to [Wo1, MMNNU] or the next section for its representation theory of $SU_q(2)$. Its irreducible representations are represented by $\{\pi_\nu\}_{\nu\in\frac{1}{2}\mathbb{Z}_{>0}}$ and its fusion rules are determined by

$$\pi_{\frac{1}{2}} \cdot \pi_\nu = \pi_{\nu+\frac{1}{2}} + \pi_{\nu-\frac{1}{2}}$$

for all $\nu \in \frac{1}{2}$.

Lemma 4.9. *Let $m : R(SU_q(2)) \longrightarrow \mathbb{Z}$ be a dimension function with $m(\pi_\nu) \leq d_{\pi_\nu}^2$ for all $\nu \in \frac{1}{2}$. Then we obtain $m(\pi_\nu) = d_{\pi_\nu}$ for all $\nu \in \frac{1}{2}$.*

Proof. Write t_0 for $m(\pi_{\frac{1}{2}})$. Define polynomials $\{p_\nu\}_{\nu \in \frac{1}{2}\mathbb{Z}}$ recursively by $p_{\nu+\frac{1}{2}}(s) = s\, p_\nu(s) - p_{\nu-\frac{1}{2}}(s)$ and $p_0 = 1$, $p_{\frac{1}{2}} = s$. Then we get $m(\pi_\nu) = p_\nu(t_0)$. From the positivity of $m(\pi_\nu)$ we have $t_0 \geq 2$. Assume $t_0 > 2$. For $s > 2$, the polynomials $p_\nu(s)$ are strictly increasing and we get a lower bound by an affine line $p_\nu(2)'(s-2) + p_\nu(2)$ where $p_\nu(s)'$ means the derivative at s. Putting $s = t_0$, we obtain an equality $m(\pi_\nu) \geq (t-2)p_\nu(2)' + 2\nu + 1$. The left hand side has upper bound by $d_{\pi_\nu}^2 = (2\nu+1)^2$. In order to derive contradiction, it suffices to show the right hand side has a polynomial degree 3 with respect to ν. By the definition of p_ν, we get $p_{\nu+\frac{1}{2}}(2)' - p_\nu(2)' = p_\nu(2)' - p_{\nu-\frac{1}{2}}(2)' + 2\nu + 1$. This immediately gives $p_\nu(2)' = \frac{2}{3}\nu(\nu+1)(2\nu+1)$. $\square$

We show the following main result in this section which asserts the existence of multiplicity vector $\boldsymbol{c}$ under a tracial condition on A.

Theorem 4.10. *Let $\{A, SU_q(2), \alpha\}$ be a compact quantum ergodic system. Assume that A has a tracial state. Then there exists a positive entry vector $\boldsymbol{c} = (c_i)_{i \in I}$ such that $\mathbb{M}(\pi_\nu)\, \boldsymbol{c} = d_{\pi_\nu}\, \boldsymbol{c}$ for all $\nu \in \frac{1}{2}$.*

Proof. By Lemma 4.8 there exists an eigenvector $\boldsymbol{c}$ such that $\mathbb{M}(\pi_{\frac{1}{2}})\, \boldsymbol{c} = t\, \boldsymbol{c}$ where $t = \big|\big|\mathbb{M}(\pi_{\frac{1}{2}})\big|\big|$. Since any π_ν is written by the polynomial P_ν of $\pi_{\frac{1}{2}}$, we can define the dimension function $m : R(SU_q(2)) \longrightarrow \mathbb{Z}$ with $\mathbb{M}(\pi_\nu)\, \boldsymbol{c} = m(\pi_\nu)\, \boldsymbol{c}$ for all $\nu \in \frac{1}{2}$. We show $m(\pi_\nu) = ||\mathbb{M}(\pi_\nu)||$ for all $\nu \in \frac{1}{2}$. The self-adjoint operator $\mathbb{M}(\pi_\nu)$ is written by the polynomial P_ν of $\mathbb{M}(\pi_{\frac{1}{2}})$. Hence we have $||\mathbb{M}(\pi_\nu)|| \geq P_\nu(\big|\big|\mathbb{M}(\pi_{\frac{1}{2}})\big|\big|) = P_\nu(t) = m(\pi_\nu)$. The converse inequality is obtained by applying the Schur test to $\mathbb{M}(\pi_\nu)\, \boldsymbol{c} = m(\pi_\nu)\, \boldsymbol{c}$. Therefore we have $m(\pi_\nu) = ||\mathbb{M}(\pi_\nu)|| \leq d_{\pi_\nu}^2$. By previous lemma, we have $m(\pi_\nu) = d_{\pi_\nu}$ for all $\nu \in \frac{1}{2}$. $\square$

The next lemma has already been proved in [Wa2, Lemma 1, Theorem 2] for $q = 1$, that is, the $SU(2)$ case.

Lemma 4.11. *Let $\{A, SU(2), \alpha\}$ be an ergodic system. Then its $\pi_{\frac{1}{2}}$-eigenvector space $X_{\pi_{\frac{1}{2}}}$ is not one-dimensional.*

Proof. Let $\xi = (a, b)$ be a $\pi_{\frac{1}{2}}$-eigenvector. Then we have

$$\alpha(a) = a \otimes x + b \otimes v\,, \quad \alpha(b) = a \otimes u + b \otimes y.$$

Consider a vector $\eta = (b^*, -q^{-1}a^*)$. By an easy calculation, we see η is also a $\pi_{\frac{1}{2}}$-eigenvector. Assume that $X_{\pi_{\frac{1}{2}}}$ is one-dimensional. Then there exists a complex number μ such that $\eta = \mu\xi$. So we get $b^* = \mu a$ and $-q^{-1}a^* = \mu b$.

It follows $-q^{-1}a = |\mu|^2 a$. Hence we obtain $0 > -q^{-1} = |\mu|^2 > 0$. This is a contradiction. $\square$

Hence we obtain the result corresponding to [Wa2, Theorem 1] for positive q.

Corollary 4.12. *Let $\{A, SU(2), \alpha\}$ be an ergodic system. Assume that A has a tracial state and q is positive. Then its multiplicity diagram is one of type $1, \mathbb{T}_n \,(n \geq 2), \mathbb{T}, SU(2), D_n^* \,(n \geq 2), D_\infty^*, A_4^*, S_4^*$ and A_5^*.*

Example 4.13. We consider the multiplicity diagrams of quantum spheres $C(S^2_{q,\lambda})$. If $\lambda_0 = c(n)$, then its spectral pattern (or finite dimensionality) allows only the diagram of type $SU(2)$ in Figure 6 (see Appendix). If $0 \leq \lambda_0 \leq 1$, then its spectral pattern derives the diagram of type $\mathbb{T}$ in Figure 5. If $\{A, SU_q(2), \alpha\}$ is an ergodic system and A is finite dimensional, then its multiplicity diagram of $\pi_{\frac{1}{2}}$ must be of type $SU(2)$ by its finite dimensionality. Hence the classification by Podleś ([Po1]) shows that A is G-isomorphic to $\mathrm{End}(H_{\pi_\nu})$ for some $\nu \in \mathbb{Z}_{\geq \frac{1}{2}}$.

We recall the definition of the McKay diagrams. Let $H \subset G$ be a quantum subgroup with the restriction map r_H and $w \in \mathbb{B}(H_w) \otimes C(G)$ be a unitary representation of G. We denote the restricted representation $(\mathrm{id} \otimes r_H)(w)$ by $w|_H$. Prepare the vertices $\{\sigma\}_{\sigma \in \widehat{H}}$. Let σ_0 be the one-dimensional trivial representation of H. First we consider the irreducible decomposition $w|_H \cdot \sigma_0 = \oplus_{\sigma \in \widehat{H}} N_\sigma^{w|_H} \sigma$ where the scalar $N_\sigma^{w|_H}$ means a multiplicity. Then we draw arrows from σ_0 to the above arising irreducible representations $\{\sigma\}_{\sigma \in \widehat{H}}$ with $N_\sigma^{w|_H}$-times, respectively. Second for each σ in the above decomposition we consider the irreducible decomposition of $w|_H \cdot \sigma$ and we draw arrows from σ to the irreducible representations in a similar way. Continue this procedure we get the McKay diagram for $H \subset G$ with respect to w. If w is self-conjugate, then the diagram is non-oriented because of the symmetry $N_\sigma^{w|_H \tau} = N^\tau_{w|_H \sigma}$. When we treat quantum subgroups of $SU_q(2)$, the McKay diagrams are drawn with respect to the fundamental representation $\pi_{\frac{1}{2}}$. Now we consider G-covariant system $\{C(H \setminus G), \delta\}$ where H is a quantum subgroup with the restriction map r_H. It is well-known that the multiplicity diagram and the McKay diagram coincide in the classical case, however, we give the proof of the general quantum group case for readers' convenience. We denote $(r_H \otimes \mathrm{id})(V_\ell)$ by $V_\ell|_H$. The C^*-algebra $C(G) \otimes \mathbb{K}(L^2(G))$ has the left and right actions α_ℓ and β_r defined by $\alpha_\ell(x) = \mathrm{Ad}\, V_{\ell 13}^* V_{\ell 12}^* (1 \otimes x)$ and $\beta_r(x) = \mathrm{Ad}\, V_{13}(x \otimes 1)$ for all $x \in C(G) \otimes \mathbb{K}(L^2(G))$, respectively. We define the left H-action α_ℓ^H by the composition $(r_H \otimes \mathrm{id}) \circ \alpha_\ell$. Consider the map $\mathrm{Ad}\, V_\ell : C(G) \otimes \mathbb{K}(L^2(G)) \longrightarrow C(G) \otimes \mathbb{K}(L^2(G))$.

Lemma 4.14.
1. *The map $\mathrm{Ad}\, V_\ell$ gives an isomorphism between $C(H \setminus G) \otimes \mathbb{K}(L^2(G))$ and ${}^H(C(G) \otimes \mathbb{K}(L^2(G)))$ where the latter one is the fixed point algebra of the left H-action α_ℓ^H. Moreover it maps $C(H \setminus G) \rtimes_\delta G$ to $\mathbb{C} \otimes {}^H\mathbb{K}(L^2(G)))$.*
2. *$\mathrm{Ad}\, V_\ell$ intertwines the right G-actions $\widetilde{\alpha}$ and β_r.*

Proof. (1) It suffices to show that the map $\operatorname{Ad} V_\ell$ intertwines left H-actions $\operatorname{Ad} V_\ell|_H \otimes \mathrm{id}$ and α_ℓ^H. This is immediately verified by the equality $V_{\ell\,23} V_\ell|_H{}^*_{12} = V_\ell|_H{}^*_{13} V_\ell|_H{}^*_{12} V_{\ell\,23}$. (2) Similarly the equality $V_{\ell\,12} W_{23} V_{13} = V_{13} V_{\ell\,12}$ gives the desired intertwining property. □

Now we choose a left irreducible H-module K_σ^ℓ for each $\sigma \in \widehat{H}$. Then we have the irreducible decomposition $L^2(G) = \oplus_{\sigma \in \widehat{H}} K_\sigma^\ell \otimes L_\sigma$ where L_σ is the multiplicity space for σ. Note that all the L_σ are non-zero, because there exists $\pi \in \widehat{G}$ such that σ is contained in $\pi|_H$. Then we get ${}^H\mathbb{K}(L^2(G))) = \oplus_{\sigma \in \widehat{H}} \mathbb{C}1_\sigma \otimes \mathbb{K}(L_\sigma)$. Let $e_\sigma = 1_\sigma \otimes p_\sigma$ be a minimal projection of $\mathbb{C}1_\sigma \otimes \mathbb{K}(L_\sigma)$. By the previous lemma we have $K_0(C(H \setminus G) \rtimes_\delta G) = \oplus_{\sigma \in \widehat{H}} \mathbb{Z}[e_\sigma]$. Hence the vertices of multiplicity maps are represented by the elements of $\widehat{H}$. Then the following proposition holds.

Proposition 4.15. *Let π be an element of $\widehat{G}$. Then we have $\mathbb{M}(\pi)_{\rho,\sigma} = N_\sigma^{\pi|_H\,\rho}$ for all ρ, σ in $\widehat{H}$.*

Proof. We use Corollary 4.4. The G-module $e_\rho C(H \setminus G) \otimes \mathbb{K}(L^2(G)) e_\sigma$ is isomorphic to ${}^H(C(G) \otimes e_\rho \mathbb{K}(L^2(G)) e_\sigma)$ by the previous lemma. Since quantum group G acts on the first tensor component, the multiplicity space for π is ${}^H(H_\pi^\ell \otimes e_\rho \mathbb{K}(L^2(G)) e_\sigma)$ which is linearly isomorphic to $\operatorname{Hom}_H(H^\ell_{\overline{\pi|_H}}, K_\rho \otimes K_{\overline{\sigma}})$. Hence its dimension is equal to $N^{\rho\overline{\sigma}}_{\overline{\pi|_H}} = N_\sigma^{\pi|_H\,\rho}$. □

Define the vector $\boldsymbol{d} = (d_\sigma)_{\sigma \in \widehat{H}}$ and we have $\mathbb{M}(\pi)\boldsymbol{d} = d_\pi \boldsymbol{d}$. Hence we have solved the eigenvector problem in the case of the quotient spaces.

Corollary 4.16. *Let H be a quantum subgroup of G and consider the ergodic system $\{C(H \setminus G), \delta\}$. Then the multiplicity diagram for $\pi \in \widehat{G}$ coincides with the McKay diagram of $H \subset G$ with respect to π.*

Next we study reduced ergodic systems associated to ergodic systems and show the heredity of information of multiplicity maps.

Definition 4.17. Let $\{A, G, \alpha\}$ and $\{B, G, \alpha\}$ be compact quantum ergodic systems and take index sets I, J with $A \rtimes_\alpha G \cong \bigoplus_{i \in I} \mathbb{K}(H_i)$ and $B \rtimes_\beta G \cong \bigoplus_{j \in J} \mathbb{K}(H_i{}')$. We say two ergodic systems are of the same type if there exists a bijective map $\theta : I \longrightarrow J$ satisfying $\mathbb{M}^B(\pi) \circ \theta = \theta \circ \mathbb{M}^A(\pi)$ for all $\pi \in G$, where $\mathbb{M}^A$ and $\mathbb{M}^B$ are multiplicity maps for ergodic systems $\{A, G, \alpha\}$ and $\{B, G, \alpha\}$ respectively.

Theorem 4.18. *Let $\{A, G, \alpha\}$ be a compact quantum ergodic system and take an index set I with $A \rtimes_\alpha G \cong \bigoplus_{i \in I} \mathbb{K}(H_i)$. Take minimal projections $\{e_i\}_{i \in I}$ from $\{\mathbb{K}(H_i)\}_{i \in I}$. Then for all $i \in I$, two ergodic systems $\{A, G, \alpha\}$ and $\{e_i(A \otimes \mathbb{K}(L^2(G)))e_i, G, \tilde{\alpha}\}$ are of the same type.*

Proof. Denote $\mathbb{K}(L^2(G))$ and $\mathbb{B}(L^2(G))$ simply by $\mathbb{K}$ and $\mathbb{B}$. We claim that the covariant system $\{A\otimes\mathbb{K}\otimes\mathbb{K}, G, \tilde{\tilde{\alpha}}\}$ is strongly equivalent to $\{\mathbb{K}\otimes A\otimes\mathbb{K}, G, \mathrm{id}\otimes\tilde{\alpha}\}$. Let us consider the map $\mathrm{Ad}(W_{23}^*) : A\otimes\mathbb{K}\otimes\mathbb{K} \longrightarrow A\otimes\mathbb{K}\otimes\mathbb{K}$. For $x \in A, k \in \mathbb{K}\otimes\mathbb{K}$ we have:

$$\begin{aligned} W_{23}^*\tilde{\tilde{\alpha}}(W_{23}(x\otimes k)W_{23}^*)W_{23} =& W_{23}^*W_{34}W_{24}W_{23}\alpha(x)_{14}(1\otimes k\otimes 1)W_{23}^*W_{24}^*W_{34}^*W_{23} \\ =& W_{34}\alpha(x)_{14}(1\otimes k\otimes 1)W_{34}^*, \end{aligned}$$

where we use the pentagonal identity for W, $W_{23}W_{13}W_{12} = W_{12}W_{23}$. Next flip the first and second tensor component of $A\otimes\mathbb{K}\otimes\mathbb{K}$ and the claim is proved. Hence we get an isomorphism $(A\otimes\mathbb{K})\rtimes_{\tilde{\alpha}} G \cong \mathbb{K}\otimes(A\rtimes_\alpha G)$. A projection $e_i\otimes 1$ is in the fixed point algebra of the multiplier algebra $M(A\otimes\mathbb{K}\otimes\mathbb{K})$ for $\tilde{\tilde{\alpha}}$. So it is mapped to a projection f_i in the multiplier algebra $M(\mathbb{K}\otimes A\rtimes_\alpha G)$. Hence we see that $\{e_i(A\otimes\mathbb{K})e_i\otimes\mathbb{K}, G, \tilde{\tilde{\alpha}}\}$ is equivalent to $\{f_i(\mathbb{K}\otimes A\otimes\mathbb{K})f_i, G, \mathrm{id}\otimes\tilde{\alpha}\}$. The projection f_i is decomposed into a direct sum of projections $\{p_j\}_{j\in I}$ which are in $\{\mathbb{B}\otimes\mathbb{B}(H_j)\}_{j\in I}$. We claim they are non-zero projections. It suffices to prove the central support of f_i in $\mathbb{B}\otimes M\rtimes_\alpha G$ is equal to 1 by passing to von Neumann algebras, that is, isomorphic covariant systems $\{e_i(M\otimes\mathbb{B})e_i\otimes\mathbb{B}, G, \tilde{\tilde{\alpha}}\}$ and $\{f_i(\mathbb{B}\otimes M\otimes\mathbb{B})f_i, G, \mathrm{id}\otimes\tilde{\alpha}\}$. Hence we show the central support of $e_i\otimes 1$ in $(M\otimes\mathbb{B}\otimes\mathbb{B})^{\tilde{\tilde{\alpha}}} = (M\otimes\mathbb{B})\rtimes_{\tilde{\alpha}} G$ is equal to 1. By duality theorem, we get the isomorphism of inclusions, $M\rtimes_\alpha G\otimes\mathbb{C} \subset (M\otimes\mathbb{B})\rtimes_{\tilde{\alpha}} G \cong \hat{\alpha}(M\rtimes_\alpha G) \subset M\rtimes_\alpha G\otimes\mathbb{B}$. Hence we study the central support of $\hat{\alpha}(e_i)$ in $M\rtimes_\alpha G\otimes\mathbb{B}$. Fix $j\in I$ and we can take $\pi\in\widehat{G}$ with $\mathbb{M}(\pi)_{j,i} > 0$ by Lemma 4.5. By the definition of $\mathbb{M}(\pi)$ we have $[\hat{\alpha}_\pi(e_i)] = \sum_{k\in I}\mathbb{M}(\pi)_{k,i}[e_k]$. Hence the j-th component of $\hat{\alpha}_\pi(e_i)$ is a non-zero projection, in particular the central support of $\hat{\alpha}(e_i)$ in $M\rtimes_\alpha G\otimes\mathbb{B}$ is equal to 1. Let ${H_j}'$ be a Hilbert space $p_j(L^2(G)\otimes H_j)$ and we have an isomorphism $e_i(A\otimes\mathbb{K})e_i\rtimes_{\tilde{\alpha}} G \cong \bigoplus_{j\in I}\mathbb{K}({H_j}')$. Next we choose minimal projections $\{q_j\}_{j\in I}$ from $\mathbb{K}({H_j}')$. Let p be a minimal projection in $\mathbb{K}$. For any $j\in I$, the projection q_j is equivalent to $p\otimes e_j$ in $\mathbb{K}\otimes\mathbb{K}(H_j) \subset (\mathbb{K}\otimes A\otimes\mathbb{K})^{\mathrm{id}\otimes\tilde{\alpha}}$. Then we have a G-isomorphism between reduced spaces $q_j(\mathbb{K}\otimes A\otimes\mathbb{K})q_k$ and $(p\otimes e_j)(\mathbb{K}\otimes A\otimes\mathbb{K})(p\otimes e_k) = \mathbb{C}p\otimes e_j(A\otimes\mathbb{K})e_k$ for all $j,k\in I$. Therefore we have proved the equality $\dim\mathrm{Hom}_G(H_\pi, q_j(\mathbb{K}\otimes A\otimes\mathbb{K})q_k) = \dim\mathrm{Hom}_G(H_\pi, e_j(A\otimes\mathbb{K})e_k)$ for all $\pi\in\widehat{G}$ and $j,k\in I$. □

Definition 4.19. Let $\{A, G, \alpha\}$ be a compact quantum ergodic system. A unital C^*-subalgebra $B\subset A$ is called a G-invariant C^*subalgebra if it satisfies $\alpha(B)\subset B\otimes C(G)$. Then $\{B, G, \alpha\}$ is called a subsystem of $\{A, G, \alpha\}$ and denote this situation by $\{B, G, \alpha\}\subset\{A, G, \alpha\}$.

Let B be a G-invariant C^*-subalgebra of A. The inclusion G-homomorphism $\iota : B\longrightarrow A$ induces the inclusion of crossed products, $B\rtimes_\alpha G\subset A\rtimes_\alpha G$. Taking index sets I, J as $B\rtimes_\alpha G\cong\bigoplus_{i\in I}\mathbb{K}(H_i)$ and $A\rtimes_\alpha G\cong\bigoplus_{j\in J}\mathbb{K}(K_i)$. Let Λ be an inclusion matrix of $B\rtimes_\alpha G\subset A\rtimes_\alpha G$, that is, $\mathbb{K}(H_i)$ is amplified into $\mathbb{K}(K_j)$ by $\Lambda_{j,i}$ times. We can easily show the next proposition by the definition of multiplicity maps.

Proposition 4.20. *Let* $\{B, G, \alpha\} \subset \{A, G, \alpha\}$ *be an inclusion of compact quantum ergodic systems. Let* $\mathbb{M}^B$ *and* $\mathbb{M}^A$ *be the multiplicity maps. Then we have* $\Lambda\, \mathbb{M}^B(\pi) = \mathbb{M}^A(\pi)\, \Lambda$ *for all* $\pi \in \widehat{G}$*.*

We recall an observation by A. Wassermann, which has an important role in his classification program in [Wa2]. Let $p_0 = (\mathrm{id} \otimes h)(V)$ be a minimal projection of $C_r^*(G)$. Note that it is central in not only $C_r^*(G)$ but also $C_\ell^*(G)$. It is also a minimal projection on $L^2(G)$. Then we have

$$(1 \otimes p_0)A \rtimes_\alpha G(1 \otimes p_0) = \big(A \otimes p_0 \mathbb{K}(L^2(G))p_0\big)^{\tilde{\alpha}} = A^\alpha \otimes \mathbb{C}p_0 = \mathbb{C}(1 \otimes p_0).$$

This shows that p_0 is a minimal projection in $A \rtimes_\alpha G$. Hence if we take minimal projections in $\{e_i\}_{i \in I}$ as before, then there exists unique i_0 such that e_{i_0} is equivalent to p_0 in $A \rtimes_\alpha G$. We often say this index corresponds to p_0. Then we obtain an isomorphism as covariant systems between $\{A, G, \alpha\}$ and $\{e_{i_0} A \otimes \mathbb{K}(L^2(G)) e_{i_0}, G, \tilde{\alpha}\}$. From this we get the following result.

Corollary 4.21. *Let* $A \subset C(G)$ *be a right coideal. Then there exists an eigenvector* $\boldsymbol{c} = (c_i)_{i \in I}$ *which satisfies the following conditions.*

1. *All entries of* $\boldsymbol{c}$ *are strictly positive integers and there exists an index* $i_0 \in I$ *which satisfies* $c_{i_0} = 1$*.*
2. *It is a common eigenvector of the multiplicity map;* $\mathbb{M}(\pi)\boldsymbol{c} = d_\pi \boldsymbol{c}$ *for all* $\pi \in \widehat{G}$*.*
3. *Two covariant systems* $\{A, G, \alpha\}$ *and* $\{e_{i_0} A \otimes \mathbb{K}(L^2(G)) e_{i_0}, G, \tilde{\alpha}\}$ *are isomorphic.*

Proof. Take a minimal projection p_π of $\mathrm{End}(H_\pi) \subset C_r^*(G)$ for $\pi \in \widehat{G}$. Let π be an element of $\widehat{G}$. Consider the dual coaction $\hat{\delta}_\pi : C_r^*(G) \longrightarrow C_r^*(G) \otimes \mathrm{End}(H_\pi)$. We want to compute $\hat{\delta}_\pi(p_0)$ in $K_0(C_r^*(G))$. Since we have the isomorphism $K_0(C_r^*(G)) \cong R(G) = K_0^G(\mathbb{C})$ by $[p_\pi] \to \pi$, we see $[\hat{\delta}_\pi(p_0)] = [p_\pi]$. Now let Λ be an inclusion matrix for $A \rtimes_\alpha G \subset C(G) \rtimes_\delta G$. It is actually a row vector because $C(G) \rtimes_\alpha G$ is isomorphic to $\mathbb{K}(L^2(G))$. By the previous proposition we have $\mathbb{M}'(\pi)\Lambda = \Lambda\mathbb{M}(\pi)$ for all $\pi \in \widehat{G}$, where $\mathbb{M}'$ and $\mathbb{M}$ are the multiplicity maps for $\{A, G, \alpha\}$ and $\{C(G), G, \delta\}$ respectively. Let us take $1 \otimes p_0$ for a minimal projection of $C(G) \rtimes_\delta G \cong \mathbb{K}(L^2(G))$. From the first part of this proof we have $\pi[p_0] = [p_\pi] = d_\pi[p_0]$ in $K_0(C(G) \rtimes_\delta G)$. Hence the action $\mathbb{M}'(\pi)$ is multiplication by d_π and we get $\Lambda\mathbb{M}(\pi) = d_\pi\Lambda$. Transposing it, we have $\mathbb{M}(\overline{\pi})^T\Lambda = d_{\overline{\pi}}{}^T\Lambda$. Define a vector $\boldsymbol{c} = {}^T\Lambda$. Let us take an index i_0 which corresponds to p_0 in $A \rtimes_\alpha G$. Since $1 \otimes p_0$ is also minimal in $C(G) \rtimes_\delta G$, we get $c_{i_0} = \Lambda_{i_0} = 1$. □

We end this section with the following proposition. We use the same notations as before.

Proposition 4.22. *Let* $A \subset C(G)$ *be a right coideal and take an eigenvector* $\boldsymbol{c} = {}^T\Lambda$ *for its multiplicity map. If there exists an index* $j \in I$ *with* $c_j = 1$*, then* *the reduced ergodic system* $e_j(A \otimes \mathbb{K}(L^2(G)))e_j$ *is* G*-equivariantly embedded into* $C(G)$*. In particular it also becomes a right coideal of* $C(G)$*.*

Proof. Let p_0 be a minimal projection in $\mathbb{K}(L^2(G))$ associated to the trivial representation. Since the ranks of e_j and p_0 in $C(G) \rtimes_\delta G \cong \mathbb{K}(L^2(G))$ are equal, we have a partial isometry v in $C(G) \rtimes_\delta G$ such that it satisfies $v^*v = e_j$ and $vv^* = p_0$. Note that v belongs to $C(G) \rtimes_\delta G = (C(G) \otimes \mathbb{K}(L^2(G)))^G$. Then we have a desired G-equivariant $*$-homomorphism $e_j(A \otimes \mathbb{K}(L^2(G)))e_j \longrightarrow C(G) \otimes \mathbb{C}p_0$ defined by $a \mapsto vav^*$. ☐

Remark 4.23. Let $\{A, G, \alpha\}$ and $\{B, G, \beta\}$ be two ergodic systems and $\theta : A \longrightarrow B$ be a G-equivariant $*$-homomorphism. Then it must be faithful as we see below. Let φ_A and φ_B be the unique faithful invariant states. Since θ is G-equivariant, $\varphi_B \circ \theta$ is an invariant state. By its uniqueness we obtain $\varphi_A = \varphi_B \circ \theta$. Hence θ is faithful.

5. ELEMENTARY RESULTS FOR $SU_q(2)$

In this section we summarize basic facts for $SU_q(2)$ for readers' convenience. Readers are referred to, for example, [Wo1, MMNNU] for its basic theory and [Po1, Po2, Ko, NM] for the results on the quantum spheres or quantum subgroups in $SU_q(2)$. We adopt the same notation as [MMNNU] in this paper. We treat a real number q in $[-1, 1] \setminus \{0\}$. The smooth function algebra $A(SU_q(2))$ is the universal $*$-algebra generated by four elements x, u, v and y with the following relations

$$ux = qxu, \quad vx = qxv, \quad yu = quy, \quad yv = qvy,$$

$$uv = vu, \quad xy - q^{-1}uv = yx - quv = 1, \quad x^* = y, \quad u^* = -q^{-1}v.$$

Its universal C^*-algebra is denoted by $C(SU_q(2))$. We often use a positive operator $\zeta = -q^{-1}uv$. We make $A(SU_q(2))$ a Hopf $*$-algebra by defining coproduct δ, counit ε and antipode κ as follows

$$\begin{pmatrix} \delta(x) & \delta(u) \\ \delta(v) & \delta(y) \end{pmatrix} = \begin{pmatrix} x \otimes 1 & u \otimes 1 \\ v \otimes 1 & y \otimes 1 \end{pmatrix} \cdot \begin{pmatrix} 1 \otimes x & 1 \otimes u \\ 1 \otimes v & 1 \otimes y \end{pmatrix},$$

$$\begin{pmatrix} \varepsilon(x) & \varepsilon(u) \\ \varepsilon(v) & \varepsilon(y) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

$$\begin{pmatrix} \kappa(x) & \kappa(u) \\ \kappa(v) & \kappa(y) \end{pmatrix} = \begin{pmatrix} y & -qu \\ -q^{-1}v & x \end{pmatrix}.$$

Then the pair $SU_q(2) = (C(SU_q(2)), \delta)$ becomes a compact quantum group and it is often called the *twisted* $SU(2)$ *group*. The maps δ and ε are extended to

$C(SU_q(2))$ norm continuously and the map κ is extended to $C(SU_q(2))$ as a closed operator. Its Woronowicz characters $\{f_z\}_{z\in\mathbb{C}}$ are given by

$$\begin{pmatrix} f_z(x) & f_z(u) \\ f_z(v) & f_z(y) \end{pmatrix} = \begin{pmatrix} |q|^z & 0 \\ 0 & |q|^{-z} \end{pmatrix} \text{ for all } z \in \mathbb{C}.$$

The equivalence classes of irreducible representations $\widehat{SU_q(2)}$ are indexed by spin numbers $\nu \in \frac{1}{2}\mathbb{Z}_{\geq 0}$ and we fix a selection of irreducible representations $\{w(\pi_\nu)\}_\nu$ corresponding to spins ν as follows. The representation space $H_\nu = H_{w(\pi_\nu)}$ is $2\nu+1$-dimensional and fix the orthonormal basis $\{\xi^\nu_r\}_{r\in I_\nu}$ where the index set I_ν is $\{-\nu, -\nu+1, \dots, \nu-1, \nu\}$. Then we define the matrix $w(\pi_\nu) \in \mathbb{B}(H_\nu)\otimes A(SU_q(2))$ by setting $w(\pi_\nu)_{i,j}$ as follows.

(1) Case $i+j \leq 0$, $i \geq j$:

$$x^{-i-j}v^{i-j}q^{(\nu+j)(j-i)}\begin{bmatrix}\nu+i\\ i-j\end{bmatrix}_{q^2}^{\frac{1}{2}}\begin{bmatrix}\nu-j\\ i-j\end{bmatrix}_{q^2}^{\frac{1}{2}} P^{(i-j,-i-j)}_{\nu+j}(\zeta;q^2),$$

(2) Case $i+j \leq 0$, $i \leq j$:

$$x^{-i-j}u^{j-i}q^{(\nu+i)(i-j)}\begin{bmatrix}\nu-i\\ j-i\end{bmatrix}_{q^2}^{\frac{1}{2}}\begin{bmatrix}\nu+j\\ j-i\end{bmatrix}_{q^2}^{\frac{1}{2}} P^{(j-i,-i-j)}_{\nu+i}(\zeta;q^2),$$

(3) Case $i+j \geq 0$, $i \leq j$:

$$q^{(j-i)(j-\nu)}\begin{bmatrix}\nu-i\\ j-i\end{bmatrix}_{q^2}^{\frac{1}{2}}\begin{bmatrix}\nu+j\\ j-i\end{bmatrix}_{q^2}^{\frac{1}{2}} P^{(j-i,i+j)}_{\nu-j}(\zeta;q^2)u^{j-i}y^{i+j},$$

(4) Case $i+j \geq 0$, $i \geq j$:

$$q^{(i-j)(i-\nu)}\begin{bmatrix}\nu+i\\ i-j\end{bmatrix}_{q^2}^{\frac{1}{2}}\begin{bmatrix}\nu-j\\ i-j\end{bmatrix}_{q^2}^{\frac{1}{2}} P^{(i-j,i+j)}_{\nu-i}(\zeta;q^2)v^{i-j}y^{i+j},$$

where we have used the *q-binomial coefficients* and the *little q-Jacobi polynomials*:

$$\begin{bmatrix}m\\ n\end{bmatrix}_q = \frac{(q;q)_m}{(q;q)_n(q;q)_{m-n}}, \quad (t;q)_m = \prod_{s=0}^{m-1}(1-tq^s),$$

$$P^{(\alpha,\beta)}_n(z;q) = \sum_{r\geq 0}\frac{(q^{-n};q)_r(q^{\alpha+\beta+n+1};q)_r}{(q;q)_r(q^{\alpha+1};q)_r}(qz)^r.$$

This yields the following formula

$$w(\pi_\nu)^*_{i,j} = (-q)^{i-j}w(\pi_\nu)_{-i,-j}$$

for all $\pi_\nu \in \widehat{SU_q(2)}$ and $i,j \in I_\nu$. For an integer n, we define the *q-integer* and its factorial by

$$(n)_q = \frac{|q|^n - |q|^{-n}}{|q|-|q|^{-1}}, \quad (n)_q! = (n)_q(n-1)_q\cdots(1)_q\,.$$

We summarize useful formulae as follows.

(1) $f_z(w(\pi_\nu)_{r,s}) = \delta_{r,s}|q|^{-2rz}$,

(2) $\sigma_t^h(w(\pi_\nu)_{r,s}) = |q|^{-2(r+s)it}w(\pi_\nu)_{r,s}$,

(3) $\tau_t(w(\pi_\nu)_{r,s}) = |q|^{-2(r-s)it}w(\pi_\nu)_{r,s}$,

(4) $R(w(\pi_\nu)_{r,s}) = (-1)^{r-s}w(\pi_\nu)_{-s,-r}$,

for all $z \in \mathbb{C}$, $t \in \mathbb{R}$, $\nu \in \frac{1}{2}\mathbb{Z}_{\geq}0$ and $r, s \in I_\nu$. The F-matrix $F_{\pi_\nu} = (\mathrm{id} \otimes f_1)(w(\pi_\nu))$ is equal to $\mathrm{diag}(q^{2\nu}, q^{2\nu-2}, \dots, q^{-(2\nu-2)}, q^{-2\nu})$. Hence we obtain $D_{\pi_\nu} = (2\nu+1)_q$. In the spin $\nu = \frac{1}{2}$ case we have the following matrices

$$\begin{pmatrix} \sigma_t^h(x) & \sigma_t^h(u) \\ \sigma_t^h(v) & \sigma_t^h(y) \end{pmatrix} = \begin{pmatrix} |q|^{2it}x & u \\ v & |q|^{-2it}y \end{pmatrix},$$

$$\begin{pmatrix} \tau_t(x) & \tau_t(u) \\ \tau_t(v) & \tau_t(y) \end{pmatrix} = \begin{pmatrix} x & |q|^{2it}u \\ |q|^{-2it}v & y \end{pmatrix},$$

$$\begin{pmatrix} R(x) & R(u) \\ R(v) & R(y) \end{pmatrix} = \begin{pmatrix} y & -v \\ -u & x \end{pmatrix}.$$

For the calculation about the Haar state, we have

$$h(\zeta^n) = \frac{1-q^2}{1-q^{2(n+1)}} \text{ for all } n \in \mathbb{Z}_{\geq 0}.$$

We recall the embedding of $\mathbb{T}$ into $SU_q(2)$ for $-1 \leq q < 1$. The torus group $\mathbb{T}$ is identified with the set $\{z \in \mathbb{C} \mid |z| = 1\}$ and its cyclic subgroup $\mathbb{T}_m$ ($m \geq 2$) is generated by $\exp(\frac{2\pi\sqrt{-1}}{m})$. For $-1 \leq q < 1$ if $\theta : C(SU_q(2)) \longrightarrow C(\mathbb{T})$ is a restriction map, then it must be as follows.

$$\begin{pmatrix} \theta(x) & \theta(u) \\ \theta(v) & \theta(y) \end{pmatrix} = \begin{pmatrix} z & 0 \\ 0 & \overline{z} \end{pmatrix} \text{ or } \begin{pmatrix} \overline{z} & 0 \\ 0 & z \end{pmatrix},$$

where $z \in \mathbb{T} \subset \mathbb{C}$ is a usual coordinating map. Let $\pi_{\mathbb{T}}$ and $\pi'_{\mathbb{T}}$ be the first and second restriction maps in the above equalities, respectively. $\mathbb{T}$ is called the *maximal torus subgroup* of $SU_q(2)$ and we always treat it with the restriction map $\pi_{\mathbb{T}}$. Note that quotient space $\mathbb{T} \setminus SU_q(2)$ or $\mathbb{T}_m \setminus SU_q(2)$ do not depend on the choice of $\pi_{\mathbb{T}}$ or $\pi'_{\mathbb{T}}$. The quotient space $C(\mathbb{T} \setminus SU_q(2))$ is often called the *canonical homogeneous sphere*, which is generated by $\{w(\pi_1)_{0,r}\}_{r \in I_1}$. The quotient space $C(\mathbb{T}_2 \setminus SU_q(2))$ is denoted by $C(SO_q(3))$ which is also a compact quantum group by restricting the coproduct δ. We can easily see $C(SO_q(3))$ has the spectral pattern: $\oplus_{k \in \mathbb{Z}_{\geq 0}}(2k+1)\pi_k$ and hence it does not depend on positivity or negativity of the parameter q. In fact we have an isomorphism $\Xi_q : C(SO_q(3)) \longrightarrow C(SO_{-q}(3))$ as compact quantum groups defined by the following equality, where 2 by 2 matrices $\begin{pmatrix} x & u \\ v & y \end{pmatrix}$ and $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ are fundamental representations of $SU_q(2)$ and $SU_{-q}(2)$ respectively:

$$\begin{pmatrix} \Xi_q(x^2) & \sqrt{1+q^2}\Xi_q(xu) & \Xi_q(u^2) \\ \sqrt{1+q^2}\Xi_q(xv) & \Xi_q(1+(q+q^{-1})uv) & \sqrt{1+q^2}\Xi_q(uy) \\ \Xi_q(v^2) & \sqrt{1+q^2}\Xi_q(vy) & \Xi_q(y^2) \end{pmatrix}$$
$$= \begin{pmatrix} a^2 & -i\sqrt{1+q^2}ab & b^2 \\ i\sqrt{1+q^2}ac & 1-(q+q^{-1})bc & i\sqrt{1+q^2}bd \\ c^2 & -i\sqrt{1+q^2}cd & d^2 \end{pmatrix}.$$

With this isomorphism, we often identify $SO_q(3)$ with $SO_{-q}(3)$.

Now we recall classical results on closed subgroups of $SU(2)$. Let us use the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

They give an orthonormal basis for a three dimensional real vector space $\mathbb{R}^3 = \mathbb{R}\sigma_1 + \mathbb{R}\sigma_2 + \mathbb{R}\sigma_3$. We define a double covering $\pi_1 : SU(2) \longrightarrow SO(3)$ by the adjoint action $\pi_1(g)(a) = gag^{-1}$ for all $g \in SU(2)$ and $a \in \mathbb{R}^3$. For a subgroup $H \subset SO(3)$ we write H^* for its inverse image by π_1 and such a subgroup is called a *binary subgroup.* A symmetric group and alternative groups A_4, S_4 and A_5 are embedded into $SO(3)$ as the *tetrahedral group*, the *octahedral group* and the *icosahedral group*, respectively. The group $D_m\,(2 \leq m \leq \infty)$ is a *dihedral group*, and $\mathbb{T}$ and $\mathbb{T}_m\,(2 \leq m)$ are the ordinary torus and the cyclic groups of order m. In $SU(2)$ all the closed subgroups are conjugate to one of the *trivial group* 1, the *cyclic group* $\mathbb{T}_n$, $SU(2)$, the maximal torus $\mathbb{T}$, the *binary dihedral group* $D^*_m\,(2 \leq m \leq \infty)$, the *binary tetrahedral group* A^*_4, the *binary octahedral group* S^*_4 and the *binary icosahedral group* A^*_5. Since we need explicit embedding of $\mathbb{T}_n$ and D_n in the final section, we prepare a few notations. Let us define rotation matrices

$$r^{12}(\theta) = \begin{pmatrix} e^{-i\frac{\theta}{2}} & 0 \\ 0 & e^{i\frac{\theta}{2}} \end{pmatrix}, r^{23}(\theta) = \begin{pmatrix} \cos\frac{\theta}{2} & -i\sin\frac{\theta}{2} \\ -i\sin\frac{\theta}{2} & \cos\frac{\theta}{2} \end{pmatrix}, r^{13}(\theta) = \begin{pmatrix} \cos\frac{\theta}{2} & \sin\frac{\theta}{2} \\ -\sin\frac{\theta}{2} & \cos\frac{\theta}{2} \end{pmatrix},$$

where $\pi_1(r^{ij}(\theta))$ gives the rotation of angle θ in σ_i-σ_j plane. The cyclic group $\mathbb{T}_n$ is specified by two angles χ and ψ and denoted by $\mathbb{T}^{\chi,\psi}_n$. It consists of rotations of the angle $\frac{2\pi}{n}$ around the axis; $-\cos\chi\sin\psi\sigma_1 - \sin\chi\sin\psi\sigma_2 + \cos\psi\sigma_3$. The dihedral group D_n is specified by three angles ϕ, χ and ψ and denoted by $D^{\phi,\chi,\psi}_n$. It is generated by $\mathbb{T}^{\chi,\psi}_n$ and the rotation of the angle π around the axis; $\cos\phi\cos\chi\cos\psi\sigma_1 + \sin\phi\sin\chi\sigma_2 + \cos\phi\sin\psi\sigma_3$. By definition we have $\mathbb{T}^{\chi,\psi}_n = \mathrm{Ad}(\pi_1(r^{12}(\chi)r^{13}(\psi)))(\mathbb{T}^{0,0}_n)$ and $D^{\phi,\chi,\psi}_n = \mathrm{Ad}(\pi_1(r^{12}(\chi)r^{13}(\psi)r^{12}(\phi)))(D^{0,0,0}_n)$.

In [Po2] he has classified all the quantum subgroups of $SU_q(2)$ for $-1 \leq q < 1$. We summarize it for readers' convenience and analysis on $SU_{-1}(2)$ later.

(1) $0 < |q| < 1$ case. Its quantum subgroup is one of 1, $\mathbb{T}_m\,(m \geq 2)$, $\mathbb{T}$, and $SU_q(2)$. For $\mathbb{T}_m\,(m \geq 2)$ and $\mathbb{T}$, their restriction map is $\pi_{\mathbb{T}}$ or $\pi'_{\mathbb{T}}$.

(2) $q=-1$ case. For $g=\begin{pmatrix}\alpha & -\overline{\gamma}\\ \gamma & \overline{\alpha}\end{pmatrix}\in SU(2)$, define the $*$-homomorphism $\upsilon_g : C(SU_{-1}(2)) \longrightarrow \mathbb{B}(\mathbb{C}^2)$ by

$$\begin{pmatrix}\upsilon_g(x) & \upsilon_g(u)\\ \upsilon_g(v) & \upsilon_g(y)\end{pmatrix} = \begin{pmatrix}\alpha\sigma_1 & \overline{\gamma}\sigma_2\\ \gamma\sigma_2 & \overline{\alpha}\sigma_1\end{pmatrix}.$$

Now we define the irreducible representation τ_g of $C(SU_{-1}(2))$ as a C^*-algebra. They give all irreducible representations of $C(SU_{-1}(2))$.

(i) $\alpha, \gamma \neq 0$ case. $\tau_g = \upsilon_g$

(ii) $\alpha = 0$ case. $\tau_g : C(SU_{-1}(2)) \longrightarrow \mathbb{C}$ is

$$\begin{pmatrix}\tau_g(x) & \tau_g(u)\\ \tau_g(v) & \tau_g(y)\end{pmatrix} = \begin{pmatrix}0 & \overline{\gamma}\\ \gamma & 0\end{pmatrix}.$$

(iii) $\gamma = 0$ case. $\tau_g : C(SU_{-1}(2)) \longrightarrow \mathbb{C}$ is

$$\begin{pmatrix}\tau_g(x) & \tau_g(u)\\ \tau_g(v) & \tau_g(y)\end{pmatrix} = \begin{pmatrix}\alpha & 0\\ 0 & \overline{\alpha}\end{pmatrix}.$$

For a compact subset $Z \subset SU(2)$ we consider the direct sum representation $\pi_Z = \oplus_{g\in Z}\pi_g$. In [Po2, Proposition 2.4.], it is characterized when $\pi_Z : C(SU_{-1}(2)) \longrightarrow \pi_Z(C(SU_{-1}(2)))$ gives a restriction map to a quantum subgroup. We write $C(G_Z) = \pi_Z(C(SU_{-1}(2)))$. Before a summary we study the embedding $\mathbb{T}_n \subset SU_{-1}(2)$ a little. For $a \in C(SU_{-1}(2))$ and $g \in \mathbb{T}_n$ we write simply $a(g)$ for $r(a)(g)$ where r is a restriction map. Let g be a generator of $\mathbb{T}_n$. Since we have $xv = -vx$, $x(g) = 0$ or $v(g) = 0$. We consider $v(g) \neq 0$. By the definition of the restriction map we have $\begin{pmatrix}x(g^2) & u(g^2)\\ v(g^2) & y(g^2)\end{pmatrix} = \begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}$. This shows n must be 2. We denote the restriction by π_{D_1}. Hence if $n \geq 3$, the embedding of $\mathbb{T}_n$ is unique and if $n = 2$, two embedding arises. Since in this paper notations $\mathbb{T}$ and $\mathbb{T}_n$ are used for the maximal torus and its cyclic subgroups, we prepare the notation D_1 for a subgroup of order 2 which is embedded by π_{D_1}. Of course subgroups D_1 and $\mathbb{T}_2$ are isomorphic, however, right coideals by them are not isomorphic.

(a) $C(G_Z)$ is an abelian C^*-algebra case (that is, G_Z is an ordinary group), then G_Z is (isomorphic to) one of trivial group 1, the maximal torus $\mathbb{T}$, the cyclic groups $\mathbb{T}_n(\subset \mathbb{T})(n \geq 2)$, D_1 and dihedral groups $D_n(2 \leq n \leq \infty)$ containing $\mathbb{T}_n$. (This isomorphism gives the conjugation β_z^L on right coideals by them.) About $D_n(1 \leq n \leq \infty)$ its corresponding subset Z_{D_n} is $\{r^{12}(\frac{2\pi k}{n}), r^{12}(\frac{2\pi k}{n})r^{23}(\pi) \mid 0 \leq k \leq n-1\}$. Note that Z_{D_n} is not a subgroup in $SU(2)$ if odd n. If G_Z is not $\mathbb{T}_n$ or D_n (odd n), Z is a binary subgroup and it shows the spectral pattern of the right coideal $C(G_Z \setminus SU_{-1}(2))$ consists of integer spins. If G_Z is, then it also has half integer spin parts.

(b) If $C(G_Z)$ is a non-abelian C^*-algebra, Z is one of binary subgroups whose image by π_1 is listed in [Po2, p. 11]. Note that it depends on the embedding of a closed subgroup into $SO(3)$. Then it is known that $C(G_Z \setminus SU_{-1}(2)) \subset C(SO_{-1}(3))$ and $\Xi_{-1}(C(G_Z \setminus SU_{-1}(2))) = C(H \setminus SO_1(3))$.

The quantum universal enveloping algebra $U_q(su_2)$ is generated by four elements k, k^{-1}, e and f which satisfy the following relations:

$$kk^{-1} = 1 = k^{-1}k,$$
$$kek^{-1} = qe,\ kfk^{-1} = q^{-1}f,$$
$$ef - fe = \frac{k^2 - k^{-2}}{q - q^{-1}}.$$

$U_q(su_2)$ is realized as a Hopf $*$-subalgebra of the algebraic functional space $A(SU_q(2))^*$ as follows.

(1) $q > 0$ case.

$$k^{\pm 1}(w(\pi_\nu)_{r,s}) = q^{\mp r}\delta_{r,s},$$
$$e(w(\pi_\nu)_{r,s}) = \delta_{r+1,s}\sqrt{(\nu + s)_q(\nu - s + 1)_q},$$
$$f(w(\pi_\nu)_{r,s}) = \delta_{r-1,s}\sqrt{(\nu - s)_q(\nu + s + 1)_q}.$$

(2) $q < 0$ case.

$$k^{\pm 1}(w(\pi_\nu)_{r,s}) = q^{\mp r}\delta_{r,s},$$
$$e(w(\pi_\nu)_{r,s}) = \delta_{r+1,s}\sqrt{-1}^{-2\nu+1}\sqrt{(\nu + s)_q(\nu - s + 1)_q},$$
$$f(w(\pi_\nu)_{r,s}) = \delta_{r-1,s}\sqrt{-1}^{-2\nu+1}\sqrt{(\nu - s)_q(\nu + s + 1)_q},$$

where $q^n = \sqrt{-1}^{2n}(-q)^n$ for a half integer n. The Hopf $*$-algebra structure of $U_q(su_2)$ is given by

(1) $k^* = k,\ e^* = f,\ f^* = e,$
(2) $\hat{\delta}(k) = k \otimes k,\ \hat{\delta}(e) = e \otimes k + k^{-1} \otimes e,\ \hat{\delta}(f) = f \otimes k + k^{-1} \otimes f,$
(3) $\hat{\kappa}(k) = k^{-1},\ \hat{\kappa}(e) = -qe,\ \hat{\kappa}(f) = -q^{-1}f.$

Hence $U_q(su_2)$ is a Hopf $*$-subalgebra of $A(SU_q(2))$. For a smooth corepresentation of $A(SU_q(2))$ $\alpha : K \longrightarrow K \otimes A(SU_q(2))$, we prepare $U_q(su_2)$-module structure on K by $\theta \cdot \xi = (\mathrm{id} \otimes \theta)(w\xi)$ for all $\theta \in U_q(su_2)$ and $\xi \in K$. This representation of $U_q(su_2)$ is called a differential representation of a corepresentation (α, K). Consider irreducible representation $(w(\pi_\nu), H_\nu)$ and we have the following formulae about its $U_q(su_2)$-module structure.

(1) $q > 0$ case.

$$k^{\pm 1} \cdot \xi^\nu_r = q^{\mp r}\xi^\nu_r\ ,$$
$$e \cdot \xi^\nu_r = \sqrt{(\nu + r)_q(\nu - r + 1)_q}\,\xi^\nu_{r-1}\ ,$$
$$f \cdot \xi^\nu_r = \sqrt{(\nu - r)_q(\nu + r + 1)_q}\,\xi^\nu_{r+1}\ .$$

(2) $q < 0$ case.

$$\begin{aligned} k^{\pm 1} \cdot \xi^\nu_r &= q^{\mp r} \xi^\nu_r \ , \\ e \cdot \xi^\nu_r &= \sqrt{-1}^{-2\nu+1} \sqrt{(\nu+r)_q (\nu - r + 1)_q}\, \xi^\nu_{r-1} \ , \\ f \cdot \xi^\nu_r &= \sqrt{-1}^{-2\nu+1} \sqrt{(\nu-r)_q (\nu + r + 1)_q}\, \xi^\nu_{r+1} \ . \end{aligned}$$

for all $r \in I_\nu$. For a half integer n a vector $\xi \in K$ is called a *highest weight vector* of *weight* n if it satisfies $k \cdot \xi = q^n \xi$ and $e \cdot \xi = 0$. For example, the vector $\xi^\nu_{-\nu} \in H_\nu$ is a highest weight vector of weight ν. It is well-known that a tensor product $U_q(su_2)$-module $H_\mu \otimes H_\nu$ is isomorphic to the direct sum $U_q(su_2)$-module $\bigoplus_{|\mu-\nu| \leq \ell \leq \mu+\nu} H_\ell$, where ℓ runs through half integers. If we want to make a highest weight vector of H_ℓ from those of H_μ and H_ν, the following well-known lemmas are useful.

Lemma 5.1. *For positive q, consider the tensor product $U_q(su_2)$-module $H_\mu \otimes H_\nu$. Let ℓ be an integer with $|\mu - \nu| \leq \ell \leq \mu + \nu$. Define the coefficients $(C^\ell_{\mu,\nu})_r$ for $0 \leq r \leq \mu + \nu - \ell$ by*

$$(C^\ell_{\mu,\nu})_r = q^{-\frac{1}{2}(\ell+1)(\mu+\nu-\ell)} (-q^{\ell+1})^r \prod_{t=1}^{r} \sqrt{\frac{(\mu+\nu-\ell+1-t)_q (\mu-\nu+\ell+t)_q}{(t)_q (2\nu - t + 1)_q}}.$$

Then a vector $\eta^\ell = \sum_{r=0}^{\mu+\nu-\ell} (C^\ell_{\mu,\nu})_r\, \xi^\mu_{-\ell+\nu-r} \otimes \xi^\nu_{-\nu+r}$ is a highest weight vector of weight ℓ in $H_\mu \otimes H_\nu$.

Proof. The action of $U_q(su_2)$ on $H_\mu \otimes H_\nu$ is given via coproduct. With this, we can easily justify $k \cdot \eta^\ell = q^{-\ell} \eta^\ell$ and $e \cdot \eta^\ell = 0$. □

We state a negative q version of Lemma 5.1 as follows.

Lemma 5.2. *For negative q, consider the tensor product $U_q(su_2)$-module $H_\mu \otimes H_\nu$. Let ℓ be an integer with $|\mu - \nu| \leq \ell \leq \mu + \nu$ and ξ^μ and ξ^ν be vectors of copy of π_μ and π_ν respectively. Define the coefficients $(C^\ell_{\mu,\nu})_r$ for $0 \leq r \leq \mu + \nu - \ell$ by*

$$(C^\ell_{\mu,\nu})_r = q_0^{-\frac{1}{2}(\ell+1)(\mu+\nu-\ell)} (-1)^{(-\mu+\nu+\ell)r} q_0^{r(\ell+1)} \prod_{t=1}^{r} \sqrt{\frac{(\mu+\nu-\ell+1-t)_q (\mu-\nu+\ell+t)_q}{(t)_q (2\nu+1-t)_q}}.$$

Then a vector $\eta^\ell = \sum_{r=0}^{\mu+\nu-\ell} (C^\ell_{\mu,\nu})_r\, \xi^\mu_{-\ell+\nu-r} \otimes \xi^\nu_{-\nu+r}$ is a highest weight vector of weight ℓ in $H_\mu \otimes H_\nu$.

Let Y_μ be a linear space of π_μ-eigenvectors of $C(SU_q(2))$. We prepare the notation of eigenvectors $\boldsymbol{w}^\mu_r := (w(\pi_\mu)_{r,t})_{t \in I_\mu}$ for $r \in I_\mu$. They give an orthonormal basis of Y_μ. Let us consider a covariant system $(A, SU_q(2), \alpha)$. For its eigenvector spaces $\{X_\nu\}_{\nu \in \frac{1}{2}\mathbb{Z}_{\geq 0}}$ we define the *product of eigenvectors* $\Psi_\ell : X_\mu \times X_\nu \longrightarrow X_\ell$ by using Lemma 5.1:

$$\Psi_\ell(\xi^\mu, \xi^\nu)_{-\ell} = \sum_{r=0}^{\mu+\nu-\ell} (C^\ell_{\mu,\nu})_r\, \xi^\mu_{-\ell+\nu-r}\, \xi^\nu_{-\nu+r}$$

for all eigenvectors ξ^μ and ξ^ν, where the coefficients $(C^\ell_{\mu,\nu})_r$ are given in Lemma 5.1 and Lemma 5.2. In [Po1], he has classified ergodic systems $\{A, SU_q(2), \alpha\}$ with $\dim A_{\pi_1} = 1$ and $A = C^*(A_{\pi_1})$ for $|q| < 1$. It is uniquely determined up to $SU_q(2)$-isomorphism and is called the *quantum sphere*. We summarize his classification. Let X_{π_1} be the π_1-eigenvector space. Since this is one-dimensional, we can take the (unique) π_1-eigenvector ξ with the following properties:

$$T\xi_\lambda = \xi_\lambda, \quad (\xi_\lambda, \xi_\lambda) = 1, \quad \Psi_1(\xi_\lambda, \xi_\lambda) = \lambda\, \xi_\lambda,$$

where λ is a non-negative real constant. From the second assumption the C^*-algebra A is generated by $(\xi_\lambda)_{-1}$, $(\xi_\lambda)_0$ and $(\xi_\lambda)_1$. Write λ_0 for $(q^{-1} - q)^{-1}\lambda$. For $n \in \mathbb{Z}_{\geq 1}$ we define a positive number $c(n) = \frac{q^{n+1}+q^{-n-1}}{\sqrt{(n)_q(n+2)_q}}$. Then the classification is done as follows.

Case 1: If $\lambda_0 > 1$, then there exists $n \in \mathbb{Z}_{\geq 1}$ such that $\lambda = c(n)$ and we obtain a G-isomorphism $A \cong \mathrm{End}(H_{\pi_{\frac{n}{2}}})$. In this case the spectral pattern of A is $\pi_0 \oplus \pi_1 \oplus \cdots \oplus \pi_n$.

Case 2: If $0 \leq \lambda_0 \leq 1$ and $q > 0$, then the map $\xi_\lambda = ((\xi_\lambda)_{-1}, (\xi_\lambda)_0, (\xi_\lambda)_1) \mapsto \left(q^{\frac{1}{2}}\sqrt{\frac{1-\lambda_0^2}{(2)_q}}, \lambda_0, -q^{-\frac{1}{2}}\sqrt{\frac{1-\lambda_0^2}{(2)_q}}\right) \cdot w(\pi_1)$ gives a G-equivariant embedding $A \hookrightarrow C(SU_q(2))$. In this case the spectral pattern of A is $\oplus_{\ell \in \mathbb{Z}_{\geq 0}} \pi_\ell$.

Case 3: If $0 \leq \lambda_0 \leq 1$ and $q < 0$, then the map $\xi_\lambda = ((\xi_\lambda)_{-1}, (\xi_\lambda)_0, (\xi_\lambda)_1) \mapsto \left((-q)^{\frac{1}{2}}\sqrt{\frac{1-\lambda_0^2}{(2)_q}}, \lambda_0, (-q)^{-\frac{1}{2}}\sqrt{\frac{1-\lambda_0^2}{(2)_q}}\right) \cdot w(\pi_1)$ gives a G-equivariant embedding $A \hookrightarrow C(SU_q(2))$. In this case the spectral pattern of A is $\oplus_{\ell \in \mathbb{Z}_{\geq 0}} \pi_\ell$.

In this paper we use the notation $C(S^2_{q,\lambda})$ for the right coideal which is defined in the above case 2 and 3. In the proof of Lemma 8.2 we obtain a π_1-eigenvector of $C(\mathbb{T}^{0,\psi} \setminus SO(3))$. Comparing it with the above embedding, the quantum sphere $C(S^2_{q,\lambda})$ is considered a q-deformation of $C(\mathbb{T}^{0,\psi} \setminus SO(3))$ with the parameter $\lambda_0 = \cos\psi$. If $\lambda_0 = 1$, $C(S_{q,(q^{-1}-q)^{-1}})$ becomes the canonical homogeneous sphere $C(\mathbb{T} \setminus SU_q(2))$. Let β^L_z be a left action of the maximal torus on $C(SU_q(2))$ defined by $(\mathrm{ev}_z \circ \pi_\mathbb{T} \otimes \mathrm{id}) \circ \delta$ for all $z \in \mathbb{T}$. It satisfies $(\beta^L_z \otimes \mathrm{id}) \circ \delta = \delta \circ \beta^L_z$, that is, β^L_z gives an $SU_q(2)$-isomorphism. Note that all embeddings of quantum spheres into $C(SU_q(2))$ are obtained by rotations of β^L for the above given embedding. Similarly we define the right action β^R of the maximal torus by $\beta^R = (\mathrm{id} \otimes \pi_\mathbb{T}) \circ \delta$.

We show the list of all the connected graphs of norm 2 in the Appendix. For its classification readers are referred to [GHJ, Lemma 1.4.1]. The labels except for A'_m correspond to closed subgroups of $SU(2)$ or $SU_{-1}(2)$ by their McKay diagrams about fundamental representation $\pi_{\frac{1}{2}}$ or multiplicity diagrams of the right coideals obtained by quotients. We will see later that type A'_m does not occur even if we investigate right coideals.

Remark 5.3. We discuss the absence of A'_m $(3 \leq m \leq \infty)$. Let $C(G)$ be a compact quantum group which has an irreducible unitary representation w generating $R(G)$ (that is, $C(G)$ is a compact matrix pseudogroup [Wo3]). We consider a compact quantum subgroup $C(H)$ whose McKay diagram about w is of type A'_m $(m \geq 3)$. We simply denote the restriction of w onto H by $w|_H$. Now vertices in A'_m corresponds to the irreducible representations of H and let $\rho_0, \rho_1 \dots$ be the irreducible representations from the left side in the Figure 14, 15. Since the entries of the Perron-Frobenius eigenvector are equal to the dimensions of the corresponding irreducible modules, $\widehat{H}$ must be a group. By definition of McKay diagram, $w|_H \cdot \rho_0 = \rho_0 + \rho_1$ holds. Hence we have $w|_H = 0 + \rho_1\rho_0^{-1}$ where 0 is the trivial representation. Then it yields $w|_H \cdot \rho_1 = \rho_1 + \rho_1\rho_0^{-1}\rho_1$, and this shows there must exist a single loop at ρ_1. Hence the McKay diagram of type A'_m $(m \geq 3)$ does not appear.

6. CLASSIFICATION OF RIGHT COIDEALS OF $C(SU_q(2))$: $0 < q < 1$ CASE

In $q = 1$ case the main theorem of [Wa2, p. 309] says that any ergodic system of $SU(2)$ is an induced system $\{\mathrm{End}(W), H, \beta\}$ where H is a closed subgroup of $SU(2)$. A closed subgroup H is conjugate to one of $1, \mathbb{T}_n\,(n \geq 2), \mathbb{T}, SU(2), D^*_n\,(n \geq 2), D^*_\infty, A^*_4, S^*_4$ and A^*_5. Their corresponding multiplicity diagrams are exactly their McKay diagrams. In $0 < q < 1$ case there are a little quantum subgroups of $SU_q(2)$ [Po2, Theorem 2.1.]. So we are interested in the absence of some multiplicity diagrams in the table. Finally we obtain the following results for right coideals.

Theorem 6.1. *Let $A \subset C(SU_q(2))$ be a right coideal. Then its multiplicity diagram is one of type $1, \mathbb{T}_n\,(n \geq 2), \mathbb{T}, SU(2)$ and D^*_∞. If it is of type $\mathbb{T}$, then it is one of the quantum spheres. Otherwise it is unique up to conjugation by β^L.*

Now we start the proof of Theorem 6.1. First we show that there do not exist right coideals whose multiplicity diagrams are of type $D^*_n\,(n \geq 2), D^*_\infty, A^*_4, S^*_4$ and A^*_5. Second we classify right coideals whose multiplicity diagrams are of type $1, \mathbb{T}_n\,(n \geq 2), \mathbb{T}, SU(2), D^*_\infty$. In all cases we make use of products of eigenvectors.

(I) A^*_4 case. A has a spectral pattern $\pi_0 \oplus \pi_3 \oplus \pi_4 \oplus 2\pi_6 \oplus \pi_7 \oplus \cdots$. Like the discussion of [Wa2, p. 321], we focus on spectral gaps: π_1, π_2 and π_5. We will soon notice the importance of using the both of even and odd spin. Let $\eta = (\eta_r)_{r \in I_3}$ be a self-conjugate π_3-eigenvector of A. Take scalars $\{c_r\}_{r \in I_3}$ such that $\eta_r = \sum_{s \in I_3} c_s w(\pi_3)_{s,r}$ for all $r \in I_3$. Applying Lemma 5.1, we obtain the following highest weight vectors

$$\Psi_5(\eta, \eta)_{-5} = q^{-3}\eta_{-2}\eta_{-3} - q^3\eta_{-3}\eta_{-2}\,,$$

$$\begin{aligned}\Psi_2(\eta,\eta)_{-2} = {} & q^{-6}\eta_1\eta_{-3} - q^{-3}\sqrt{\frac{(4)_q(3)_q}{(6)_q}}\eta_0\eta_{-2} + \frac{(4)_q(3)_q}{\sqrt{(6)_q(5)_q(2)_q}}\eta_{-1}\eta_{-1} \\ & - q^{3}\sqrt{\frac{(4)_q(3)_q}{(6)_q}}\eta_{-2}\eta_0 + q^{6}\eta_{-3}\eta_1\,,\end{aligned}$$

$$\begin{aligned}\Psi_1(\eta,\eta)_{-1} = {} & q^{-5}\eta_2\eta_{-3} - q^{-3}\sqrt{\frac{(5)_q(2)_q}{(6)_q}}\eta_1\eta_{-2} \\ & + q^{-1}\sqrt{\frac{(4)_q(3)_q}{(6)_q}}\eta_0\eta_{-1} - q\sqrt{\frac{(4)_q(3)_q}{(6)_q}}\eta_{-1}\eta_0 \\ & + q^{3}\sqrt{\frac{(5)_q(2)_q}{(6)_q}}\eta_{-2}\eta_1 - q^{5}\eta_{-3}\eta_2\,.\end{aligned}$$

They are actually 0 vectors because of the absence of spectra. Applying the lowering operator $f \in U_q(su_2)$ to these vectors, we obtain

$$f\cdot\Psi_5(\eta,\eta)_{-5} = \sqrt{(5)_q(2)_q}\eta_{-1}\eta_{-3} + \sqrt{(6)_q}(q^{-5}-q^{5})\eta_{-2}^2 - \sqrt{(5)_q(2)_q}\eta_{-3}\eta_{-1}\,,$$

$$\begin{aligned}f^2\cdot\Psi_5(\eta,\eta)_{-5} = {} & q^{3}\sqrt{(4)_q(3)_q}\eta_0\eta_{-3} + (q^{-3}+q^{-1}-q^{7})\sqrt{(6)_q(5)_q(2)_q}\eta_{-1}\eta_{-2} \\ & + (q^{-7}-q-q^{3})\sqrt{(6)_q(5)_q(2)_q}\eta_{-2}\eta_{-1} + q^{-3}\sqrt{(4)_q(3)_q}\eta_{-3}\eta_0\end{aligned}$$

and

$$\begin{aligned}\sqrt{(6)_q}f\cdot\Psi_1(\eta,\eta)_{-1} = {} & q^{-2}(6)_q\eta_3\eta_{-3} + \big(-q^{-1}(5)_q(2)_q + q^{-3}(6)_q\big)\eta_2\eta_{-2} \\ & + \big((4)_q(3)_q - q^{-2}(5)_q(2)_q\big)\eta_1\eta_{-1} + (q^{-1}-q)(4)_q(3)_q\eta_0^2 \\ & + \big(q^{2}(5)_q(2)_q - (4)_q(3)_q\big)\eta_{-1}\eta_1 + \big(q(5)_q(2)_q - q^{3}(6)_q\big)\eta_{-2}\eta_2 - \\ & - q^{2}(6)_q\eta_{-3}\eta_3\,.\end{aligned}$$

Recall a surjective $*$-homomorphism $\pi_{\mathbb{T}} : C(SU_q(2)) \longrightarrow C(\mathbb{T})$. It sends $w(\pi_\nu)_{r,s}$ to $\delta_{r,s}z^{-2r}$ for $\nu \in \frac{1}{2}\mathbb{Z}_{\geq 0}$ and $r,s \in I_\nu$. Then we have $\pi_{\mathbb{T}}(\eta_r) = z^{-2r}c_r$ for $r \in I_3$. From $0 = \pi_{\mathbb{T}}(f\cdot\Psi_5(\eta,\eta)_{-5})$, we obtain $c_{-2} = 0$. Then from $0 = \pi_{\mathbb{T}}(\Psi_1(\eta,\eta)_{-1}) = \pi_{\mathbb{T}}(f^2\cdot\Psi_5(\eta,\eta)_{-5})$, we have $c_{-3}c_0 = c_{-1}c_0 = 0$. If c_0 is not 0, then c_{-3} and c_{-1} are equal to 0. Hence $\{c_r\}_{r\in I_3}$ are all zero except for $r = 0$. From $0 = \pi_{\mathbb{T}}(f\cdot\Psi_1(\eta,\eta)_{-1})$ we have $c_0 = 0$, this is contradiction. Therefore c_0 must be 0. Then the last equality deduces to $(6)_q c_3 c_{-3} = (5)_q(2)_q c_1 c_{-1}$. Because of the self-conjugacy of η we have $c_{-3} = -q^3\overline{c_3}$ and $c_{-1} = -q\overline{c_1}$. Then we have

$$(6)_q|c_3|^2 = q^{-2}(5)_q(2)_q|c_1|^2\,.$$

From the second one we get

$$(q^{-6}+q^{6})c_1c_{-3}+\frac{(4)_q(3)_q}{\sqrt{(6)_q(5)_q(2)_q}}c_{-1}^2=0\,.$$

We can easily see that there does not exist a solution of the above two equalities except for the case $c_3 = c_1 = 0$. Therefore we have proved no existence of right coideal of type A_4^*.

Remark 6.2. If we consider the $q = 1$ or $q = -1$ case, then products of integer spin eigenvectors to odd integer spin eigenvectors become a trivial zero map. Therefore arguments in the above and below are non-trivial only in the case of $|q| \neq 1$.

(II) S_4^* case. A has a spectral pattern $\pi_0 \oplus \pi_4 \oplus \pi_6 \oplus \pi_8 \oplus \pi_9 \oplus \pi_{10} \oplus \cdots$. There are spectral gaps of π_ν for $\nu = 1, 2, 3, 5, 7$. Let $\eta = (\eta_s)_{s\in I_4} = \sum_{s\in I_4} c_s \boldsymbol{w}_s^4$ be a non zero self-conjugate π_4-eigenvector of A. We derive contradiction by showing the coefficients c_s are all zero. Let us consider eigenvectors

$$\begin{aligned}
\Psi_7(\eta,\eta)_{-7} =& q^{-4}\eta_{-3}\eta_{-4} - q^{4}\eta_{-4}\eta_{-3}\,,\\
\Psi_5(\eta,\eta)_{-5} =& q^{-9}\eta_{-1}\eta_{-4} - q^{-3}\sqrt{\frac{(6)_q(3)_q}{(8)_q}}\eta_{-2}\eta_{-3} + q^{3}\sqrt{\frac{(6)_q(3)_q}{(8)_q}}\eta_{-3}\eta_{-2}\\
&- q^{9}\eta_{-4}\eta_{-1},\\
\Psi_3(\eta,\eta)_{-3} =& q^{-10}\eta_{1}\eta_{-4} - q^{-6}\sqrt{\frac{(5)_q(4)_q}{(8)_q}}\eta_{0}\eta_{-3} + q^{-2}\frac{(5)_q(4)_q}{\sqrt{(8)_q(7)_q(2)_q}}\eta_{-1}\eta_{-2}\\
&- q^{2}\frac{(5)_q(4)_q}{\sqrt{(8)_q(7)_q(2)_q}}\eta_{-2}\eta_{-1} + q^{6}\sqrt{\frac{(5)_q(4)_q}{(8)_q}}\eta_{-3}\eta_{0} - q^{10}\eta_{-4}\eta_{1},\\
\Psi_2(\eta,\eta)_{-2} =& q^{-9}\eta_{2}\eta_{-4} - q^{-6}\sqrt{\frac{(6)_q(3)_q}{(8)_q}}\eta_{1}\eta_{-3}\\
&+ q^{-3}\sqrt{\frac{(6)_q(5)_q(4)_q(3)_q}{(8)_q(7)_q(2)_q}}\eta_{0}\eta_{-2} - \frac{(5)_q(4)_q}{(8)_q(7)_q(2)_q}\eta_{-1}^2\\
&+ q^{3}\sqrt{\frac{(6)_q(5)_q(4)_q(3)_q}{(8)_q(7)_q(2)_q}}\eta_{-2}\eta_{0} - q^{6}\sqrt{\frac{(6)_q(3)_q}{(8)_q}}\eta_{1}\eta_{-3} + q^{9}\eta_{-4}\eta_{2},
\end{aligned}$$

$$\begin{aligned}
\Psi_1(\eta,\eta)_{-1} =& q^{-7}\eta_{3}\eta_{-4} - q^{-5}\sqrt{\frac{(7)_q(2)_q}{(8)_q}}\eta_{2}\eta_{-3} + q^{-3}\sqrt{\frac{(6)_q(3)_q}{(8)_q}}\eta_{1}\eta_{-2}\\
&- q^{-1}\sqrt{\frac{(5)_q(4)_q}{(8)_q}}\eta_{0}\eta_{-1} + q\sqrt{\frac{(5)_q(4)_q}{(8)_q}}\eta_{-1}\eta_{0} - q^{3}\sqrt{\frac{(6)_q(3)_q}{(8)_q}}\eta_{-2}\eta_{1}
\end{aligned}$$

$$+ q^5\sqrt{\frac{(7)_q(2)_q}{(8)_q}}\eta_{-3}\eta_2 - q^7\eta_{-4}\eta_3.$$

They are in fact 0 vectors by the gap at each spectrum. Applying lowering operator f to these vectors, we get

$$f\cdot\Psi_7(\eta,\eta)_{-7} = \sqrt{(7)_q(2)_q}\eta_{-2}\eta_{-4} + (q^{-7}-q^7)\sqrt{(8)_q}\eta_{-3}^2 - \sqrt{(7)_q(2)_q}\eta_{-4}\eta_{-2},$$

$$\begin{aligned}
f\cdot\Psi_5(\eta,\eta)_{-5} = {} & q^{-5}\sqrt{(5)_q(4)_q}\eta_0\eta_{-4} + \frac{q^{-10}(8)_q-(6)_q(3)_q}{\sqrt{(8)_q}}\eta_{-1}\eta_{-3} \\
& + (q^5-q^{-5})\sqrt{\frac{(7)_q(6)_q(3)_q(2)_q}{(8)_q}}\eta_{-2}\eta_{-2} + \frac{-q^{10}(8)_q+(6)_q(3)_q}{\sqrt{(8)_q}}\eta_{-3}\eta_{-1} \\
& - q^5\sqrt{(5)_q(4)_q}\eta_{-4}\eta_0,
\end{aligned}$$

$$\begin{aligned}
f\cdot\Psi_3(\eta,\eta)_{-3} = {} & q^{-6}\sqrt{(6)_q(3)_q}\eta_2\eta_{-4} + \frac{q^{-9}\sqrt{(8)_q}-q^{-3}(5)_q(4)_q}{\sqrt{(8)_q}}\eta_1\eta_{-3} \\
& + ((5)_q(4)_q - q^{-6}(7)_q(2)_q)\frac{(5)_q(4)_q}{(8)_q(7)_q(2)_q}\eta_0\eta_{-2} \\
& + (q^{-3}-q^3)(5)_q(4)_q\frac{(6)_q(3)_q}{(8)_q(7)_q(2)_q}\eta_{-1}\eta_{-1} \\
& + (q^6(7)_q(2)_q - (5)_q(4)_q)\frac{(5)_q(4)_q}{(8)_q(7)_q(2)_q}\eta_{-2}\eta_0 \\
& + \frac{q^3(5)_q(4)_q - q^9\sqrt{(8)_q}}{\sqrt{(8)_q}}\eta_{-3}\eta_1 - q^6\sqrt{(6)_q(3)_q}\eta_{-4}\eta_2,
\end{aligned}$$

$$\begin{aligned}
f\cdot\Psi_1(\eta,\eta)_{-1} = {} & q^{-3}\sqrt{(8)_q}\eta_4\eta_{-4} + \frac{q^{-4}(8)_q - q^{-2}(7)_q(2)_q}{\sqrt{(8)_q}}\eta_3\eta_{-3} \\
& + \frac{q^{-1}(6)_q(3)_q - q^{-3}(7)_q(2)_q}{\sqrt{(8)_q}}\eta_2\eta_{-2} + \frac{q^{-2}(6)_q(3)_q-(5)_q(4)_q}{\sqrt{(8)_q}}\eta_1\eta_{-1} \\
& + \frac{(q-q^{-1})(5)_q(4)_q}{\sqrt{(8)_q}}\eta_0\eta_0 + \frac{(5)_q(4)_q - q^2(6)_q(3)_q}{\sqrt{(8)_q}}\eta_{-1}\eta_1 \\
& + \frac{q^3(7)_q(2)_q - q(6)_q(3)_q}{\sqrt{(8)_q}}\eta_{-2}\eta_2 + \frac{q^2(7)_q(2)_q - q^4(8)_q}{\sqrt{(8)_q}}\eta_{-3}\eta_3 \\
& - q^3\sqrt{(8)_q}\eta_{-4}\eta_4\,.
\end{aligned}$$

Applying $\pi_{\mathbb{T}}$ to these vectors, we obtain the following equations:

$$c_{-3} = 0\,, \tag{6.1}$$

$$(q^{-5}-q^{5})\sqrt{(5)_q(4)_q}\,c_0c_{-4} + (q^{5}-q^{-5})\sqrt{\frac{(7)_q(6)_q(3)_q(2)_q}{(8)_q}}\,c_{-2}^2 = 0\,, \tag{6.2}$$

$$\begin{aligned}(q^{-6}-q^{6})\sqrt{(6)_q(3)_q}\,c_2c_{-4} &+ (q^{6}-q^{-6})\frac{(5)_q(4)_q}{(8)_q}c_0c_{-2}\\ &+ (q^{-3}-q^{3})(5)_q(4)_q\frac{(6)_q(3)_q}{(8)_q(7)_q(2)_q}c_{-1}^2 = 0\,,\end{aligned} \tag{6.3}$$

$$\begin{aligned}(q^{-3}-q^{3})\sqrt{(8)_q}\,c_4c_{-4} &+ \frac{(q^{-1}-q)(6)_q(3)_q-(q^{3}-q^{-3})(7)_q(2)_q}{\sqrt{(8)_q}}c_2c_{-2}\\ &+ \frac{(q^{-2}-q^{2})(6)_q(3)_q}{\sqrt{(8)_q}}c_1c_{-1} + \frac{(q-q^{-1})(5)_q(4)_q}{\sqrt{(8)_q}}c_0^2 = 0\,.\end{aligned} \tag{6.4}$$

Also noticing $c_{-3}=0$, we get the following equations via original eigenvectors $\Psi_\nu(\eta,\eta)$ for $\nu = 5,3,2$:

$$(q^{-9}-q^{9})c_{-1}c_{-4} = 0\,, \tag{6.5}$$

$$(q^{-10}-q^{10})c_1c_{-4} + (q^{-2}-q^{2})\frac{(5)_q(4)_q}{(8)_q(7)_q(2)_q}c_{-1}c_{-2} = 0\,, \tag{6.6}$$

$$(q^{-9}+q^{9})c_2c_{-4} + (q^{3}+q^{-3})\sqrt{\frac{(6)_q(5)_q(4)_q(3)_q}{(8)_q(7)_q(2)_q}}\,c_0c_{-2} - \frac{(5)_q(4)_q}{(8)_q(7)_q(2)_q}c_{-1}^2 = 0. \tag{6.7}$$

From (6.5) c_{-1} or c_{-4} are 0. If c_{-4} is 0, then we have $c_{-2}=c_{-1}=c_0=0$ by (6.2), (6.3) and (6.4). This is contradiction. Hence c_{-4} is not 0 and c_{-1} is equal to 0. Next we can derive $c_2c_{-4}=c_0c_{-2}=0$ by (6.3) and (6.7). Then we have $c_{-2}=0$. This yields $c_0=0$ through (6.2). Finally we get $c_{-4}=0$ by (6.4), however, this is a contradiction.

(III) A_5^* case. A C^*-algebra A has a spectral pattern $\pi_0\oplus\pi_6\oplus\pi_{10}\oplus\pi_{12}\oplus\cdots$. Let $\eta = (\eta_s)_{s\in I_6} = \sum_{s\in I_6} c_s\boldsymbol{w}_s^6$ be a non-zero self-conjugate π_6-eigenvector. Equality $c_{-s} = (-q)^s\overline{c_s}$ follows from the self-conjugacy. We show coefficients c_s are all zero and derive contradiction. In order to do it we make use of spectral gaps at π_ν for $\nu = 11,9,8,7,4,2$ and 1.

First we see the gap at 11. We construct a π_{11}-eigenvector $\Psi_{11}(\eta,\eta)$ and apply the lowering operator f to $\Psi_{11}(\eta,\eta)_{-11}$. Then we have the following equalities

$$\begin{aligned}\Psi_{11}(\eta,\eta)_{-11} =& q^{-6}\eta_{-5}\eta_{-6} - q^{6}\eta_{-6}\eta_{-5}\,,\\ f\cdot\Psi_{11}(\eta,\eta)_{-11} =& \sqrt{(11)_q(2)_q}\,\eta_{-4}\eta_{-6} + (q^{-11}-q^{11})\sqrt{(12)_q}\,\eta_{-5}^2 - \sqrt{(11)_q(2)_q}\,\eta_{-6}\eta_{-4}.\end{aligned}$$

Applying $\pi_{\mathbb{T}}$ to the second one, we get $c_{-5}=0$.

Next we see the gap at 9. We make $\Psi_9(\eta,\eta)_{-9}$ and $f\cdot\Psi_9(\eta,\eta)_{-9}$ as follows.

$$\begin{aligned}
\sqrt{(12)_q}\,\Psi_9(\eta,\eta)_{-9} ={}& q^{-15}\sqrt{(12)_q}\eta_{-3}\eta_{-6}-q^{-5}\sqrt{(10)_q(3)_q}\eta_{-4}\eta_{-5}\\
&+q^{5}\sqrt{(10)_q(3)_q}\eta_{-5}\eta_{-4}-q^{15}\sqrt{(12)_q}\eta_{-6}\eta_{-3}\\
\sqrt{(12)_q}\,f\cdot\Psi_9(\eta,\eta)_{-9} ={}& q^{-9}\sqrt{(12)_q(9)_q(4)_q}\eta_{-2}\eta_{-6}+\big(q^{-18}(12)_q-(10)_q(3)_q\big)\eta_{-3}\eta_{-5}\\
&+(q^{9}-q^{-9})\sqrt{(11)_q(10)_q(3)_q(2)_q}\eta_{-4}\eta_{-4}\\
&+\big((10)_q(3)_q-q^{18}(12)_q\big)\eta_{-5}\eta_{-3}-q^{9}\sqrt{(12)_q(9)_q(4)_q}\eta_{-6}\eta_{-2}.
\end{aligned}$$

Applying $\pi_{\mathbb{T}}$ to the first one, we get $c_{-3}c_{-6}=0$. Similarly from the second one we get the following equality

$$\sqrt{(12)_q(9)_q(4)_q}c_{-2}c_{-6}-\sqrt{(11)_q(10)_q(3)_q(2)_q}c_{-4}^2=0\,. \tag{6.8}$$

Next we look at the gap at 8. In this case we need two operators

$$\begin{aligned}
&\sqrt{(12)_q(11)_q(2)_q}\,\Psi_8(\eta,\eta)_{-8}\\
={}&q^{-18}\sqrt{(12)_q(11)_q(2)_q}\eta_{-2}\eta_{-6}-q^{-9}\sqrt{(11)_q(9)_q(4)_q(2)_q}\eta_{-3}\eta_{-5}\\
&+\sqrt{(10)_q(9)_q(4)_q(3)_q}\eta_{-4}\eta_{-4}-q^{9}\sqrt{(11)_q(9)_q(4)_q(2)_q}\eta_{-5}\eta_{-3}\\
&+q^{18}\sqrt{(12)_q(11)_q(2)_q}\eta_{-6}\eta_{-2}\,,\\
&\sqrt{(12)_q(11)_q(2)_q}\,f^2\cdot\Psi_8(\eta,\eta)_{-8}\\
={}&q^{-6}\sqrt{(12)_q(11)_q(8)_q(7)_q(6)_q(5)_q(2)_q}\eta_0\eta_{-6}\\
&+\big((q^{-13}+q^{-15})(12)_q-q(9)_q(4)_q\big)\sqrt{(11)_q(8)_q(5)_q(2)_q}\eta_{-1}\eta_{-5}\\
&+\big\{q^{8}(10)_q(9)_q(4)_q(3)_q+q^{-22}(12)_q(11)_q(2)_q-(q^{-6}+q^{-8})(11)_q(8)_q(4)_q(2)_q\big\}\eta_{-2}\eta_{-4}\\
&+\big((10)_q(3)_q-(q^{-15}+q^{15})(11)_q\big)(2)_q\sqrt{(10)_q(9)_q(4)_q(3)_q}\eta_{-3}\eta_{-3}\\
&+\big\{q^{-8}(10)_q(9)_q(4)_q(3)_q+q^{22}(12)_q(11)_q(2)_q-(q^{6}+q^{8})(11)_q(8)_q(4)_q(2)_q\big\}\eta_{-4}\eta_{-2}\\
&+\big((q^{13}+q^{15})(12)_q-q^{-1}(9)_q(4)_q\big)\sqrt{(11)_q(8)_q(5)_q(2)_q}\eta_{-5}\eta_{-1}\\
&+q^{6}\sqrt{(12)_q(11)_q(8)_q(7)_q(6)_q(5)_q(2)_q}\eta_{-6}\eta_0.
\end{aligned}$$

From the first equality we get the following one

$$(q^{-18}+q^{18})\sqrt{(12)_q(11)_q(2)_q}c_{-2}c_{-6}+\sqrt{(10)_q(9)_q(4)_q(3)_q}c_{-4}^2=0\,. \tag{6.9}$$

Equalities (6.8) and (6.9) shows $c_{-2}c_{-6} = 0$ and $c_{-4} = 0$. Hence by the above second equality we get

$$\begin{aligned}(q^{-6}+q^{6})&\sqrt{(12)_q(11)_q(8)_q(7)_q(6)_q(5)_q(2)_q}c_0c_{-6}\\&+\big((10)_q(3)_q-(q^{-15}+q^{15})(11)_q\big)(2)_q\sqrt{(10)_q(9)_q(4)_q(3)_q}c_{-3}^2=0.\end{aligned}\tag{6.10}$$

Next we see the gap at 7. The operators $\sqrt{(12)_q(11)_q(2)_q}\,\Psi_7(\eta,\eta)_{-7}$ and $\sqrt{(12)_q(11)_q(2)_q}\,f\cdot\Psi_7(\eta,\eta)_{-7}$ are as follows.

$$\begin{aligned}&\sqrt{(12)_q(11)_q(2)_q}\,\Psi_7(\eta,\eta)_{-7}\\&\qquad=q^{-20}\sqrt{(12)_q(11)_q(2)_q}\eta_{-1}\eta_{-6}-q^{-12}\sqrt{(11)_q(8)_q(5)_q(2)_q}\eta_{-2}\eta_{-5}\\&\qquad\quad+q^{-4}\sqrt{(9)_q(8)_q(5)_q(4)_q}\eta_{-3}\eta_{-4}-q^{4}\sqrt{(9)_q(8)_q(5)_q(4)_q}\eta_{-4}\eta_{-3}\\&\qquad\quad+q^{12}\sqrt{(11)_q(8)_q(5)_q(2)_q}\eta_{-5}\eta_{-2}-q^{20}\sqrt{(12)_q(11)_q(2)_q}\eta_{-6}\eta_{-1},\end{aligned}$$

$$\begin{aligned}\sqrt{(12)_q(11)_q(2)_q}\,f\cdot\Psi_7(\eta,\eta)_{-7}=&q^{-14}\sqrt{(12)_q(11)_q(7)_q(6)_q(2)_q}\eta_0\eta_{-6}\\&+\big(q^{-21}(12)_q-q^{-7}(8)_q(5)_q\big)\sqrt{(11)_q(2)_q}\eta_{-1}\eta_{-5}\\&+\big((9)_q(4)_q-q^{-14}(11)_q(2)_q\big)\sqrt{(8)_q(5)_q}\eta_{-2}\eta_{-4}\\&+(q^{-7}-q^{7})\sqrt{(10)_q(9)_q(8)_q(5)_q(4)_q(3)_q}\eta_{-3}\eta_{-3}\\&+\big((9)_q(4)_q-q^{-14}(11)_q(2)_q\big)\sqrt{(8)_q(5)_q}\eta_{-4}\eta_{-2}\\&+\big(q^{7}(8)_q(5)_q-q^{21}(12)_q\big)\sqrt{(11)_q(2)_q}\eta_{-5}\eta_{-1}\\&-q^{14}\sqrt{(12)_q(11)_q(7)_q(6)_q(2)_q}\eta_{-6}\eta_0.\end{aligned}$$

From the second one we get the following equation

$$\begin{aligned}(q^{-14}-q^{14})&\sqrt{(12)_q(11)_q(7)_q(6)_q(2)_q}c_0c_{-6}\\&+(q^{-7}-q^{7})\sqrt{(10)_q(9)_q(8)_q(5)_q(4)_q(3)_q}c_{-3}^2=0\,.\end{aligned}\tag{6.11}$$

On equations (6.10) and (6.11) we can easily show that the determinant of the following matrix is not 0 for $0<q<1$

$$\begin{pmatrix}(q^{-6}+q^{6}) & \big((10)_q(3)_q-(q^{-15}+q^{15})(11)_q\big)(2)_q\\(q^{-14}-q^{14}) & (q^{-7}-q^{7})(8)_q(5)_q\end{pmatrix}.$$

Hence we obtain $c_0c_{-6}=0$ and $c_{-3}=0$.

Next we see the gap at 4. We use an eigenvector $\Psi_4(\eta,\eta)$

$$
\begin{aligned}
&\sqrt{(12)_q(11)_q(10)_q(9)_q(4)_q(3)_q(2)_q}\,\Psi_4(\eta,\eta)_{-4}\\
&\qquad = q^{-20}\sqrt{(12)_q(11)_q(10)_q(9)_q(4)_q(3)_q(2)_q}\eta_2\eta_{-6}\\
&\qquad\quad - q^{-15}\sqrt{(11)_q(10)_q(9)_q(8)_q(5)_q(4)_q(3)_q(2)_q}\eta_1\eta_{-5}\\
&\qquad\quad + q^{-10}\sqrt{(10)_q(9)_q(8)_q(7)_q(6)_q(5)_q(4)_q(3)_q}\eta_0\eta_{-4}\\
&\qquad\quad - q^{-5}(7)_q(6)_q\sqrt{(9)_q(8)_q(5)_q(4)_q}\eta_{-1}\eta_{-3}\\
&\qquad\quad + (8)_q(7)_q(6)_q(5)_q\eta_{-2}^2\\
&\qquad\quad - q^{5}(7)_q(6)_q\sqrt{(9)_q(8)_q(5)_q(4)_q}\eta_{-3}\eta_{-1}\\
&\qquad\quad + q^{10}\sqrt{(10)_q(9)_q(8)_q(7)_q(6)_q(5)_q(4)_q(3)_q}\eta_{-4}\eta_0\\
&\qquad\quad - q^{15}\sqrt{(11)_q(10)_q(9)_q(8)_q(5)_q(4)_q(3)_q(2)_q}\eta_{-5}\eta_1\\
&\qquad\quad + q^{-20}\sqrt{(12)_q(11)_q(10)_q(9)_q(4)_q(3)_q(2)_q}\eta_{-6}\eta_2\,.
\end{aligned}
$$

We know $c_2c_{-6}=0$ because of $c_{-2}c_{-6}=0$. Hence the above equality derives $c_{-2}=0$.

Next we see the gap at 2. The operator $\Psi_2(\eta,\eta)_{-2}$ is

$$
\begin{aligned}
&\sqrt{(12)_q(11)_q(2)_q}\,\Psi_2(\eta,\eta)_{-2}\\
&\qquad = q^{-15}\sqrt{(12)_q(11)_q(2)_q}\eta_4\eta_{-6} - q^{-12}\sqrt{(11)_q(10)_q(3)_q(2)_q}\eta_3\eta_{-5}\\
&\qquad\quad + q^{-9}\sqrt{(10)_q(9)_q(4)_q(3)_q}\eta_2\eta_{-4} - q^{-6}\sqrt{(9)_q(8)_q(5)_q(4)_q}\eta_1\eta_{-3}\\
&\qquad\quad + q^{-3}\sqrt{(8)_q(7)_q(6)_q(5)_q}\eta_0\eta_{-2} - (7)_q(6)_q\eta_{-1}^2\\
&\qquad\quad + q^{3}\sqrt{(8)_q(7)_q(6)_q(5)_q}\eta_{-2}\eta_0 - q^{6}\sqrt{(9)_q(8)_q(5)_q(4)_q}\eta_{-3}\eta_1\\
&\qquad\quad + q^{9}\sqrt{(10)_q(9)_q(4)_q(3)_q}\eta_{-4}\eta_2 - q^{12}\sqrt{(11)_q(10)_q(3)_q(2)_q}\eta_{-5}\eta_3\\
&\qquad\quad + q^{15}\sqrt{(12)_q(11)_q(2)_q}\eta_{-6}\eta_4\,.
\end{aligned}
$$

This shows $c_{-1}=0$.

Finally we see the gap at 1. The operators $\Psi_1(\eta,\eta)_{-1}$ and $f\cdot\Psi_1(\eta,\eta)_{-1}$ are

$$
\begin{aligned}
\sqrt{(12)_q}\,\Psi_1(\eta,\eta)_{-1} &= q^{-11}\sqrt{(12)_q}\eta_5\eta_{-6} - q^{-9}\sqrt{(11)_q(2)_q}\eta_4\eta_{-5} + q^{-7}\sqrt{(10)_q(3)_q}\eta_3\eta_{-4}\\
&\quad - q^{-5}\sqrt{(9)_q(4)_q}\eta_2\eta_{-3} + q^{-3}\sqrt{(8)_q(5)_q}\eta_1\eta_{-2} - q^{-1}\sqrt{(7)_q(6)_q}\eta_0\eta_{-1}
\end{aligned}
$$

$$
\begin{aligned}
&+ q\sqrt{(6)_q(7)_q}\eta_{-1}\eta_0 - q^3\sqrt{(5)_q(8)_q}\eta_{-2}\eta_1 + q^5\sqrt{(4)_q(9)_q}\eta_{-3}\eta_2 \\
&+ q^7\sqrt{(10)_q(3)_q}\eta_{-4}\eta_3 - q^9\sqrt{(11)_q(2)_q}\eta_{-5}\eta_4 - q^{11}\sqrt{(12)_q}\eta_{-6}\eta_5\,,
\end{aligned}
$$

$$
\begin{aligned}
&\sqrt{(12)_q}\, f \cdot \Psi_1(\eta,\eta)_{-1} \\
&\quad = q^{-5}(12)_q\eta_6\eta_{-6} + \big(q^{-6}(12)_q - q^{-4}(11)_q(2)_q\big)\eta_5\eta_{-5} \\
&\qquad + \big(q^{-3}(10)_q(3)_q - q^{-5}(11)_q(2)_q\big)\eta_4\eta_{-4} + \big(q^{-4}(10)_q(3)_q - q^{-2}(9)_q(4)_q\big)\eta_3\eta_{-3} \\
&\qquad + \big(q^{-1}(8)_q(5)_q - q^{-3}(9)_q(4)_q\big)\eta_2\eta_{-2} + \big(q^{-2}(8)_q(5)_q - (7)_q(6)_q\big)\eta_1\eta_{-1} \\
&\qquad + (q-q^{-1})(7)_q(6)_q\eta_0\eta_0 + \big((7)_q(6)_q - q^2(8)_q(5)_q\big)\eta_{-1}\eta_1 \\
&\qquad + \big(q^3(9)_q(4)_q - q(8)_q(5)_q\big)\eta_{-2}\eta_2 + \big(q^2(9)_q(4)_q - q^4(10)_q(3)_q\big)\eta_{-3}\eta_3 \\
&\qquad + \big(q^5(11)_q(2)_q - q^3(10)_q(3)_q\big)\eta_{-4}\eta_4 + \big(q^4(11)_q(2)_q - q^6(12)_q\big)\eta_{-5}\eta_5 \\
&\qquad - q^5(12)_q\eta_{-6}\eta_6\,.
\end{aligned}
$$

Since we have $c_s = 0$ for $s = 5, 4, 3, 2$ and 1, the above equality shows

$$(q^{-5} - q^5)(12)_q c_6 c_{-6} + (q - q^{-1})(7)_q(6)_q c_0^2 = 0\,.$$

We also know $c_0 c_6 = 0$ because of $c_0 c_{-6} = 0$. Hence we obtain $c_{-6} = c_0 = 0$.

(IV)D^*_∞ case. We conclude that this case actually occurs and a right coideal of this type is unique up to conjugation by β^L_z.

Lemma 6.3. *Consider the quantum sphere $C(S^2_{q,0})$. Then a vector*

$$\Big(q\sqrt{(3)_q!}, 0, -\sqrt{(4)_q}, 0, q^{-1}\sqrt{(3)_q!}\Big)\, w(\pi_2)$$

is a π_2-eigenvector of $C(S^2_{q,0})$.

Proof. Recall a highest weight vector of weight 1 $(\xi^1_0)_{-1}$ in $C(S^2_{q,0})$, $(\xi^1_0)_{-1} = q^{\frac{1}{2}}\sqrt{(2)_q}^{-1}x^2 - q^{-\frac{1}{2}}\sqrt{(2)_q}^{-1}v^2$. Then we can easily compute a highest weight vector of weight 2 in $C(S^2_{q,0})$ as follows, $(\xi^1_0)^2_{-1} = q(2)_q^{-1}w(\pi_2)_{-2-2} - (q^{-2} + q^2)\sqrt{(4)_q!}^{-1}w(\pi_2)_{0,-2} + q^{-1}(2)_q^{-1}w(\pi_2)_{2,-2}$. Multiply $(2)_q\sqrt{(3)_q!}$ to both sides we get the desired highest weight vector. □

Let $\xi = (\xi_{-1}, \xi_0, \xi_1)$ be the canonical π_1-eigenvector of $C(S^2_{q,0})$. By its definition, $\{\xi_r\}_{r\in I_1}$ satisfies the following conditions

$$\xi^*_{-1} = -q\,\xi_1,\ \xi^*_0 = \xi_0,\ \xi_{\pm1}\xi_0 = q^{\pm2}\xi_0\xi_{\pm1}, \tag{6.12}$$

$$\xi_1\xi_{-1} = (q+q^{-1})^{-1}(q^2\xi_0^2 - 1), \tag{6.13}$$

$$\xi_{-1}\xi_1 = (q+q^{-1})^{-1}(q^{-2}\xi_0^2 - 1). \tag{6.14}$$

An embedding of $C(S^2_{q,0})$ into $C(SU_q(2))$ is given by

$$\xi = (\sqrt{1+q^{-2}}^{-1}, 0, -\sqrt{1+q^2}^{-1})\, w(\pi_1)\,,$$

or more precisely

$$\xi_{-1} = \sqrt{1+q^{-2}}^{-1} x^2 - \sqrt{1+q^2}^{-1} v^2 \,, \quad \xi_0 = qxu - vy \,,$$
$$\xi_{-1} = \sqrt{1+q^{-2}}^{-1} u^2 - \sqrt{1+q^2}^{-1} y^2 \,.$$

Consider the smooth part of $C(S^2_{q,0})$, that is, the $*$-algebra generated by $\{\xi_r\}_{r\in I_1}$. Denote it by $\mathcal{B}$. Let $\mathcal{B}_{\mathrm{e}}$ be a $SU_q(2)$-invariant $*$-subalgebra of $\mathcal{B}$ generated by $\{\xi_r\xi_s\}_{r,s\in I_1}$. In [DK, Proposition 2.9.], it is shown that $\{\xi_0^m \xi_{-1}^n\}_{m,n\in\mathbb{Z}_{\geq 0}}$ and $\{\xi_0^m \xi_1^n\}_{m,n\in\mathbb{Z}_{\geq 0}}$ are basis for a vector space $\mathcal{B}$. Since the $*$-subalgebra $\mathcal{B}_{\mathrm{e}}$ is generated by words of even length (this is because of the previous equalities), we see ξ_r is not in $\mathcal{B}_{\mathrm{e}}$ for $r \in I_1$. In particular, we have $\mathcal{B}_{\mathrm{e}} \subsetneq \mathcal{B}$. Let B_{e} be the norm closure of $\mathcal{B}_{\mathrm{e}}$. By $SU_q(2)$-invariance of $\mathcal{B}_{\mathrm{e}}$ we see that B_{e} is also $SU_q(2)$-invariant and $B_{\mathrm{e}} \subsetneq C(S^2_{q,0})$. Next we prove that π_3-spectral subspace of B_{e} is 0. Since $\xi_{-1}^3 \neq 0$ is the highest weight vector in $C(S^2_{q,0})_{\pi_3}$, it is not contained in $\mathcal{B}_{\mathrm{e}}$. Again $SU_q(2)$-invariance yields that ξ_{-1}^3 is not contained in B_{e}. In a similar way we can prove that all the odd spin spectral subspaces of B_{e} are 0. Of course, even spin spectral subspaces of B_{e} are not 0 because of $\xi_{-1}^{2n} \neq 0$ for all $n \in \mathbb{Z}_{\geq 0}$. The spectral pattern $\oplus_{k\in\mathbb{Z}_{\geq 0}} \pi_{2k}$ follows from the spectral pattern of $C(S^2_{q,0}) \oplus_{k\in\mathbb{Z}_{\geq 0}} \pi_k$. Therefore, we have shown that B_{e} is of type D^*_∞. Next lemma says that a right coideal of D^*_∞-type is unique up to conjugation of the automorphism β^L_z.

Lemma 6.4. *Let $A \subset C(SU_q(2))$ be a right coideal of type D^*_∞. Then there exists $z \in \mathbb{C}$ such that A is $\beta^L_z(B_e)$.*

Proof. By assumption A has the spectral pattern $\pi_0 \oplus \pi_2 \oplus \pi_4 \oplus \cdots$. In the following discussion we do not need to use the gap at π_3. Let $\eta = (\eta_s)_{s\in I_2} = \sum_{s\in I_2} c_s \boldsymbol{w}^2_s$ be a self conjugate π_2-eigenvector of A. We consider zero vectors $\Psi_1(\eta,\eta)_{-1}$ and $f \cdot \Psi_1(\eta,\eta)_{-1}$. They become as follows.

$$\Psi_1(\eta,\eta)_{-1} = q^{-3}\,\eta_1\eta_{-2} - q^{-1}\sqrt{\frac{(3)_q(2)_q}{(4)_q}}\,\eta_0\eta_{-1} + q\sqrt{\frac{(3)_q(2)_q}{(4)_q}}\,\eta_{-1}\eta_0 - q^3\,\eta_{-2}\eta_1 \,,$$

$$\begin{aligned} f \cdot \Psi_1(\eta,\eta)_{-1} = {} & q^{-1}\sqrt{(4)_q}\,\eta_2\eta_{-2} + \sqrt{(4)_q}^{-1}(q^{-2}(4)_q - (3)_q!)\,\eta_1\eta_{-1} \\ & + \sqrt{(4)_q}^{-1}(q-q^{-1})(3)_q!\,\eta_0^2 + \sqrt{(4)_q}^{-1}((3)_q! - q^2(4)_q)\,\eta_{-1}\eta_1 \\ & - q\sqrt{(4)_q}\,\eta_{-2}\eta_2 \,. \end{aligned}$$

Apply the map $\pi_{\mathbb{T}}$ to the above equalities and we obtain

$$(q^{-3} - q^3)c_1c_{-2} + (q - q^{-1})\sqrt{\frac{(3)_q!}{(4)_q}}c_{-1}c_0 = 0 \,,$$

$$(q^{-1} - q)\sqrt{(4)_q}c_2c_{-2} + (q^{-2} - q^2)\sqrt{(4)_q}c_1c_{-1} \,,$$

$$+ (q - q^{-1})\sqrt{(4)_q}^{-1}(3)_q!c_0^2 = 0 \,.$$

For simplicity, we may assume that c_1 is a real number by applying a conjugation β_z^L. Then we have $c_{-1} = -qc_1$ by the self-conjugacy of η. Assume c_1 is not 0. Then from the first equality we get $c_{-2} = -q\sqrt{\frac{(2)_q}{(4)_q(3)_q}}c_0$. If we put this one to the second equality, we get

$$\frac{(2)_q - (3)_q^2(2)_q}{(3)_q}c_0^2 + (2)_q(4)_q c_1 c_{-1} = 0\,.$$

Since we know $(3)_q > 1$ and $c_1c_{-1} = -qc_1^2 < 0$, the left hand side is strictly negative unless $c_0 = c_1 = 0$. This also shows $c_2 = 0$ and this is a contradiction. Hence we get $c_1 = 0$. Then from the second equality we get $|c_2| = q^{-1}\sqrt{\frac{(3)_q!}{(4)_q}}|c_0|$. We may assume $c_0 = -\sqrt{(4)_q}$ by scalar multiplication and $c_2 = q^{-1}\sqrt{(3)_q!}$ by a conjugation of β_z^L.

□

From now on, the right coideal of type D_∞^* generated by $q\sqrt{(3)_q!}w(\pi_2)_{-2,s} - \sqrt{(4)_q}w(\pi_2)_{0,s} + q^{-1}\sqrt{(3)_q!}w(\pi_2)_{2,s}$ for $s \in I_2$ is denoted by $A_{D_\infty^*}$. We remark that $\boldsymbol{w}_0^2$ does not generates a right coideal of type D_∞^*. In fact, it generates the canonical homogeneous sphere $C(\mathbb{T}\setminus SU_q(2))$. This is essentially due to the effect of $q \neq 1$ (see also Lemma 7.5). The C^*-algebra $A_{D_\infty^*}$ is actually the Toeplitz algebra as we see below. By [MNW, Lemma 3.2] there exists the matrix units $\{e_{i,j}^+\}_{i,j\in\mathbb{Z}_{\geq 0}}$ and $\{e_{i,j}^-\}_{i,j\in\mathbb{Z}_{\geq 0}}$ in $C(S_{q,0}^2)$ which satisfy the following equalities

$$\xi_{-1}e_{k,k}^{\pm} = \pm\gamma_k e_{k+1,k}^{\pm},\ \xi_1 e_{k,k}^{\pm} = \mp q^{-1}\gamma_{k-1}e_{k-1,k}^{\pm}, \tag{6.15}$$

$$\xi_0 = \sum_{k=0}^{\infty} -q^{2k+1}e_{k,k}^{-} + \sum_{k=0}^{\infty} q^{2k+1}e_{k,k}^{+} \tag{6.16}$$

for all $k \in \mathbb{Z}_{\geq 0}$, where $\gamma_k = q^{\frac{1}{2}}(q+q^{-1})^{-\frac{1}{2}}(1-q^{4k+4})^{\frac{1}{2}}$. Note the equality $1 = \sum_{k=0}^{\infty} e_{k,k}^{-} + \sum_{k=0}^{\infty} e_{k,k}^{+}$ in $C(SU_q(2))$ where the summation converges in the strong operator topology in $\mathbb{B}(L^2(SU_q(2)))$. In order to verify it, it suffices to show $1 = \sum_{k=0}^{\infty} h(e_{k,k}^{-}) + \sum_{k=0}^{\infty} h(e_{k,k}^{+})$, because the Haar state h is normal and faithful. This equality holds by the following lemma.

Lemma 6.5. *We have* $h(e_{k,k}^{\pm}) = 2^{-1}q^{2k}(1-q^2)$ *for all* $k \in \mathbb{Z}_{\geq 0}$.

Proof. Let $X_{\pi_1} = \mathbb{C}\xi$ be the π_1-eigenvector space of $C(S_{q,0}^2)$. Recall the conjugation map T and the one-parameter unitary group U_t on X_{π_1} which are defined by $(T\xi)_r = (-q)^{-r}\xi_{-r}^*$ and $(U_t\xi)_r = (F_{\pi_1})_{r,r}^{-it}\sigma_t^h(\xi_r)$ for all $r \in I_{\pi_1}$ and $t \in \mathbb{R}$, where σ_t^h is the modular automorphism group with respect to the invariant state h on $C(S_{q,0}^2)$. As is proved in Lemma 3.7, we have $IT^*TI^* = U_i$ where the unitary map I is defined by $(I\xi)_r = \xi_{-r}$ for $r \in I_{\pi_1}$. It shows that the spectrum of U_i is inverse-closed and hence $U_i = \mathrm{id}$. It yields the formula $\sigma_t^h(\xi_r) = q^{-2rit}\xi_r$. We have the equality $h(\xi_1 e^{\pm}_{k,k}\xi_{-1}) = -q^{-1}\gamma^2_{k-1}h(e^{\pm}_{k-1,k-1})$ by (6.15). The left hand side is equal to

$$\begin{aligned} h(\xi_1 e^{\pm}_{k,k}\xi_{-1}) &= h(\sigma^h_i(\xi_{-1})\xi_1 e^{\pm}_{k,k}) \\ &= q^{-2}h(\xi_{-1}\xi_1 e^{\pm}_{k,k}) \\ &= q^{-2}(q+q^{-1})^{-1}(q^{4k}-1)h(e^{\pm}_{k,k}), \end{aligned}$$

where we used (6.13) for the third equality. Hence we have $h(e^{\pm}_{k,k}) = q^{2k}h(e^{\pm}_{0,0})$ for $k \geq 1$. Since the equality $h(\xi_0) = 0$ holds, we get $h(e^{+}_{0,0}) = h(e^{-}_{0,0})$. Using Lemma 3.6, we see $h(\xi_0^2) = (3)^{-1}$ and obtain $h(e^{\pm}_{0,0}) = 2^{-1}(1-q^2)$. ☐

Now we consider the invariant subalgebra $A_{D^*_\infty} \subset C(S^2_{q,0})$. Let us define the matrix units $e_{i,j} = e^{-}_{i,j} + e^{+}_{i,j}$ for $i,j \in \mathbb{Z}_{\geq 0}$. The C^*-algebra $A_{D^*_\infty}$ is generated by them and the unilateral shift. It also concludes that the right coideal von Neumann algebra $A_{D^*_\infty}{}''$ is isomorphic to $\mathbb{B}(\ell^2(\mathbb{Z}_{\geq 0}))$. Especially the quantum group $SU_q(2)$ can act ergodically on every separable type I factor.

(V) D^*_m ($m \geq 2$) case. We will derive the non-existence of this case. We study a C^*-algebra A whose spectral pattern is $\oplus_{k\in\mathbb{Z}_{\geq 0}}\bigl(\frac{1+(-1)^k}{2} + \bigl[\frac{k}{m}\bigr]\bigr)\pi_k$. Making use of a gap at π_1 as Lemma 6.4, we can similarly prove that elements of one π_2-eigenvector generates a right coideal of type D^*_∞. (the π_2-eigenvector space is two-dimensional when m is equal to 2.) So we may assume there is a π_2-eigenvector ξ_0^2 in A. Recall that ξ^2_0 is defined by $\xi^2_0 = \Bigl(q\sqrt{(3)_q(2)_q}, 0, -\sqrt{(4)_q}, 0, q^{-1}\sqrt{(3)_q(2)_q}\Bigr)\, w(\pi_2)$. Note that the index 0 of ξ^2_0 means $\lambda = 0$ of $C(S^2_{q,\lambda})$.

Lemma 6.6. *Let n be a half integer with $n \geq 2$. Then we have the following equalities for all $t \in I_n$*

$$\Psi_{n-2}(\boldsymbol{w}^2_{-2}, \boldsymbol{w}^n_t) = q^{-2t+6}\sqrt{\frac{(n+t)_q(n+t-1)_q(n+t-2)_q(n+t-3)_q}{(2n)_q(2n-1)_q(2n-2)_q(2n-3)_q}}\,\boldsymbol{w}^{n-2}_{t-2}\,,$$

$$\Psi_{n-2}(\boldsymbol{w}^2_{0}, \boldsymbol{w}^n_t) = q^{-2n-2t+4}\sqrt{\frac{(4)_q(3)_q}{(2)_q}}\sqrt{\frac{(n+t)_q(n+t-1)_q(n-t)_q(n-t-1)_q}{(2n)_q(2n-1)_q(2n-2)_q(2n-3)_q}}\,\boldsymbol{w}^{n-2}_{t},$$

$$\Psi_{n-2}(\boldsymbol{w}^2_{2}, \boldsymbol{w}^n_t) = q^{-4n-2t+2}\sqrt{\frac{(n-t)_q(n-t-1)_q(n-t-2)_q(n-t-3)_q}{(2n)_q(2n-1)_q(2n-2)_q(2n-3)_q}}\,\boldsymbol{w}^{n-2}_{t+2}\,.$$

Proof. For the first equality we use $w(\pi_2)_{-2,2-r} = \bigl[{4 \atop r}\bigr]^{\frac{1}{2}}_{q^2} x^r u^{4-r}$ for $r \in I_2$. From this we obtain

$$\Psi_{n-2}(\boldsymbol{w}^2_{-2}, \boldsymbol{w}^n_t)_{-(n-2)} = \sum_{r=0}^{4}(C^{2,n}_{n-2})_r \begin{bmatrix}4\\ r\end{bmatrix}^{\frac{1}{2}}_{q^2} x^r u^{4-r} w(\pi_n)_{t,-n+r}$$

for all $t \in I_n$ where the coefficients $(C^{2,n}_{n-2})_r$ are given as follows.

$$(C^{2,n}_{n-2})_0 = q^{-2(n-1)}\,,\ (C^{2,n}_{n-2})_1 = -q^{-(n-1)}\sqrt{\tfrac{(4)_q}{(2n)_q}}\,,$$
$$(C^{2,n}_{n-2})_2 = \sqrt{\tfrac{(4)_q(3)_q}{(2n)_q(2n-1)_q}}\,,\ (C^{2,n}_{n-2})_3 = -q^{n-1}\sqrt{\tfrac{(4)_q(3)_q(2)_q}{(2n)_q(2n-1)_q(2n-2)_q}}\,,$$
$$(C^{2,n}_{n-2})_4 = q^{2(n-1)}\sqrt{\tfrac{(4)_q(3)_q(2)_q}{(2n)_q(2n-1)_q(2n-2)_q(2n-3)_q}}\,.$$

This must be a scalar multiple of $x^{n-t}v^{n+t-4}$. So we may compute only the $r=4$ term

$$(C^{2,n}_{n-2})_4\, x^4 w(\pi_n)_{t,-n+4}.$$

If $w(\pi_n)_{t,-n+4}$ contains the word u such that $t \lneq -n+4$, we can disregard the effect of this term. In order to simplify calculations we use the symbol $\sim$ which ignores terms with a power of u. Assume $t \geq -n+4$ then we have

$$w(\pi_n)_{t,-n+4} \sim q^{4(-n-t+4)} \left[\begin{smallmatrix} n+t \\ 4 \end{smallmatrix}\right]^{\frac{1}{2}}_{q^2} \left[\begin{smallmatrix} 2n-4 \\ n-t \end{smallmatrix}\right]^{\frac{1}{2}}_{q^2} x^{n-t-4}v^{n+t-4} \text{ if } t-n+4 \leq 0$$
$$w(\pi_n)_{t,-n+4} \sim q^{(n+t-4)(t-n)} \left[\begin{smallmatrix} n+t \\ 4 \end{smallmatrix}\right]^{\frac{1}{2}}_{q^2} \left[\begin{smallmatrix} 2n-4 \\ n-t \end{smallmatrix}\right]^{\frac{1}{2}}_{q^2} v^{n+t-4}y^{-n+t+4} \text{ if } t-n+4 \geq 0.$$

In both cases we have the same result

$$\begin{aligned} x^4 w(\pi_n)_{t,-n+4} &\sim q^{4(-n-t+4)} \begin{bmatrix} n+t \\ 4 \end{bmatrix}^{\frac{1}{2}}_{q^2} \begin{bmatrix} 2n-4 \\ n-t \end{bmatrix}^{\frac{1}{2}}_{q^2} x^{n-t}v^{n+t-4} \\ &= q^{4(-n-t+4)} \begin{bmatrix} n+t \\ 4 \end{bmatrix}^{\frac{1}{2}}_{q^2} w(\pi_{n-2})_{t-2,-(n-2)} \end{aligned}$$

by simple calculation. Hence we obtain the first equality

$$\begin{aligned} \Psi_{n-2}(\boldsymbol{w}^2_{-2}, \boldsymbol{w}^n_t)_{-(n-2)} &= q^{2(n-1)} q^{4(-n-t+4)} \sqrt{\frac{(4)_q!}{(2n)_q \cdots (2n-3)_q}} \\ &\quad \cdot \begin{bmatrix} n+t \\ 4 \end{bmatrix}^{\frac{1}{2}}_{q^2} w(\pi_{n-2})_{t-2,-(n-2)} \\ &= q^{-2t+6} \sqrt{\frac{(n+t)_q \cdots (n+t-3)_q}{(2n)_q \cdots (2n-3)_q}}\, w(\pi_{n-2})_{t-2,-(n-2)}\,. \end{aligned}$$

Next we show the second equality. Then the matrix element $w(\pi_2)_{0,2-r}$ is as follows.

$$w(\pi_2)_{0,2-r} \sim q^{(4-r)(2-r)} \left[\begin{smallmatrix} 2 \\ r-2 \end{smallmatrix}\right]^{\frac{1}{2}}_{q^2} \left[\begin{smallmatrix} r \\ 2 \end{smallmatrix}\right]^{\frac{1}{2}}_{q^2} x^{r-2}v^{r-2} \qquad \text{if } r \geq 2\,,$$
$$w(\pi)_{0,2-r} \sim 0 \qquad \text{if } r \leq 1.$$

Hence we may only compute the coefficient of $x^{n-t-2}v^{n+t-2}$ of the next one

$$\sum_{r\geq 2}(C^{2,n}_{n-2})_r q^{(4-r)(2-r)} \begin{bmatrix} 2 \\ r-2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} r \\ 2 \end{bmatrix}_{q^2}^{\frac{1}{2}} x^{r-2}v^{r-2}w(\pi_n)_{t,-n+r}.$$

If $t \geq -n+4$, then we have

$$w(\pi_2)_{0,2-r}w(\pi_n)_{t,-n+r} \sim q^{r^2-(2t+4)r-2n+2t+8} \begin{bmatrix} n+t \\ r \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-r \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}}$$
$$\begin{bmatrix} 2 \\ r-2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} r \\ 2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ n+t-2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} w(\pi_{n-2})_{t,-(n-2)}\,.$$

When $t < -n+4$, then $w(\pi_n)_{t,-n+r}$ contains the word u if and only if $2n-r < n-t$. In this case the binomial $\left[\begin{smallmatrix} 2n-r \\ n-t \end{smallmatrix}\right]_{q^2}^{\frac{1}{2}}$ is 0 by definition. From this observation we may continue the computation under $t \geq -n+4$ in order to get the result about general t. Then we can carry out the following calculation

$$\begin{aligned}
&\sum_{r=2}^{4}(C^{2,n}_{n-2})_r q^{r^2-(2t+4)r-2n+2t+8} \begin{bmatrix} n+t \\ r \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-r \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2 \\ r-2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} r \\ 2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ n+t-2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&= (C^{2,n}_{n-2})_2 q^{-2n-2t+4} \begin{bmatrix} n+t \\ 2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-2 \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ n+t-2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&\quad + (C^{2,n}_{n-2})_3 q^{-2n-4t+5} \begin{bmatrix} n+t \\ 3 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-3 \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2 \\ 1 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 3 \\ 2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ n+t-2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&\quad + (C^{2,n}_{n-2})_4 q^{-2n-4t+5} \begin{bmatrix} n+t \\ 4 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2 \\ 1 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 4 \\ 2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ n+t-2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&= \begin{bmatrix} 2n-4 \\ n+t-2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \sqrt{\frac{(4)_q(3)_q}{(2)_q}} \sqrt{\frac{(2n-4)_q!}{(2n)_q\cdots(2n-3)_q}} \sqrt{\frac{(n+t)_q!}{(n-t)_q!}}\, q^{\frac{1}{2}n^2-\frac{1}{2}t^2-n-t+1} \\
&\quad \cdot \left\{ \frac{(2n-2)_q(2n-3)_q}{(n+t-2)_q!} q^{-n+t+1} - \frac{(2n-3)_q(2)_q}{(n+t-3)_q!} + \frac{q^{n-t-1}}{(n+t-4)_q!} \right\} \\
&= q^{-2n-2t+4} \sqrt{\frac{(4)_q(3)_q}{(2)_q}} \sqrt{\frac{(n+t)_q(n+t-1)_q(n-t)_q(n-t-1)_q}{(2n)_q(2n-1)_q(2n-2)_q(2n-3)_q}}\,.
\end{aligned}$$

Hence we have proved the second equality.

Finally we prove the third equality. Then the matrix element $w(\pi_2)_{2,2-r}$ is as follows.

$$w(\pi_2)_{2,2-r} = \begin{bmatrix} 4 \\ r \end{bmatrix}_{q^2}^{\frac{1}{2}} v^r y^{4-r}$$

for all $0 \leq r \leq 4$. If $t \geq -n+4$, we have

$$\begin{aligned}
v^r y^{4-r} w(\pi_n)_{t,-n+r} &\sim q^{r(r-2t-4)} \begin{bmatrix} n+t \\ r \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-r \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} x^{n-t-4} v^{n+t} \\
&= q^{r(r-2t-4)} \begin{bmatrix} n+t \\ r \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-r \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} w(\pi_{n-2})_{t+2,-(n-2)} \, .
\end{aligned}$$

For $t < -n+4$, $w(\pi_n)_{t,-n+r}$ contains the word u if and only if $n+t < r$. Then the binomial $\left[\begin{smallmatrix} n+t \\ r \end{smallmatrix}\right]_{q^2}$ is equal to 0. As in the proof of the second equality, we may treat only $t \geq -n+4$. We compute the coefficient of the desired equality as follows.

$$\begin{aligned}
&\sum_{r=0}^{4} (C_{n-2}^{2,n})_r \begin{bmatrix} 4 \\ r \end{bmatrix}_{q^2}^{\frac{1}{2}} q^{r(r-2t-4)} \begin{bmatrix} n+t \\ r \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-r \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&= q^{-2(n-1)} \begin{bmatrix} 2n \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&\quad - q^{-(n-1)} q^{-2t-3} \sqrt{\frac{(4)_q}{(2)_q}} \begin{bmatrix} 4 \\ 1 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} n+t \\ 1 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-1 \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&\quad + q^{2(-2t-2)} \sqrt{\frac{(4)_q(3)_q}{(2n)_q(2n-1)_q}} \begin{bmatrix} 4 \\ 2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} n+t \\ 2 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-2 \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&\quad - q^{n-1} q^{3(-2t-1)} \sqrt{\frac{(4)_q!}{(2n)_q \cdots (2n-2)_q}} \begin{bmatrix} 4 \\ 3 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} n+t \\ 3 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-3 \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&\quad + q^{2(n-1)} q^{-8t} \sqrt{\frac{(4)_q!}{(2n)_q \cdots (2n-3)_q}} \begin{bmatrix} n+t \\ 4 \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ n-t \end{bmatrix}_{q^2}^{\frac{1}{2}} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\[1em]
&= q^{\frac{1}{2}n^2 - \frac{1}{2}t^2 - 2t - 4} \sqrt{\frac{(n+t)_q!}{(n-t)_q!}} (n+t)_q!^{-1} \frac{\sqrt{(2n)_q!}}{(2n)_q \cdots (2n-3)_q} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&\quad \cdot \Big\{ q^{-2n+2t+6} (2n)_q \cdots (2n-3)_q - q^{-n+t+3} (4)_q (2n-1)_q \cdots (2n-3)_q (n+t)_q \\
&\qquad + \frac{(4)_q(3)_q}{(2)_q} (2n-2)_q (2n-3)_q (n+t)_q (n+t-1)_q \\
&\qquad - q^{n-t-3} (4)_q (2n-3)_q (n+t)_q \cdots (n+t-2)_q \\
&\qquad + q^{2n-2t-6} (n+t)_q \cdots (n+t-3)_q \Big\} \\
&= q^{\frac{1}{2}n^2 - \frac{1}{2}t^2 - 2t - 4} \sqrt{\frac{(n+t)_q!}{(n-t)_q!}} \frac{1}{(n+t)_q!} \frac{\sqrt{(2n)_q!}}{(2n)_q \cdots (2n-3)_q} \begin{bmatrix} 2n-4 \\ t+2 \end{bmatrix}_{q^2}^{-\frac{1}{2}} \\
&\quad \cdot q^{-6n-2t+6} (n-t)_q \cdots (n-t-3)_q
\end{aligned}$$

$$=q^{-4n-2t+2}\sqrt{\frac{(n-t)_q\cdots(n-t-3)_q}{(2n)_q\cdots(2n-3)_q}}.$$

□

Let n be the smallest odd integer which satisfies $\left[\frac{n}{m}\right]=1$. Then π_n appears in the spectral pattern of A once and π_{n-2} does not. Therefore $\Psi_{n-2}: X_2\times X_n \longrightarrow X_{n-2}$ is a 0-map. Let $\eta=\sum_{t\in I_n} d_t\boldsymbol{w}_t^n$ be a self-conjugate π_n-eigenvector. From the self-conjugacy we obtain $d_{-t}=(-q)^t\overline{d_t}$ for all $t\in I_n$. We shall show the complex numbers $\{d_t\}_{t\in I_n}$ are all 0. It derives non-existence of type D_m^* for $m\geq 2$.

Lemma 6.7. *The complex numbers $\{d_t\}_{t\in I_n}$ satisfy the following recurrence formula for $t\in I_n=\{-n,-n+1,\dots,n-1,n\}$.*

$$\begin{aligned}&q^{2n-1}(2)_q\sqrt{(n+t+2)_q(n+t+1)_q(n+t)_q(n+t-1)_q}\,d_{t+2}\\&-(4)_q\sqrt{(n+t)_q(n+t-1)_q(n-t)_q(n-t-1)_q}\,d_t\\&+q^{-2n+1}(2)_q\sqrt{(n-t+2)_q(n-t+1)_q(n-t)_q(n-t-1)_q}\,d_{t-2}=0,\end{aligned}$$

where $d_t=0$ if $|t|\geq n+1$.

Proof. Take the π_2-eigenvector $\xi_0^2=\sum_{s\in I_2}c_s\boldsymbol{w}_s$ where $c_{\pm 2}=q^{\mp 1}\sqrt{(3)_q!}$, $c_{\pm 1}=0$ and $c_0=-\sqrt{(4)_q}$. Then we have $\Psi_{n-2}(\xi_0^2,\zeta)=0$. Multiplying $q^{2n-4}\sqrt{(2n)_q\cdots(2n-3)_q}$ to the left hand side, we can derive

$$\begin{aligned}\sum_{t\in I_n}\Big(&c_{-2}q^{2n-2t-2}\sqrt{(n+t+2)_q\cdots(n+t-1)_q}\,d_{t+2}\\&+c_0q^{-2t}\sqrt{\frac{(4)_q(3)_q}{(2)_q}}\sqrt{(n+t)_q(n+t-1)_q(n-t)_q(n-t-1)_q}\,d_t\\&+c_2q^{-2n-2t+2}\sqrt{(n-t+2)_q\cdots(n-t-1)_q}\,d_{t-2}\Big)\boldsymbol{w}_t^{n-2}=0.\end{aligned}$$

Then we obtain the desired formula. □

Let us write

$$\begin{aligned}\alpha_t&=q^{2n-1}(2)_q\sqrt{(n+t+2)_q(n+t+1)_q(n+t)_q(n+t-1)_q}\,,\\\beta_t&=-(4)_q\sqrt{(n+t)_q(n+t-1)_q(n-t)_q(n-t-1)_q}\,,\\\gamma_t&=q^{-2n+1}(2)_q\sqrt{(n-t+2)_q(n-t+1)_q(n-t)_q(n-t-1)_q}\,.\end{aligned}$$

Then $\{d_t\}_{t\in I_n}$ satisfies

$$\alpha_t d_{t+2}+\beta_t d_t+\gamma_t d_{t-2}=0\,. \tag{6.17}$$

By definition we obtain $\alpha_{-t} = q^{4n-2}\gamma_t$ and $\beta_{-t} = \beta_t$. From the above recurrence formula at $-t$, we obtain

$$q^{4n-2}\gamma_t d_{-t+2} + \beta_t d_{-t} + q^{-4n+2}\alpha_t d_{-t-2} = 0\,.$$

Since we have assumed the self-conjugacy $d_{-t} = (-q)^t\overline{d_t}$, we obtain the following formula.

$$q^{-4n+4}\alpha_t d_{t+2} + \beta_t d_t + q^{4n-4}\gamma_t d_{t-2} = 0\,. \tag{6.18}$$

Through formulae (6.17) and (6.18), we obtain

$$d_{t+2} = q^{-2}\sqrt{\frac{(n-t+2)_q\cdots(n-t-1)_q}{(n+t+2)_q\cdots(n+t-1)_q}}\,d_{t-2} \tag{6.19}$$

for $t \geq -n+2$. Again from (6.17), we have $\beta_t d_t = (-1-q^{4n-4})\gamma_t d_{t-2}$. Hence for $|t| \leq n-2$, we obtain the following formula.

$$d_t = q^{-1}(q^{-2n+2}+q^{2n-2})\frac{(2)_q}{(4)_q}\sqrt{\frac{(n-t+2)_q(n-t+1)_q}{(n+t)_q(n+t-1)_q}}\,d_{t-2}\,. \tag{6.20}$$

Then we consider the above equalities (6.19) and (6.20). For $-n+2 \leq t \leq n-4$ we obtain by using (6.20) twice

$$d_{t+2} = q^{-2}(q^{-2n+2}+q^{2n-2})^2\frac{(2)_q^2}{(4)_q^2}\sqrt{\frac{(n-t+2)_q\cdots(n-t-1)_q}{(n+t+2)_q\cdots(n+t-1)_q}}\,d_{t-2}\,. \tag{6.21}$$

By (6.19) and (6.21) we have a equality

$$d_{t-2} = (q^{-2n+2}+q^{2n-2})^2\frac{(2)_q^2}{(4)_q^2}\,d_{t-2} \tag{6.22}$$

for $-n+2 \leq t \leq n-4$. We easily see that $1 = (q^{-2n+2}+q^{2n-2})^2\frac{(2)_q^2}{(4)_q^2}$ holds if and only if $n = 0$ or 2, however, this does not occur because n is an odd number. Thus we have $d_{t-2} = 0$ for $-n+2 \leq t \leq n-4$. It follows $d_t = 0$ for $-n \leq t \leq n-6$. For $n \geq 7$ we can derive $d_t = 0$ immediately. So we have to consider the cases $n = 3$ and $n = 5$.

(1) $n = 5$ case. We have already known $d_t = 0$ for $-5 \leq t \leq -1$. Using (6.20), we have $d_0 = 0$. Hence we have $d_t = 0$ for all $t \in I_5$.

(2) $n = 3$ case. We have already known $d_{-3} = 0$. Using (6.20) for $t = -1, 1$, we have $d_t = 0$ for odd t. From (6.19) and (6.20) for $t = 0$ we obtain

$$\begin{aligned} d_2 &= q^{-2}d_{-2}\,,\\ d_0 &= q^{-1}(q^{-4}+q^4)\frac{(2)_q}{(3)_q}\sqrt{\frac{(5)_q(4)_q}{(3)_q(2)_q}}\,d_{-2}\,. \end{aligned}$$

This shows d_{-2} and d_2 are real numbers. Now we consider a π_1-eigenvector $\Psi_1(\eta,\eta)=0$. As in the case A_4^*, we obtain

$$\begin{aligned}\sqrt{(6)_q}f\cdot\Psi_1(\eta,\eta)_{-1}=\,&q^{-2}(6)_q\eta_3\eta_{-3}+\big(-q^{-1}(5)_q(2)_q+q^{-3}(6)_q\big)\eta_2\eta_{-2}\\&+\big((4)_q(3)_q-q^{-2}(5)_q(2)_q\big)\eta_1\eta_{-1}+(q^{-1}-q)(4)_q(3)_q\eta_0^2\\&+(q^2(5)_q(2)_q-(4)_q(3)_q)\eta_{-1}\eta_1+(q(5)_q(2)_q-q^3(6)_q)\eta_{-2}\eta_2\\&-q^2(6)_q\eta_{-3}\eta_3\,.\end{aligned}$$

Applying $\pi_{\mathbb{T}}$ to the above both sides, we have

$$0=\big((q-q^{-1})(5)_q(2)_q+(q^{-3}-q^3)(6)_q\big)d_{-2}d_2+(q^{-1}-q)(4)_q(3)_qd_0^2\,.$$

Since d_{-2} and d_0 are real numbers, we obtain

$$q^{-2}((6)_q(3)_q-(5)_q(2)_q)d_{-2}^2+(4)_q(3)_qd_0^2=0\,,$$

however, the left hand side is positive because the above d_t are real. Hence we can get $d_{-2}=d_0=d_2=0$.

(VI) $\mathbb{T}_m\,(m\geq 2)$ case. We treat a C^*-algebra A whose spectral pattern is one of the following

$\mathbb{T}_{2\ell-1}$ $(\ell\geq 2)$ $\oplus_{k\in\mathbb{Z}_{\geq 0}}\big(1+2\big[\frac{k}{2\ell-1}\big]\big)\pi_k\oplus\oplus_{k\in\mathbb{Z}_{\geq 0}}2\big[\frac{k+\ell}{2\ell-1}\big]\pi_{k+\frac{1}{2}}$,

$\mathbb{T}_{2\ell}$ $(\ell\geq 1)$ $\oplus_{k\in\mathbb{Z}_{\geq 0}}(1+2\big[\frac{k}{\ell}\big])\pi_k$.

If A is of type $\mathbb{T}_2$, then we can easily derive $A=C(\mathbb{T}_2\setminus SU_q(2))$. Hence we study the case $\mathbb{T}_m$ for $m\geq 3$. In the cases, the π_1-multiplicity are one, so the linear subspace A_{π_1} generates a quantum sphere $C(S^2_{q,\lambda})$. We shall prove this is the canonical homogeneous sphere $C(\mathbb{T}\setminus SU_q(2))$, that is, the parameter $\lambda_0=(q^{-1}-q)^{-1}\lambda$ is equal to 1. In order to do this, we take the same strategy as in the case of type D_m^*. We have to prepare the following lemma which is proved in similar way to Lemma 6.6 and we omit the proof. Recall a π_1-eigenvector of $C(S^2_{q,\lambda})$

$$\xi^1_\lambda=q^{\frac{1}{2}}\sqrt{\frac{1-\lambda_0^2}{(2)_q}}\boldsymbol{w}^1_{-1}+\lambda_0\boldsymbol{w}^1_0-q^{-\frac{1}{2}}\sqrt{\frac{1-\lambda_0^2}{(2)_q}}\boldsymbol{w}^1_1.$$

Lemma 6.8. *Let n be a half integer with $n\geq 1$. Then we have the following equalities for all $t\in I_n$.*

$$\Psi_{n-1}(\boldsymbol{w}^1_{-1},\boldsymbol{w}^n_t)=q^{-t+2}\sqrt{\tfrac{(n+t)_q(n+t-1)_q}{(2n)_q(2n-1)_q}}\boldsymbol{w}^{n-1}_{t-1}\,,$$

$$\Psi_{n-1}(\boldsymbol{w}^1_0,\boldsymbol{w}^n_t)=-q^{-n-t+1}\sqrt{\tfrac{(2)_q(n-t)_q(n+t)_q}{(2n)_q(2n-1)_q}}\boldsymbol{w}^{n-1}_t\,,$$

$$\Psi_{n-1}(\boldsymbol{w}^1_1,\boldsymbol{w}^n_t)=q^{-2n-t}\sqrt{\tfrac{(n-t)_q(n-t-1)_q}{(2n)_q(2n-1)_q}}\boldsymbol{w}^{n-1}_{t+1}\,.$$

We immediately derive the following lemma.

Lemma 6.9. *Let n be a half integer with $n \geq 1$. Then we have the following equalities for all $t \in I_n$.*

$$\begin{aligned}
&q^{n-1}\sqrt{(2)_q(2n)_q(2n-1)_q}\,\Psi_{n-1}(\xi^1_\lambda, \boldsymbol{w}^n_t)\\
&\qquad\qquad\qquad = \; q^{n-t+\frac{3}{2}}\sqrt{1-\lambda_0^2}\,\sqrt{(n+t)_q(n+t-1)_q}\,\boldsymbol{w}^{n-1}_{t-1}\\
&\qquad\qquad\qquad\quad - q^{-t}(2)_q\lambda_0\sqrt{(n-t)_q(n+t)_q}\,\boldsymbol{w}^{n-1}_t\\
&\qquad\qquad\qquad\quad - q^{-n-t-\frac{3}{2}}\sqrt{1-\lambda_0^2}\,\sqrt{(n-t)_q(n-t-1)_q}\,\boldsymbol{w}^{n-1}_{t+1}\,.
\end{aligned}$$

We need another lemma whose proof is also same as for Lemma 6.7.

Lemma 6.10. *Let n be a half integer with $n \geq 1$ and $\eta = \sum_{t\in I_n} d_t\boldsymbol{w}^n_t$ be a π_n-eigenvector of $C(SU_q(2))$ such that $\Psi_{n-1}(\xi^1_\lambda, \eta) = 0$. Then $\{d_t\}_{t\in I_n}$ satisfies the following recurrence equation for all $t \in I_n = \{-n, -n+1, \ldots, n-1, n\}$.*

$$\begin{aligned}
&q^{n+\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{(n+t+1)_q(n+t)_q}\,d_{t+1}\\
&- (2)_q\lambda_0\sqrt{(n+t)_q(n-t)_q}\,d_t\\
&- q^{-n-\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{(n-t+1)_q(n-t)_q}\,d_{t-1} = 0
\end{aligned}$$

where we define $d_t = 0$ if $|t| \geq n+1$.

We show $A = C(\mathbb{T}_m \setminus SU_q(2))$ for in each case of odd and even order cyclic groups.

(1) $\mathbb{T}_{2\ell-1}$ $(\ell \geq 2)$ case. We focus on $\pi_{\ell-\frac{1}{2}}$-spectral subspace. Write n for $\ell - \frac{1}{2}$. Since π_n-eigenvector space is two-dimensional and π_{n-1}-eigenvector space is 0, the map $\Psi_{n-1}(\xi^1_\lambda, \cdot) : X_{\pi_n} \longrightarrow X_{\pi_{n-1}}$ is 0-map. Let $\eta = \sum_{t\in I_n} d_t\boldsymbol{w}^n_t$ be a non-zero π_n-eigenvector for A. Recall its conjugate eigenvector $T\eta = \sum_{t\in I_n}(-q)^{-t}\overline{d_{-t}}\boldsymbol{w}^n_t$, where we have defined $(-q)^s = \sqrt{-1}^{2s}|q|^s$ for all real number s. Then η and $T\eta$ are in the kernel of $\Psi_{n-1}(\xi^1_\lambda, \cdot)$. By Lemma 6.10 we obtain the following recurrence equation.

$$\alpha_t d_{t+1} + \beta_t d_t + \gamma_t d_{t-1} = 0 \tag{6.23}$$

$$q^{-2n}\alpha_t d_{t+1} + \beta_t d_t + q^{2n}\gamma_t d_{t-1} = 0\,, \tag{6.24}$$

where we put

$$\begin{aligned}
\alpha_t &= q^{n+\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{(n+t+1)_q(n+t)_q}\,,\\
\beta_t &= -\,(2)_q\lambda_0\sqrt{(n+t)_q(n-t)_q}\,,\\
\gamma_t &= -\,q^{-n-\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{(n-t+1)_q(n-t)_q}\,.
\end{aligned}$$

Assume that λ_0 is not equal to 1. Then we know $\alpha_t = 0$ if and only if $t = -n$ and $\gamma_t = 0$ if and only if $t = n$. By (6.23) and (6.24) we obtain

$$\alpha_t d_{t+1} = q^{2n} \gamma_t d_{t-1} \,. \tag{6.25}$$

Hence again by (6.23) we have

$$\beta_t d_t = -(1 + q^{2n}) \gamma_t d_{t-1} \,.$$

Assume λ_0 is equal to 0. Then the above equality shows $d_{t-1} = 0$ for $-n \leq t \leq n-1$. By (6.25) d_{n-1} and d_n are also equal to 0, however, this is a contradiction to non triviality of η. Next we consider the case $0 < \lambda_0 < 1$. From the above equality d_t for $-n+1 \leq t \leq n-1$ are uniquely determined by d_{-n}. The number d_n is determined by the previous equation as $d_n = q^{2n} \alpha_{n-1}^{-1} \gamma_{n-1} d_{n-2}$ where we use the fact that α_{n-1} is never 0 for $n \geq \frac{3}{2}$. Therefore $(d_t)_{t \in I_n}$ is a scalar multiple of a vector. In particular the π_n-eigenvector space X_{π_n} becomes one-dimensional. This is a contradiction. It concludes λ_0 is equal to 1 in this case. From simple computation we obtain $X_n = \mathbb{C} \boldsymbol{w}_{-n}^n + \mathbb{C} \boldsymbol{w}_n^n$. We show $A = C(\mathbb{T}_{2\ell-1} \setminus SU_q(2))$. The C^*-algebra $C(\mathbb{T}_{2\ell-1} \setminus SU_q(2))$ is generated by the matrix elements $w(\pi_\nu)_{ns,t}$ for all $\nu \in \frac{1}{2}\mathbb{Z}_{\geq 0}$ and $t \in I_\nu$, where s is an integer such that $ns \in I_\nu$. If A contains $C(\mathbb{T}_{2\ell-1} \setminus SU_q(2))$, then they coincide because of its common spectral pattern. By $SU_q(2)$-invariance of A, it suffices to show $w(\pi_\nu)_{ns,-\nu} = \left[\begin{smallmatrix} 2\nu \\ \nu+ns \end{smallmatrix}\right]_{q^2}^{\frac{1}{2}} x^{\nu+ns} v^{\nu-ns}$ are contained in A for non-negative s. This is equal to $\left[\begin{smallmatrix} 2\nu \\ \nu+ns \end{smallmatrix}\right]_{q^2}^{\frac{1}{2}} x^{2ns} x^{\nu-ns} v^{\nu-ns}$. Since A contains $xv \in C(\mathbb{T} \setminus SU_q(2))$ and $x^{2n} = \left[\begin{smallmatrix} 2\nu \\ \nu+ns \end{smallmatrix}\right]_{q^2}^{-\frac{1}{2}} w(\pi_n)_{-n,-n}$, the element $x^{\nu+ns} v^{\nu-ns}$ is in A. This shows A is in fact $C(\mathbb{T}_{2\ell-1} \setminus SU_q(2))$.

(2) $\mathbb{T}_{2\ell}$ ($\ell \geq 2$) case. Write n for ℓ in this case. We analyze the recurrence equation in Lemma 6.10 under the condition that a π_n-eigenvector $\eta = \sum_{t \in I_n} d_t \boldsymbol{w}_t^n$ is self-conjugate. Assume that λ_0 is not equal to 1. When λ_0 is equal to 0, η is the zero vector as in the previous case. In the case $0 < \lambda_0 < 1$ we know that the space of its solution is one dimensional and we can assume that d_t are all real number. Then we have

$$d_t = -(1 + q^{2n}) \beta_t^{-1} \gamma_t d_{t-1}$$

for all $|t| \leq n-1$. For $-n+1 \leq t \leq n-2$ we obtain the following equality.

$$\begin{aligned} d_{-t} &= -(1+q^{2n}) \frac{\gamma_{-t}}{\beta_{-t}} d_{-t-1} \\ &= -(-q)^{t+1} (1+q^{2n}) \frac{\gamma_{-t}}{\beta_{-t}} d_{t+1} \\ &= (-q)^{t+1} (1+q^{2n})^2 \frac{\gamma_{-t}\gamma_{t+1}}{\beta_{-t}\beta_{t+1}} d_t \\ &= -q(1+q^{2n})^2 \frac{\gamma_{-t}\gamma_{t+1}}{\beta_{-t}\beta_{t+1}} d_{-t} \,. \end{aligned}$$

Since its coefficient $\frac{\gamma_{-t}\gamma_{t+1}}{\beta_{-t}\beta_{t+1}}$ is strictly positive, we have $d_t = 0$ for $-n+2 \leq t \leq n-1$. We now treat a positive integer $n \geq 2$. It deduces $-n+2 \leq n-2$ and we obtain $d_n = 0$ by (6.25). Hence we have $d_t = 0$ for all $t \in I_n$ because of $-n+2 \leq 0$ and the self-conjugacy of η. Under the condition $\lambda_0 < 1$ we have shown $\eta = 0$ if η is self-conjugate and $\Psi_{n-1}(\xi_\lambda^1, \eta) = 0$. Now we consider the map $\Psi_{n-1}(\xi_\lambda^1, \cdot) : X_n \longrightarrow X_{n-1}$. We have known X_n is three-dimensional and X_{n-1} is one-dimensional. Take two linearly independent π_n-eigenvectors ζ_1 and ζ_2 from X_n which satisfy $\Psi_{n-1}(\xi_\lambda^1, \zeta_1) = \Psi_{n-1}(\xi_\lambda^1, \zeta_2) = 0$. For conjugate eigenvectors $T\zeta_1$ and $T\zeta_2$, we can take complex numbers μ_1 and μ_2 which are defined by $\Psi_{n-1}(\xi_\lambda^1, T\zeta_1) = \mu_1 \Psi_{n-1}(\xi_\lambda^1, \xi_\lambda^n)$ and $\Psi_{n-1}(\xi_\lambda^1, T\zeta_2) = \mu_2 \Psi_{n-1}(\xi_\lambda^1, \xi_\lambda^n)$, where ξ_λ^ν is a self-conjugate π_ν-eigenvector for $C(S_{q,\lambda}^2)$. Note $\Psi_{n-1}(\xi_\lambda^1, \xi_\lambda^n)$ is not the zero vector. Multiplying complex numbers to ζ_1 and ζ_2 respectively, we may assume μ_1 and μ_2 are real numbers. If μ_1 is equal to 0, then the self-conjugate vectors $\zeta_1 + T\zeta_1$ and $\sqrt{-1}(\zeta_1 - T\zeta_1)$ are in the kernel of $\Psi_{n-1}(\xi_\lambda^1, \cdot)$. From the above discussion they are 0. This shows immediately $\zeta_1 = 0$. This is a contradiction. Hence we have a non-zero real number μ_1. Similarly we show μ_2 is non-zero. We may assume they are equal to 1. For $i = 1, 2$ we have $\Psi_{n-1}(\xi_\lambda^1, \zeta_i + T\zeta_i - \xi_\lambda^n) = 0$. Then it yields $\zeta_i + T\zeta_i = \xi_\lambda^n$ because of the self-conjugacy of $\zeta_i + T\zeta_i - \xi_\lambda^n$ for each i. It follows $\zeta_1 - \zeta_2 = -T(\zeta_1 - \zeta_2)$. In particular $T(\zeta_1 - \zeta_2)$ is in the kernel of $\Psi_{n-1}(\xi_\lambda^1, \cdot)$. This shows $\zeta_1 - \zeta_2 = 0$ but this is a contradiction to linear independence of ζ_1 and ζ_2. Therefore we can derive $\lambda_0 = 1$. Then we obtain $X_n = \mathbb{C}\boldsymbol{w}_{-n}^n + \mathbb{C}\boldsymbol{w}_0^n + \mathbb{C}\boldsymbol{w}_n^n$. As in the proof of the case $\mathbb{T}_{2\ell-1}$, we can prove $A = C(\mathbb{T}_{2\ell} \setminus SU_q(2))$.

7. CLASSIFICATION OF RIGHT COIDEALS OF $C(SU_q(2))$: $-1 < q < 0$ CASE

We use the same strategy as in the previous section in order to classify right coideals of $C(SU_q(2))$ for negative q. We prepare a positive parameter q_0 defined by $q_0 = -q$. Recall we have defined q-integer $(n)_q$ by q_0-integer, more precisely $(n)_q = (n)_{q_0}$ and $q^n = \sqrt{-1}^{2n} q_0^n$ for a half integer n.

Now we investigate each case of multiplicity diagrams. Contrary to the case of positive q, we have to treat other graphs which have a single loop at a vertex as listed in Appendix. We recall the notion of the quantum spheres $C(S_{q,\lambda}^2)$. We use an embedding it into $C(SU_q(2))$ by defining its π_1-eigenvector as

$$\xi_\lambda^1 = \left(q_0^{\frac{1}{2}} \sqrt{\frac{1-\lambda_0^2}{(2)_q}}, \lambda_0, q_0^{-\frac{1}{2}} \sqrt{\frac{1-\lambda_0^2}{(2)_q}} \right) w(\pi_1) .$$

If a right coideal A is of type $\mathbb{T}$, then it is in fact $\beta_z^L(C(S_{q,\lambda}^2))$ for some $z \in \mathbb{T}$ by Podleś' classification results on the quantum spheres [Po1, Theorem 1]. We know that $C(SO_q(3))$ and $C(SO_{-q}(3))$ are isomorphic as compact quantum groups. Hence we have the complete collection of right coideals which have only spectral subspaces with integer spins. In this way we can deny the existence of type A_4^*, S_4^*, A_5^* and $D_m^*\,(m \geq 2)$. For right coideals of D_∞^* and $\mathbb{T}_{2n}\,(n \geq 1)$ we have shown their uniqueness up to conjugation by β^L. Hence we have to investigate right coideals of type $\mathbb{T}_n$ (odd $n \geq 3$), D_n (odd $n \geq 1$), A'_m ($m \geq 3$). The main classification result is as follows.

Theorem 7.1. *Let $A \subset C(SU_q(2))$ be a right coideal. Then its multiplicity diagram is one of type $1, \mathbb{T}_n\,(n \geq 2), \mathbb{T}, SU(2), D^*_\infty$ and D_1. If it is of type $\mathbb{T}$, it is one of the quantum spheres. Otherwise it is unique up to conjugation by β^L.*

(I) $\mathbb{T}_{2\ell-1}\,(\ell \geq 2)$ case. We treat a C^*-algebra A whose spectral pattern is $\oplus_{k\in\mathbb{Z}_{\geq 0}}\big(1+2\big[\frac{k}{2\ell-1}\big]\big)\pi_k \oplus \oplus_{k\in\mathbb{Z}_{\geq 0}} 2\big[\frac{k+\ell}{2\ell-1}\big]\pi_{k+\frac{1}{2}}$. As in the previous section we study the C^*-subalgebra generated by π_1-spectral subspace and assert that is the canonical homogeneous sphere. We begin classifying this case by stating a negative q version of Lemma 6.8.

Lemma 7.2. *Let n be a half integer with $n \geq 1$. Then we have the following equalities for all $t \in I_n$.*

$$\Psi_{n-1}(\boldsymbol{w}^1_{-1}, \boldsymbol{w}^n_t) = q_0^{-t+2}\sqrt{\tfrac{(n+t)_q(n+t-1)_q}{(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_{t-1}\,,$$

$$\Psi_{n-1}(\boldsymbol{w}^1_{0}, \boldsymbol{w}^n_t) = -(-1)^{n-t}q_0^{-n-t+1}\sqrt{\tfrac{(2)_q(n-t)_q(n+t)_q}{(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_{t}\,,$$

$$\Psi_{n-1}(\boldsymbol{w}^1_{1}, \boldsymbol{w}^n_t) = q_0^{-2n-t}\sqrt{\tfrac{(n-t)_q(n-t-1)_q}{(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_{t+1}\,.$$

If necessary, we consider the conjugation by β^L and may assume $C(S^2_{q,\lambda}) \subset A$.

Lemma 7.3. *Let n be a half integer with $n \geq 1$. Then we get the following equality for $t \in I_n$.*

$$\begin{aligned}\Psi_{n-1}(\xi^1_\lambda, \boldsymbol{w}^n_t) =\, & q_0^{-t+\frac{5}{2}}\sqrt{1-\lambda_0^2}\sqrt{\frac{(n+t)_q(n+t-1)_q}{(2)_q(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_{t-1}\\ & -(-1)^{n-t}q_0^{-n-t+1}\lambda_0\sqrt{\frac{(2)_q(n-t)_q(n+t)_q}{(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_{t}\\ & + q_0^{-2n-t-\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{\frac{(n-t)_q(n-t-1)_q}{(2)_q(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_{t+1}\,.\end{aligned}$$

Lemma 7.4. *Let n be a half integer with $n \geq 1$ and $\eta = \sum_{t\in I_n} d_t \boldsymbol{w}^n_t$ be a π_n-eigenvector with $\Psi_{n-1}(\xi^1_\lambda, \eta) = 0$. Then we get the following recurrence equation for $t \in I_n$.*

$$\begin{aligned}& q_0^{n+\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{(n+t)_q(n+t+1)_q}\,d_{t+1}\\ & -(-1)^{n-t}\lambda_0(2)_q\sqrt{(n-t)_q(n+t)_q}\,d_t\\ & + q_0^{-n-\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{(n-t)_q(n-t+1)_q}\,d_{t-1} = 0\,,\end{aligned}$$

where we define $d_t = 0$ for $|t| \geq n+1$.

We write n for $\ell-\frac{1}{2}$. Then π_n-eigenvector space of A is two-dimensional. We may assume π_1-part generates a quantum sphere $C(S^2_{q,\lambda})$ as usual. We derive $\lambda_0 = 1$ and it immediately derives $A = C(\mathbb{T}_{2\ell-1}\backslash C(SU_q(2)))$. Now Let $\eta = \sum_{t\in I_n} d_t\boldsymbol{w}^n_t$ be a π_n-eigenvector of A. By the absence of π_{n-1}-part we have $\Psi_{n-1}(\xi^1_\lambda, \eta) = 0 = \Psi_{n-1}(\xi^1_\lambda, T\eta)$. From the previous lemmas we get

$$\begin{aligned}\alpha_t d_{t+1} + \beta_t d_t + \gamma_t d_{t-1} &= 0\,,\\ q_0^{-2n}\alpha_t d_{t+1} - \beta_t d_t + q_0^{2n}\gamma_t d_{t-1} &= 0\,,\end{aligned}$$

where we put

$$\begin{aligned}\alpha_t &= q_0^{n+\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{(n+t)_q(n+t+1)_q}\,,\\ \beta_t &= -(-1)^{n-t}\lambda_0(2)_q\sqrt{(n-t)_q(n+t)_q}\,,\\ \gamma_t &= q_0^{-n-\frac{1}{2}}\sqrt{1-\lambda_0^2}\sqrt{(n-t)_q(n-t+1)_q}\,.\end{aligned}$$

Then we can prove $(d_t)_{t\in I_n}$ is a scalar multiple of a vector as in the $\mathbb{T}_{2\ell-1}$ case in the previous section. This is a contradiction. Hence we have proved $\lambda_0 = 1$ and $A = C(\mathbb{T}_{2\ell-1}\setminus SU_q(2))$.

(II) D_n (odd $n \geq 3$) case. Before proof of the non-existence of this case, we shall state some basic lemmas without proofs which are proved similarly by direct calculations as before.

We state a negative q and Ψ_{n-1} version of Lemma 6.6.

Lemma 7.5. *Let n be a half integer with $n \geq \frac{3}{2}$. Then we have the following equalities for all $t \in I_n$.*

$$\begin{aligned}\Psi_{n-1}(\boldsymbol{w}^2_{-2}, \boldsymbol{w}^n_t) &= (-1)^{n-t}q_0^{-\frac{1}{2}n-2t+6}\frac{1}{(2n-2)_q}\\ &\qquad\sqrt{\frac{(4)_q(n-t+1)_q(n+t)_q(n+t-1)_q(n+t-2)_q}{(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_{t-2}\,,\\ \Psi_{n-1}(\boldsymbol{w}^2_0, \boldsymbol{w}^n_t) &= (-1)^{n-t}q_0^{-\frac{3}{2}n-2t+3}\frac{q_0^{-n-1}(n-t-1)_q - q_0^{n+1}(n+t-1)_q}{(2n-2)_q}\\ &\qquad\sqrt{\frac{(3)_q!(n+t)_q(n-t)_q}{(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_t\,,\\ \Psi_{n-1}(\boldsymbol{w}^2_2, \boldsymbol{w}^n_t) &= -(-1)^{n-t}q_0^{-\frac{5}{2}n-2t}\frac{1}{(2n-2)_q}\\ &\qquad\sqrt{\frac{(4)_q(n+t+1)_q(n-t)_q(n-t-1)_q(n-t-2)_q}{(2n)_q(2n-1)_q}}\,\boldsymbol{w}^{n-1}_{t+2}\,.\end{aligned}$$

We take a vector $\xi_0^2 = (q_0\sqrt{(3)_q!}, 0, -\sqrt{(4)_q}, 0, q_0^{-1}\sqrt{(3)_q!}\,)\, w(\pi_2)$ as a π_2-eigenvector of $C(S_{q,0}^2)$. Recall its entries generate the right coideal of type D_∞^*. We denote it by $A_{D_\infty^*}$.

Lemma 7.6. *Let n be a half integer with $n \geq \frac{3}{2}$. Then we have the following equalities for all $t \in I_n$.*

$$\begin{aligned}
&-q_0^{\frac{3}{2}n-3}\sqrt{\frac{(2n)_q(2n-1)_q}{(4)_q!}}\Psi_{n-1}(\xi_0^2, \boldsymbol{w}_t^n) \\
&\quad= -(-1)^{n-t}q_0^{n-2t+4}\sqrt{(n-t+1)_q(n+t)_q(n+t-1)_q(n+t-2)_q}\,\boldsymbol{w}_{t-2}^{n-1} \\
&\qquad+ (-1)^{n-t}q_0^{-2t}(q_0^{-n-1}(n-t-1)_q - q_0^{n+1}(n+t-1)_q)\sqrt{(n+t)_q(n-t)_q}\boldsymbol{w}_t^{n-1} \\
&\qquad+ (-1)^{n-t}q_0^{-n-2t-4}\sqrt{(n+t+1)_q(n-t)_q(n-t-1)_q(n-t-2)_q}\,\boldsymbol{w}_{t+2}^{n-1}\,.
\end{aligned}$$

Lemma 7.7. *Let n be a half integer with $n \geq \frac{3}{2}$ and $\eta = \sum_{t\in I_n} d_t\boldsymbol{w}_t^n$ be a π_n-eigenvector of A such that it satisfies $\Psi_{n-1}(\xi_0^2, \eta) = 0$. Then we have the following recurrence equation for $t \in I_n$.*

$$\begin{aligned}
&-q_0^n\sqrt{(n-t-1)_q(n+t+2)_q(n+t+1)_q(n+t)_q}\,d_{t+2} \\
&+(q_0^{-n-1}(n-t-1)_q - q_0^{n+1}(n+t-1)_q)\sqrt{(n+t)_q(n-t)_q}\,d_t \\
&+q_0^{-n}\sqrt{(n+t-1)_q(n-t+2)_q(n-t+1)_q(n-t)_q}\,d_{t-2} = 0,
\end{aligned}$$

where we define $d_t = 0$ for $|t| \geq n+1$.

Lemma 7.8. *Let n be a half integer in $\frac{3}{2} + \mathbb{Z}_{\geq 0}$ and A be a right coideal in $C(SU_q(2))$ such that A contains $A_{D_\infty^*}$. If a π_n-eigenvector η of A satisfies $\Psi_{n-1}(\xi_0^2, \eta) = 0$ and $\Psi_{n-1}(\xi_0^2, T\eta) = 0$, then η is the zero vector.*

Proof. Take complex numbers $\{d_t\}_{t\in I_n}$ with $\eta = \sum_{t\in I_n} d_t\boldsymbol{w}_t^n$. Note that $T\eta = \sum_{t\in I_n}(-q)^{-t}\overline{d_{-t}}\boldsymbol{w}_t^n$. Let us prepare the following notations α_t, β_t and γ_t.

$$\begin{aligned}
\alpha_t &= -q_0^n\sqrt{(n-t-1)_q(n+t+2)_q(n+t+1)_q(n+t)_q} \\
\beta_t &= (q_0^{-n-1}(n-t-1)_q - q_0^{n+1}(n+t-1)_q)\sqrt{(n+t)_q(n-t)_q} \\
\gamma_t &= q_0^{-n}\sqrt{(n+t-1)_q(n-t+2)_q(n-t+1)_q(n-t)_q}\,.
\end{aligned}$$

From our assumption on η and $T\eta$, we get the following recurrence equations for $t \in I_n$ by the previous lemma.

$$\begin{aligned}
\alpha_t d_{t+2} + \beta_t d_t + \gamma_t d_{t-2} &= 0\,, \\
q_0^{-2n+2}\alpha_t d_{t+2} - \beta_{-t}d_t + q_0^{2n-2}\gamma_t d_{t-2} &= 0\,.
\end{aligned} \tag{7.1}$$

In the above equations put $t = -n + 1$ and we get

$$-q_0^n \sqrt{(2n-2)_q (3)_q!}\, d_{-n+3} + q_0^{-n-1} (2n-2)_q \sqrt{(2n-1)_q}\, d_{-n+1} = 0\,,$$
$$-q_0^{-n+2} \sqrt{(2n-2)_q (3)_q!}\, d_{-n+3} + q_0^{n+1} (2n-2)_q \sqrt{(2n-1)_q}\, d_{-n+1} = 0\,.$$

This derives $d_{-n+1} = d_{-n+3} = 0$ because $0 < q_0 < 1$. We know $\alpha_t = 0$ if and only if $t = -n, n-1$. By using (7.1) inductively, we get $d_{-n+2k-1} = 0$ for $k = 1, \dots n + \frac{1}{2}$. Similarly we get $d_{n-2k+1} = 0$ for $k = 1, \dots n + \frac{1}{2}$. Hence we have proved $\eta = 0$. □

Let A be a right coideal of type D_n (odd $n \geq 3$). Then its spectral pattern is

$$\bigoplus_{k \in \mathbb{Z}_{\geq 0}} \left(\left[\frac{k}{n}\right] + \frac{1 + (-1)^k}{2} \right) \pi_k \oplus \bigoplus_{k \in \mathbb{Z}_{\geq 0}} \left(\left[\frac{2k+1}{n}\right] - \left[\frac{k}{n}\right] \right) \pi_{k+\frac{1}{2}}\,.$$

If we look at the integer spin spectral pattern, then we have $\pi_0 \oplus \pi_2 \oplus \cdots$. Therefore the π_2-spectral subspace A_{π_2} generates a right coideal of type D_∞^*. Considering $\beta_z^L(A)$ for some $z \in \mathbb{T}$, we may assume A contains $A_{D_\infty^*}$. Next look at the half integer part and we see that $\pi_{\frac{n}{2}}$ appears once and $\pi_{\frac{n}{2}-1}$ does not there. Then we can make use of Lemma 7.8 and conclude such a right coideal does not exist.

(III) D_1 case. We show the existence of this case and its uniqueness up to conjugation. In order to do we need some elementary lemmas.

Lemma 7.9. *Take a positive integer n. Then we have the following equalities.*

$$f \cdot x^n = \sqrt{-1}^{\,-n+1} q_0^{\frac{n-1}{2}} (n)_q x^{n-1} u\,, \quad f \cdot v^n = \sqrt{-1}^{\,-n+1} q_0^{\frac{n-1}{2}} (n)_q v^{n-1} y\,.$$

Proof. We know $x^n = w(\pi_{\frac{n}{2}})_{-\frac{n}{2}, -\frac{n}{2}}$. Hence we have:

$$\begin{aligned} f \cdot x^n &= \sqrt{-1}^{\,-n+1} \sqrt{(n)_q}\, w(\pi_{\frac{n}{2}})_{-\frac{n}{2}, -\frac{n}{2}+1} \\ &= \sqrt{-1}^{\,-n+1} \sqrt{(n)_q} \begin{bmatrix} n \\ 1 \end{bmatrix}^{\frac{1}{2}} x^{n-1} u \\ &= \sqrt{-1}^{\,-n+1} q_0^{\frac{n-1}{2}} (n)_q x^{n-1} u\,. \end{aligned}$$

Similarly we get the assertion on $f \cdot v^n$. □

Lemma 7.10. *For $r, s \geq 0$ we have the following equality.*

$$h(x^r y^r u^s v^s) = q_0^{r(s+1)} \frac{(r)_q! (s)_q!}{(r+s+1)_q!}\,.$$

Proof. Recall $f \cdot y^r = f \cdot u^{s-1} = 0$ and we get

$$\begin{aligned}
f \cdot x^{r+1} y^r u^{s-1} v^s &= (f \cdot x^{r+1}) \cdot (k \cdot y^r u^{s-1} v^s) + (k^{-1} \cdot x^{r+1}) \cdot (f \cdot y^r u^{s-1} v^s) \\
&= \sqrt{-1}^{\,-r} q_0^{\frac{r}{2}} (r+1)_q \cdot q^{-\frac{r}{2} - \frac{s-1}{2} + \frac{s}{2}} x^r u y^r u^{s-1} v^s \\
&\quad + q^{\frac{-(r+1)}{2}} \cdot q^{\frac{r}{2} + \frac{s-1}{2}} \sqrt{-1}^{\,-s+1} q_0^{\frac{s}{2}} (s)_q x^{r+1} y^r u^{s-1} v^{s-1} y \\
&= \sqrt{-1}^{\,-r} q_0^{\frac{r}{2}} (r+1)_q \cdot q^{-\frac{3r}{2} + \frac{1}{2}} x^r y^r u^s v^s \\
&\quad + q^{\frac{s}{2} - 1} \sqrt{-1}^{\,-s+1} q_0^{\frac{s}{2}} (s)_q q^{-2(s-1)} x^{r+1} y^{r+1} u^{s-1} v^{s-1} \\
&= \sqrt{-1}\, q_0^{-r+\frac{1}{2}} (r+1)_q \cdot x^r y^r u^s v^s \\
&\quad - \sqrt{-1}\, q_0^{-s+\frac{1}{2}} (s)_q x^{r+1} y^{r+1} u^{s-1} v^{s-1} .
\end{aligned}$$

The Haar state h has the property that $h(f \cdot a) = 0$ for $a \in A(SU_q(2))$. Apply h to the above both sides and we have

$$h(x^{r+1} y^{r+1} u^{s-1} v^{s-1}) = q_0^{s-r} \frac{(r+1)_q}{(s)_q} h(x^r y^r u^s v^s) .$$

Inductively we can calculate as desired. $\square$

Recall the $\pi_{\frac{1}{2}}$-spectral subspace $Y_{\frac{1}{2}} = \mathbb{C}x + \mathbb{C}v + \mathbb{C}u + \mathbb{C}y$. Let $P_{\frac{1}{2}} : C(SU_q(2)) \longrightarrow Y_{\frac{1}{2}}$ be the projection introduced in the second section. It is an orthogonal projection with respect to the Haar state.

Lemma 7.11. *For $r, s \geq 0$ we have the following equalities.*

$$\begin{aligned}
P_{\frac{1}{2}}(x^r y^r u^s v^s u) &= q_0^{r(s+2)} \frac{(2)_q (r)_q! (s+1)_q!}{(r+s+2)_q!} u , \\
P_{\frac{1}{2}}(x^r y^r u^s v^s y) &= q_0^{r(s+1)-s} \frac{(2)_q (r+1)_q! (s)_q!}{(r+s+2)_q!} u .
\end{aligned}$$

Proof. Recall two maps $\beta^L = (\pi_{\mathbb{T}} \otimes \mathrm{id}) \circ \delta$ and $\beta^R = (\mathrm{id} \otimes \pi_{\mathbb{T}}) \circ \delta$ which act on x, v, u and y as follows.

$$\begin{aligned}
\begin{pmatrix} \beta_z^L(x) & \beta_z^L(u) \\ \beta_z^L(v) & \beta_z^L(y) \end{pmatrix} &= \begin{pmatrix} zx & zu \\ \bar{z}v & \bar{z}y \end{pmatrix} , \\
\begin{pmatrix} \beta_z^R(x) & \beta_z^R(u) \\ \beta_z^R(v) & \beta_z^R(y) \end{pmatrix} &= \begin{pmatrix} zx & \bar{z}u \\ zv & \bar{z}y \end{pmatrix} ,
\end{aligned}$$

where z runs on the torus $\mathbb{T} \subset \mathbb{C}$. Hence the above four elements have the different spectrums with respect to β^L and β^R. It is easy to see that the element $x^r y^r u^s v^s u$ has the same spectrum as u. Hence it has the expansion in $L^2(SU_q(2))$ $x^r y^r u^s v^s u = \lambda u + \cdots$ where λ is a complex number. By orthogonality of u and

$w(\pi_\nu)_{i,j}$ for $\nu \geq 1$ and $i,j \in I_\nu$, we get

$$\begin{aligned}\lambda &= h(uu^*)^{-1}h(x^ry^ru^sv^suu^*)\\ &= q_0(2)_q(-q^{-1})h(x^ry^ru^{s+1}v^{s+1})\\ &= q_0^{r(s+2)}\frac{(2)_q(r)_q!(s+1)_q!}{(r+s+2)_q!}\,.\end{aligned}$$

We also get the desired result on y in the same way.

□

We propose the following lemma. Later we see this equality guarantees the existence of the right coideal of type D_1. Let us write $p_r = \prod_{t=1}^{r}(1+q^{-t})$.

Lemma 7.12. *For $k \geq 0$ we have the following equality.*

$$\sum_{r=0}^{k}\begin{bmatrix}k\\ r\end{bmatrix}_q(-1)^{k-r}q^{r(r-k+1)}p_rp_{k-r+1} = q^{-k}\sum_{r=0}^{k}\begin{bmatrix}k\\ r\end{bmatrix}_q(-1)^{k-r}q^{r(r-k+2)}p_{r+1}p_{k-r}\,.$$

Moreover, this is precisely equal to $(1+q^{-1})(q^{-2};q^{-1})_k$.

Proof. Denote the above left and right hand side by a_k and b_k, respectively. We want to lead the recurrence formula of them. In computations below, we use the following equalities.

$$\begin{bmatrix}n+1\\ r\end{bmatrix}_q = \begin{bmatrix}n\\ r-1\end{bmatrix}_q + q^r\begin{bmatrix}n\\ r\end{bmatrix}_q\,, \quad p_{n+1} = p_n + q^{-n-1}p_n\,.$$

Change variables r to $k-r$ in the formula of a_k and b_k and we get

$$\begin{aligned}a_k &= q^k\sum_{r=0}^{k}\begin{bmatrix}k\\ r\end{bmatrix}_q(-1)^rq^{r(r-k-1)}p_{r+1}p_{k-r}\,,\\ b_k &= q^k\sum_{r=0}^{k}\begin{bmatrix}k\\ r\end{bmatrix}_q(-1)^rq^{r(r-k-2)}p_rp_{k-r+1}\,.\end{aligned}$$

For a_k we compute as follows.

$$\begin{aligned}
q^{-k-1}a_{k+1} =& \sum_{r=0}^{k+1} \begin{bmatrix} k+1 \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-2)} p_{r+1}p_{k-r+1} \\
=& \sum_{r=0}^{k+1} \left(\begin{bmatrix} k \\ r-1 \end{bmatrix}_q + q^r \begin{bmatrix} k \\ r \end{bmatrix}_q \right) (-1)^r q^{r(r-k-2)} p_{r+1}p_{k-r+1} \\
=& \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^{r+1} q^{(r+1)(r-k-1)} p_{r+2}p_{k-r} + \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-1)} p_{r+1}p_{k-r+1} \\
=& - q^{-k-1} \left(\sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^{r+1} q^{r(r-k)} (1+q^{-r-2}) p_{r+1}p_{k-r} \right) \\
& + \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-1)} (1+q^{-k+r-1}) p_{r+1}p_{k-r} \\
=& (1-q^{-k-3}) \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-1)} p_{r+1}p_{k-r} \\
=& q^{-k}(1-q^{-k-3}) a_k \, .
\end{aligned}$$

Hence we get $a_{k+1} = (q - q^{-k-2})a_k$. Similarly we compute b_{k+1} as follows.

$$\begin{aligned}
q^{-k-1}b_{k+1} =& \sum_{r=0}^{k+1} \begin{bmatrix} k+1 \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-3)} p_r p_{k-r+2} \\
=& \sum_{r=0}^{k+1} \left(\begin{bmatrix} k \\ r-1 \end{bmatrix}_q + q^r \begin{bmatrix} k \\ r \end{bmatrix}_q \right) (-1)^r q^{r(r-k-3)} p_r p_{k-r+2} \\
=& \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^{r+1} q^{(r+1)(r-k-2)} p_{r+1}p_{k-r+1} + \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-2)} p_r p_{k-r+2} \\
=& - q^{-k-2} \left(\sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-1)} (p_r + q^{-r-1}p_r) p_{k-r+1} \right) \\
& + \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-2)} (1+q^{-k+r-2}) p_r p_{k-r+1} \\
=& (-q^{-k-3}+1) \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^r q^{r(r-k-2)} p_r p_{k-r+1} \\
=& q^{-k}(1-q^{-k-3}) b_k \, .
\end{aligned}$$

Hence we get $b_{k+1} = (q - q^{-k-2})b_k$, which is the same form as a_{k+1}. We can easily check $a_0 = p_1 = b_0$ and it deduces the desired equality. $\square$

At last we prove the existence of a right coideal A of type D_1. It has a spectral pattern, $\oplus_{k\in\mathbb{Z}_{\geq 0}}\big(k+\frac{1+(-1)^k}{2}\big)\pi_k \oplus \oplus_{k\in\mathbb{Z}_{\geq 0}}(k+1)\pi_{k+\frac{1}{2}}$. If A really exists, its spectral subspace $A_{\pi_{\frac{1}{2}}}$ generates the whole C^*-algebra A. In fact the generated C^*-algebra contains $\pi_{\frac{1}{2}}$ and the only D_1 case admits such an algebra. We have to clarify a self-conjugate $\pi_{\frac{1}{2}}$-eigenvector $\eta = \sum_{s\in I_{\frac{1}{2}}} d_s \boldsymbol{w}_s^{\frac{1}{2}}$. By the self-conjugacy we get $d_{-\frac{1}{2}} = (-q)^{\frac{1}{2}}\overline{d_{\frac{1}{2}}}$. We may assume $d_{\frac{1}{2}}$ is real number by applying β_z^L for some $z \in \mathbb{T}$. Uniqueness up to conjugation follows from this observation. Hence we conclude $\eta = (\sqrt{q_0}x+v, \sqrt{q_0}u+y)$ is a desired $\pi_{\frac{1}{2}}$-vector. Let us write $a = \sqrt{q_0}x+v$ and $b = \sqrt{q_0}u + y$. We study the C^*-algebra A, which is generated by a and b, and derive the result that it is really of type D_1. By direct calculation we obtain

$$a^* = \sqrt{q_0}\, b\,, \quad \sqrt{q_0}\, ab + \sqrt{q_0}^{\,-1} ba = 1 + q_0\,.$$

Because of this equality the smooth part of A is linearly spanned by $a^k b^\ell$ for $k, \ell \geq 0$. We shall show $P_{\frac{1}{2}}(a^k b^\ell) \in \mathbb{C}a + \mathbb{C}b$. If it is done, then it shows the $\pi_{\frac{1}{2}}$-multiplicity is exactly one. Since other types which have a single loop at a vertex do not occur, we can conclude A is of type D_1. Now we start a proof. Applying β^R to a and b, we get $\beta_z^R(a) = za$ and $\beta_z^L(b) = \bar{z}b$ for $z \in \mathbb{T}$. Hence the element $P_{\frac{1}{2}}(a^k b^\ell)$ must have the following form.

$$\begin{cases} P_{\frac{1}{2}}(a^{\ell+1}b^{\ell}) = \lambda a\,, \\ P_{\frac{1}{2}}(a^{k}b^{k+1}) = \mu b\,, \\ \quad P_{\frac{1}{2}}(a^{k}b^{\ell}) = \quad 0 \ \text{ if } |k-\ell| \neq 1\,, \end{cases}$$

where λ and μ are complex number. If we show $a^k b^{k+1} = \mu b + \cdots$, then we have $a^{k+1}b^k = \bar{\mu}a + \cdots$. Hence it suffices to show $P_{\frac{1}{2}}(a^k b^{k+1}) = \mu b$. Recall the following well-known binomial equality.

$$(c+d)^n = \sum_{r=0}^{n} \begin{bmatrix} n \\ r \end{bmatrix}_q c^r d^{n-r} \quad \text{if} \quad dc = qcd\,.$$

Then we have

$$a^k b^{k+1} = \sum_{\substack{0\leq r\leq k \\ 0\leq s\leq k+1}} \begin{bmatrix} k \\ r \end{bmatrix}_q \begin{bmatrix} k+1 \\ s \end{bmatrix}_q \sqrt{q_0}^{\,s+r} x^r v^{k-r} u^s y^{k+1-s}\,.$$

We want to take out the coefficient of u and y. For the sake of this, we make use of β^L. Since we have $\beta^L_z(x^r v^{k-r} u^s y^{k+1-s}) = z^{2r+2s-2k-1} x^r v^{k-r} u^s y^{k+1-s}$, we obtain

$$\begin{aligned}
P_{\frac{1}{2}}(a^k b^{k+1}) &= \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q \begin{bmatrix} k+1 \\ k+1-r \end{bmatrix}_q \sqrt{q_0}^{k+1} P_{\frac{1}{2}}(x^r v^{k-r} u^{k+1-r} y^r) \\
&\quad + \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q \begin{bmatrix} k+1 \\ k-r \end{bmatrix}_q \sqrt{q_0}^{k} P_{\frac{1}{2}}(x^r v^{k-r} u^{k-r} y^{r+1}) \\
&= \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q \begin{bmatrix} k+1 \\ r \end{bmatrix}_q \sqrt{q_0}^{k+1} q^{-r(-2r+2k+1)} P_{\frac{1}{2}}(x^r y^r u^{k-r} v^{k-r} u) \\
&\quad + \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q \begin{bmatrix} k+1 \\ k-r \end{bmatrix}_q \sqrt{q_0}^{k} q^{-r(-2r+2k)} P_{\frac{1}{2}}(x^r y^r u^{k-r} v^{k-r} y) \\
&= \frac{\sqrt{q_0}^{k+1}(2)_q}{(k+2)_q} \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q \begin{bmatrix} k+1 \\ r \end{bmatrix}_q (-1)^{r(k-r)} q^{r(r-k+1)} \frac{(r)_q!(k-r+1)_q!}{(k+1)_q!}\, u \qquad (7.2) \\
&\quad + \frac{\sqrt{q_0}^{k} q_0^{-k}(2)_q}{(k+2)_q} \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q \begin{bmatrix} k+1 \\ k-r \end{bmatrix}_q (-1)^{r(k-r)+k} q^{r(r-k+2)} \frac{(r+1)_q!(k-r)_q!}{(k+1)_q!} y,
\end{aligned}$$

where we have used Lemma 7.11 in the last equality. Moreover we use the following formula.

$$\frac{(n)_q!}{(r)_q!(n-r)_q!} = (-1)^{r(n-r)} \frac{p_n}{p_r p_{n-r}} \begin{bmatrix} n \\ r \end{bmatrix}_q .$$

Then (7.2) is equal to

$$\begin{aligned}
(7.2) &= \frac{(-1)^k \sqrt{q_0}^{k+1}(2)_q}{(k+2)_q p_{k+1}} \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^{k-r} q^{r(r-k+1)} p_r p_{k-r+1}\, u \\
&\quad + \frac{(-1)^k \sqrt{q_0}^{k} q_0^{-k}(2)_q}{(k+2)_q! p_{k+1}} \sum_{r=0}^{k} \begin{bmatrix} k \\ r \end{bmatrix}_q (-1)^{k-r} q^{r(r-k+2)} p_{r+1} p_{k-r}\, y \\
&= \frac{(-1)^k \sqrt{q_0}^{k}(2)_q}{(k+2)_q p_{k+1}} (1+q^{-1})(q^{-2};q^{-1})_k (\sqrt{q_0}u + y)\,,
\end{aligned}$$

where we have used Lemma 7.12. Hence we have obtained $P_{\frac{1}{2}}(a^k b^{k+1}) \in \mathbb{C}(\sqrt{q_0}u + y)$ and this completes the proof of the existence of a right coideal of type D_1.

(IV) $A'_m\,(3 \leq m \leq \infty)$ case. We assume that there exists a right coideal of type A'_m. Although we only treat the finite m case here, we can similarly derive a contradiction in the case of $m = \infty$. Look at the first vertex from the left of Figure 14. Since the entry of the Perron-Frobenius eigenvector of this vertex is 1, the corresponding ergodic system becomes a right coideal of $C(SU_q(2))$ by Lemma 4.22. Let us denote the right coideal by A. It is also of type A'_m by Theorem 4.18. The spectral subspace $A_{\pi_{\frac{1}{2}}}$ generates a right coideal of type D_1, which is denoted by B. Now we consider the subsystem $B \subset A$. Let Λ be the inclusion matrix of $B \rtimes_\delta SU_q(2) \subset A \rtimes_\delta SU_q(2)$. Apply Proposition 4.20 and we get the equality $\Lambda\, \mathbb{M}^B(\pi_{\frac{1}{2}}) = \mathbb{M}^A(\pi_{\frac{1}{2}})\, \Lambda$, where $\mathbb{M}^B$ and $\mathbb{M}^A$ are the multiplicity maps of B and A, respectively. They have the following form.

$$\mathbb{M}^B(\pi_{\frac{1}{2}}) = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \quad \mathbb{M}^A(\pi_{\frac{1}{2}}) = \begin{pmatrix} 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 1 & \ddots & \ddots & 0 & 0 \\ 0 & 1 & 0 & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \ddots & \ddots & \ddots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & 1 & 0 \\ 0 & 0 & \dots & 0 & 1 & 0 & 1 \\ 0 & 0 & \dots & 0 & 0 & 1 & 1 \end{pmatrix}.$$

We know that the minimal projection $p_0 \in C_r^*(SU_q(2))$ is also minimal in $B \rtimes_\delta SU_q(2)$ and $A \rtimes_\delta SU_q(2)$. We discuss the corresponding vertex of p_0. About $B \rtimes_\delta SU_q(2)$ we may assume p_0 sits at the left vertex of Figure 12. About $A \rtimes_\delta SU_q(2)$ we may assume p_0 sits at the first vertex from the left in Figure 14 because the reduced system $p_0(A \otimes \mathbb{K}(L^2(SU_q(2))))p_0$ is canonically isomorphic to A. Hence the inclusion matrix Λ must be as the following form.

$$\Lambda = \begin{pmatrix} 1 & \lambda_1 \\ 0 & \lambda_2 \\ \vdots & \vdots \\ 0 & \lambda_{m-1} \\ 0 & \lambda_m \end{pmatrix}.$$

Then we want to solve the equation $\Lambda\, \mathbb{M}^B(\pi_{\frac{1}{2}}) = \mathbb{M}^A(\pi_{\frac{1}{2}})\, \Lambda$ about the non-negative integers $\lambda_1 \dots \lambda_m$, however, we immediately see it has no solutions. This is a contradiction and hence there is not a right coideal of type A'_m.

8. Classification of right coideals of $C(SU_{-1}(2))$

In this final section we complete the classification program of right coideals associated to the quantum $SU(2)$ group. Its representation theory such as actions of $U_{-1}(su_2)$ on $C(SU_{-1}(2))$ or the Clebsh-Gordan coefficients is obtained by the limit of $q_0 \to 1$. The continuous function algebra $C(SU_{-1}(2))$ is generated by x, u, v and y which satisfy

$$ux = -xu\,, \quad vx = -xv\,, \quad uy = -yu\,, \quad vy = -yv\,, \quad uv = vu\,,$$

$$xy + uv = yx + uv = 1\,, \quad x^* = y\,, \quad u^* = v\,.$$

By simple calculation, we see $\kappa^2 = \mathrm{id}$. Hence $C(SU_{-1}(2))$ is a compact Kac algebra. Refer the theory of general Kac algebras to [ES]. One of differences between cases of $q = -1$ and $q^2 \neq 1$ is amount of the quantum subgroups (or right coideals). In fact as we have seen there are not a lot of right coideals in $C(SU_q(2))$, on the contrary, in [Po2] he completely collects plentiful quantum subgroups of $SU_{-1}(2)$. The main result in this section is as follows. The definition of $\eta^{\frac{n}{2}}$ and $\widehat{\eta}^{\frac{n}{2}}$ is given in the case of type $\mathbb{T}_n$ with odd $n \geq 3$.

Theorem 8.1. *If a right coideal A is not of type $\mathbb{T}_n$ (odd $n \geq 3$) or D_n (odd $n \geq 1$), there exists a closed subgroup H in $SO_1(3)$ such that A is $C(H \backslash SO_{-1}(3))$. If a right coideal A is of type $\mathbb{T}_n$ (odd $n \geq 3$), A is conjugated to $C(\mathbb{T}_n \setminus SU_{-1}(2))$ or $C^*(\eta^{\frac{n}{2}}, \widehat{\eta}^{\frac{n}{2}})$. If a right coideal A is of type D_1, then A is conjugated to $C(D_1 \setminus SU_{-1}(2))$. If a right coideal A is of type D_n (odd $n \geq 3$), A is conjugated to $C(D_n \setminus SU_{-1}(2))$ or $C^*(\eta^{\frac{n}{2}})$. Here conjugation is given by β_z^L for some $z \in \mathbb{T}$.*

Although $C(\mathbb{T}_n \setminus SU_{-1}(2))$ is not isomorphic to $C^*(\eta^{\frac{n}{2}}, \widehat{\eta}^{\frac{n}{2}})$ as a $SU_{-1}(2)$-covariant system, in Proposition 8.11 we show that $C(D_n \backslash SU_{-1}(2))$ is isomorphic to $C^*(\eta^{\frac{n}{2}})$ as a $SU_{-1}(2)$-covariant system. As we have said in the previous section, we have the isomorphism between $C(SO_{-1}(3))$ and $C(SO_1(3))$ as compact quantum groups. Hence we can conclude that right coideals are quotient by subgroups when they are one of type A_4^*, S_4^*, A_5^*, D_m^* ($m \geq 2$), D_∞^*, $\mathbb{T}_{2n}$ ($n \geq 1$). Hence we have to study other types: $\mathbb{T}_n$ (odd $n \geq 3$), A'_m ($m \geq 3$), D_n (odd $n \geq 1$).

(I) A'_m ($m \geq 3$) case. The entirely same proof as in the previous section derives a contradiction well.

(II) D_1 case. If we consider the limit $q_0 \to 1$ in the previous section, we can show its existence. In that procedure we do not need any lemma stated in that section, because $\pi_{\frac{1}{2}}$-eigenvector is uniquely determined and in [Po2, Proposition 3.8] it has been shown that there exists a right coideal by subgroup D_1 whose $\pi_{\frac{1}{2}}$-multiplicity is one.

(III) $\mathbb{T}_n$ (odd $n \geq 3$) case. A has the spectral pattern

$$\bigoplus_{k \in \mathbb{Z}_{\geq 0}} \left(1 + 2\left[\frac{k}{n}\right]\right)\pi_k \oplus \bigoplus_{k \in \mathbb{Z}_{\geq 0}} 2\left[\frac{2k+n+1}{2n}\right]\pi_{k+\frac{1}{2}}.$$

Since its integer part becomes a right coideal in $C(SO_{-1}(3))$, it must be quotient by a subgroup $H \subset SO_{-1}(3)$. From the spectral pattern there exist angles $0 \leq \chi < 2\pi$ and $0 \leq \psi < \pi$ such that $H = \mathbb{T}_n^{\chi,\psi}$. We begin to determine a π_1-eigenvector of $C(\mathbb{T}_n^{\chi,\psi} \setminus SO_{-1}(3))$.

Lemma 8.2. *The following vector is a π_1-eigenvector for $C(\mathbb{T}_n^{\chi,\psi} \setminus SO_{-1}(3))$.*

$$\xi^{\chi,\psi} = \sqrt{2}^{-1}(-ie^{i\chi}\sin\psi, \sqrt{2}\cos\psi, ie^{-i\chi}\sin\psi)w(\pi_1).$$

Proof. First we consider a π_1-eigenvector $\zeta^{\chi,\psi} = \sum_{t \in I_1} c_t \boldsymbol{w}_t^1$ of $C(\mathbb{T}_n^{\chi,\psi} \setminus SO(3))$. The left action of $\pi_1(g) \in \mathbb{T}_n^{\chi,\psi}$ is given by multiplication of $w(\pi_1)(g)$ to $w(\pi_1)$ from the left. Hence we have to get a vector $(c_{-1}, c_0, c_1)w(\pi_1)(g) = (c_{-1}, c_0, c_1)$ for all $g \in SU(2)$ with $\pi_1(g) \in \mathbb{T}_n^{\chi,\psi}$. If χ and ψ are equal to 0, we can easily get

$c_{\pm 1} = 0$ and $c_0 = 1$. Since we know $\mathbb{T}_n^{\chi,\psi} = \mathrm{Ad}(\pi_1(r^{12}(\chi)r^{13}(\psi)))(\mathbb{T}_n^{0,0})$, we obtain $(c_{-1}, c_0, c_1) = (0,1,0)w(\pi_1)(r^{13}(-\psi)r^{12}(-\chi))$. Using two matrices

$$w(\pi_1)(r^{13}(-\psi)) = \begin{pmatrix} \cos^2\frac{\psi}{2} & -\sqrt{2}\sin\frac{\psi}{2}\cos\frac{\psi}{2} & \sin^2\frac{\psi}{2} \\ \sqrt{2}\sin\frac{\psi}{2}\cos\frac{\psi}{2} & 1-2\sin^2\frac{\psi}{2} & -\sqrt{2}\sin\frac{\psi}{2}\cos\frac{\psi}{2} \\ \sin^2\psi & -\sqrt{2}\sin\frac{\psi}{2}\cos\frac{\psi}{2} & \cos^2\frac{\psi}{2} \end{pmatrix}$$

and

$$w(\pi_1)(r^{12}(-\chi)) = \begin{pmatrix} e^{i\chi} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-i\chi} \end{pmatrix},$$

we have $c_{\pm 1} = \mp\sqrt{2}^{-1}e^{\mp i\chi}\sin\psi$ and $c_0 = \cos\psi$. Next we make use of $\Xi_1 : C(SO_1(3)) \longrightarrow C(SO_{-1}(3))$ and get the desired eigenvector. □

Next we determine the $\pi_{\frac{n}{2}}$-eigenvector space which is two-dimensional. Let us write m for $\frac{n}{2}$. For $\eta = \sum_{t\in I_m} d_t \boldsymbol{w}_t^m$ we use $q=-1$ version of Lemma 7.2 and we obtain

$$\begin{aligned}\sqrt{2m(2m-1)}\Psi_{m-1}(\xi^{\chi,\psi},\eta) = \sum_{t\in I_m}\bigl\{ &- ie^{i\chi}\sin\psi\sqrt{(m+t)(m+t+1)}d_{t+1} \\ &- (-1)^{m-t}2\cos\psi\sqrt{(m-t)(m+t)}d_t \\ &+ ie^{-i\chi}\sin\psi\sqrt{(m-t)(m-t+1)}d_{t-1}\bigr\}\boldsymbol{w}_t^m .\end{aligned}$$

Define α_t, β_t and γ_t by

$$\begin{aligned}\alpha_t &= -ie^{i\chi}\sin\psi\sqrt{(m+t)(m+t+1)}\,, \\ \beta_t &= -(-1)^{m-t}2\cos\psi\sqrt{(m-t)(m+t)}\,, \\ \gamma_t &= ie^{-i\chi}\sin\psi\sqrt{(m-t)(m-t+1)}\,.\end{aligned}$$

Then we have $\alpha_{-t} = \overline{\gamma_t}$ and $\beta_{-t} = -\beta_t$. If η is a π_m-eigenvector for A, then we have the following recurrence formula for $t \in I_m$.

$$\alpha_t d_{t+1} + \beta_t d_t + \gamma_t d_{t-1} = 0\,. \tag{8.1}$$

If we assume η is self-conjugate, we have

$$\alpha_t d_{t+1} - \beta_t d_t + \gamma_t d_{t-1} = 0$$

where we define $d_t = 0$ for $|t| \geq m+1$. From (8.1) we get $\beta_t d_t = 0$ and $\alpha_t d_{t+1} + \gamma_t d_{t-1} = 0$.

(a) $\psi \neq 0, \frac{\pi}{2}$ case. Since β_t is equal to zero if $|t| = m$, we have $d_t = 0$ for $|t| \leq m-1$. We also have $\alpha_{m-1}d_m + \gamma_{m-1}d_{m-2} = 0$. Then we get $d_m = 0$ by $\alpha_{m-1} \neq 0$ and $m-2 \geq -m+1$. This shows d_t are all equal to zero. This is not appropriate.

(b) $\psi = 0$ case. Then similarly we have $d_t = 0$ for $|t| \leq m-1$. Since π_m-eigenvector space is two dimensional, it is spanned by $\boldsymbol{w}_{\pm m}^m$. This shows $A = C(\mathbb{T}_n \setminus SU_{-1}(2))$.

(c) $\psi = \frac{\pi}{2}$ case. We have only a non-trivial equation $\alpha_t d_{t+1} + \gamma_t d_{t-1} = 0$. Its solution space is two dimensional. We give an explicit solution as follows. The proof is straightforward.

Lemma 8.3. *Let m be a half integer in $\frac{3}{2} + \mathbb{Z}_{\geq 0}$ and $0 \leq \chi < 2\pi$. Consider the recurrence equation,*

$$-ie^{i\chi}\sqrt{(m+t)(m+t+1)}d_{t+1} + ie^{-i\chi}\sqrt{(m-t)(m-t+1)}d_{t-1} = 0.$$

Then its solution is a linear combination of $d_t = e^{i(m-t)\chi}\left[\begin{smallmatrix} 2m \\ m-t \end{smallmatrix}\right]^{\frac{1}{2}}$ and $d_t = (-1)^{m-t}e^{i(m-t)\chi}\left[\begin{smallmatrix} 2m \\ m-t \end{smallmatrix}\right]^{\frac{1}{2}}$.

Define two π_m-eigenvectors of $C(SU_{-1}(2))$ η^m_χ and $\widehat{\eta}^m_\chi$ by

$$\eta^{m,\chi} = \sum_{t \in I_m} e^{i(m-t)\chi}\begin{bmatrix} 2m \\ m-t \end{bmatrix}^{\frac{1}{2}} \boldsymbol{w}^m_t\,,$$

$$\widehat{\eta}^{m,\chi} = \sum_{t \in I_m} (-1)^{m-t}e^{i(m-t)\chi}\begin{bmatrix} 2m \\ m-t \end{bmatrix}^{\frac{1}{2}} \boldsymbol{w}^m_t\,.$$

Since the π_m-eigenvector space of A is two-dimensional, it is spanned by the above vectors. Conversely we consider the right coideal B which is generated by $\eta^{m,\chi}$ and $\widehat{\eta}^{m,\chi}$. We want to conclude that B is a right coideal of type $\mathbb{T}_n$ and hence B coincides with A as a result. For the sake of this, it suffices to show that the π_ℓ-eigenvector space of B is trivial for all $\ell \in \{\frac{1}{2}, \ldots, m-1\}$. We prepare the element $g^{\theta,\chi}$ of $SU(2)$ defined by $g^{\theta,\chi} = \begin{pmatrix} \cos\frac{\theta}{2} & -ie^{-i\chi}\sin\frac{\theta}{2} \\ -ie^{i\chi}\sin\frac{\theta}{2} & \cos\frac{\theta}{2} \end{pmatrix}$. Recall the $*$-homomorphism $\upsilon_{g^{\theta,\chi}} : C(SU_{-1}(2)) \longrightarrow \mathbb{B}(\mathbb{C}^2)$

$$\begin{pmatrix} \upsilon_{g^{\theta,\chi}}(x) & \upsilon_{g^{\theta,\chi}}(u) \\ \upsilon_{g^{\theta,\chi}}(v) & \upsilon_{g^{\theta,\chi}}(y) \end{pmatrix} = \begin{pmatrix} \cos\frac{\theta}{2}\sigma_1 & ie^{-i\chi}\sin\frac{\theta}{2}\sigma_2 \\ -ie^{i\chi}\sin\frac{\theta}{2}\sigma_2 & \cos\frac{\theta}{2}\sigma_1 \end{pmatrix}.$$

Define a $*$-homomorphism $\rho_{g^{\theta,\chi}} = (\upsilon_{g^{\theta,\chi}} \otimes \mathrm{id}) \circ \delta$ and we have

$$\begin{pmatrix} \rho_{g^{\theta,\chi}}(x) \, \rho_{g^{\theta,\chi}}(u) \\ \rho_{g^{\theta,\chi}}(v) \, \rho_{g^{\theta,\chi}}(y) \end{pmatrix} = \begin{pmatrix} \cos\frac{\theta}{2}\sigma_1 \otimes x + ie^{-i\chi}\sin\frac{\theta}{2}\sigma_2 \otimes v & \cos\frac{\theta}{2}\sigma_1 \otimes u + ie^{-i\chi}\sin\frac{\theta}{2}\sigma_2 \otimes y \\ -ie^{i\chi}\sin\frac{\theta}{2}\sigma_2 \otimes x + \cos\frac{\theta}{2}\sigma_1 \otimes v & -ie^{i\chi}\sin\frac{\theta}{2}\sigma_2 \otimes u + \cos\frac{\theta}{2}\sigma_1 \otimes y \end{pmatrix}.$$

Lemma 8.4. *Let m be a half integer in $\frac{1}{2} + \mathbb{Z}_{\geq 0}$. For all $s \in I_m$ we have the following equalities.*

$$\rho_{g^{\theta,\chi}}((\eta^{m,\chi})_s) = \sum_{r \in I_m} (\cos m\theta\sigma_1 + (-1)^{m-r} i \sin m\theta\sigma_2) \otimes e^{i(m-r)\chi}\begin{bmatrix} 2m \\ m-r \end{bmatrix}^{\frac{1}{2}} w(\pi_m)_{r,s},$$

$$\rho_{g^{\theta,\chi}}((\widehat{\eta}^{m,\chi})_s) = \sum_{r \in I_m} ((-1)^{m-r}\cos m\theta\sigma_1 - i\sin m\theta\sigma_2) \otimes e^{i(m-r)\chi}\begin{bmatrix} 2m \\ m-r \end{bmatrix}^{\frac{1}{2}} w(\pi_m)_{r,s}.$$

Proof. Since $\rho_{g^{\theta,\chi}}$ commutes with the lowering operator f, it is enough to show those equalities with $s = -m$. Here it is proved by induction on m. We can easily show $\rho_{g^{\theta,\chi}} \circ \beta^L_z = (\mathrm{id} \otimes \beta^L_z) \circ \rho_{g_{\theta,0}}$ where z is equal to $e^{i\frac{\chi}{2}}$. Moreover we have $\eta^{m,\chi}_{-m} = e^{im\chi}\beta^L_z(\eta^{m,0}_{-m})$ and $\widehat{\eta}^{m,\chi}_{-m} = e^{im\chi}\beta^L_z(\widehat{\eta}^{m,0}_{-m})$. Hence it yields $\rho_{g^{\theta,\chi}}(\eta^{m,\chi}_{-m}) = e^{im\chi}(\mathrm{id} \otimes \beta^L_z) \circ \rho_{g_{\theta,0}}(\eta^{m,0}_{-m})$ and $\rho_{g^{\theta,\chi}}(\widehat{\eta}^{m,\chi}_{-m}) = e^{im\chi}(\mathrm{id} \otimes \beta^L_z) \circ \rho_{g_{\theta,0}}(\widehat{\eta}^{m,0}_{-m})$. Hence we may assume χ is equal to 0. For simplicity of notations we write g^θ, η^m and $\widehat{\eta}^m$ for $g_{\theta,0}$, $\eta^{m,0}$ and $\widehat{\eta}^{m,0}$, respectively. When m is equal to $\frac{1}{2}$, the desired equalities are easily obtained. We assume that the desired equalities hold for m. We can justify an equality about $\rho_{g^\theta}(\eta^{m+1}_{-(m+1)})$ as follows. Making use of the equality $\eta^{m+1}_{-(m+1)} = x^2\eta^m_{-m} + 2x\eta^m_{-m}v + \eta^m_{-m}v^2$, the computation of $\rho_{g^\theta}(x^2\eta^m_{-m})$, $\rho_{g^\theta}(2x\eta^m_{-m}v)$ and $\rho_{g^\theta}(\eta^m_{-m}v^2)$ is carried out in the following way.

$$\begin{aligned}
\rho_{g^\theta}(x^2\eta^m_{-m}) &= (\cos^2\frac{\theta}{2} \otimes x^2 - 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}\sigma_3 \otimes xv - \sin^2\frac{\theta}{2} \otimes v^2) \\
&\quad \cdot \sum_{r \in I_m} (\cos m\theta\sigma_1 + (-1)^{m-r} i \sin m\theta\sigma_2) \otimes \begin{bmatrix} 2m \\ m-r \end{bmatrix} x^{m-r}v^{m+r} \\
&= \sum_{r \in I_{m+1}} \Bigg(\cos^2\frac{\theta}{2}(\cos m\theta\sigma_1 + (-1)^{m+1-r} i \sin m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r-1 \end{bmatrix} \\
&\qquad + 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}(-\sin m\theta\sigma_1 + (-1)^{m+1-r} i \cos m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r \end{bmatrix} \\
&\qquad - \sin^2\frac{\theta}{2}(\cos m\theta\sigma_1 + (-1)^{m+1-r} i \sin m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r+1 \end{bmatrix} \Bigg) \\
&\qquad \otimes x^{m+1-r}v^{m+1+r},
\end{aligned}$$

$$\begin{aligned}
\rho_{g^\theta}(x\eta^m_{-m}v) = \sum_{r \in I_{m+1}} \Bigg(& \sin\frac{\theta}{2}\cos\frac{\theta}{2}(-\sin m\theta\sigma_1 + (-1)^{m+r} i \cos m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r-1 \end{bmatrix} \\
& + (\cos^2\frac{\theta}{2} - \sin^2\frac{\theta}{2})(\cos m\theta\sigma_1 - (-1)^{m-r} i \sin m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r \end{bmatrix} \\
& + \sin\frac{\theta}{2}\cos\frac{\theta}{2}(-\sin m\theta\sigma_1 - (-1)^{m-r} i \cos m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r+1 \end{bmatrix} \Bigg) \\
& \otimes x^{m+1-r}v^{m+1+r}
\end{aligned}$$

and

$$\begin{aligned}
\rho_{g^\theta}(\eta^m_{-m}v^2) = \sum_{r \in I_{m+1}} \Bigg(& -\sin^2\frac{\theta}{2}(\cos m\theta\sigma_1 - (-1)^{m-r} i \sin m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r-1 \end{bmatrix} \\
& + 2\sin\frac{\theta}{2}\cos\frac{\theta}{2}(-\sin m\theta\sigma_1 + (-1)^{m+r} i \cos m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r \end{bmatrix} \\
& + \cos^2\frac{\theta}{2}(\cos m\theta\sigma_1 - (-1)^{m-r} i \sin m\theta\sigma_2) \begin{bmatrix} 2m \\ m-r+1 \end{bmatrix} \Bigg) \\
& \otimes x^{m+1-r}v^{m+1+r}.
\end{aligned}$$

Then we can derive $\rho_{g_\theta}(\eta^{m+1}_{-(m+1)}) = \sum_{r\in I_{m+1}}(\cos(m+1)\theta + (-1)^{m+1-r} i \sin(m+1)\theta) \otimes \left[{2m+2 \atop m+1-r}\right] x^{m+1-r} v^{m+1+r}$ where we use the formula $\left[{2m+2 \atop m+1-r}\right] = \left[{2m \atop m-r-1}\right] + 2\left[{2m \atop m-r}\right] + \left[{2m \atop m-r+1}\right]$. By induction the assertion about $\rho_{g^\theta}(\eta^m_{-m})$ is justified. About $\rho_{g^\theta}(\widehat{\eta}^m_{-m})$ we make use of $\rho_{g^\theta} \circ \beta^L_i = (\mathrm{id} \otimes \beta^L_i) \circ \rho_{g^{-\theta}}$ and $\widehat{\eta}^m_{-m} = i^{2m}\beta^L_i(\eta^m_{-m})$. $\square$

Especially putting $\theta = \frac{\pi}{m}k$ $(k = 0, \dots, n-1)$ and $z = e^{-i\frac{\chi}{2}}$, we get

$$\begin{aligned} \rho_{g_{\frac{\pi}{m}k,\chi}}(\eta^{m,\chi}_s) &= (-1)^k \sigma_1 \otimes \eta^{m,\chi}_s, \\ \rho_{g_{\frac{\pi}{m}k,\chi}}(\widehat{\eta}^{m,\chi}_s) &= (-1)^k \sigma_1 \otimes \widehat{\eta}^{m,\chi}_s\,. \end{aligned}$$

Therefore B is contained in a C^*-algebra $C^{n,\chi} = \{a \in C(SU_{-1}(2)) \mid \rho_{g_{\frac{\pi}{m}k,\chi}}(a) \in (\mathbb{C} + \mathbb{C}\sigma_1) \otimes \mathbb{C}a \text{ for all } 0 \le k < n\}$. Actually, $C^{n,\chi}$ is a right coideal because of $(\rho_{g_{\frac{\pi}{m}k,\chi}} \otimes \mathrm{id}) \circ \delta = (\mathrm{id} \otimes \delta) \circ \rho_{g_{\frac{\pi}{m}k,\chi}}$.

Lemma 8.5. *The following equalities hold.*

$$\begin{aligned} \Psi_1(\eta^{m,\chi}, \widehat{\eta}^{m,\chi}) &= -\, i\sqrt{m}2^{2m}e^{2im\chi}\xi^{\chi,\frac{\pi}{2}}, \\ \Psi_1(\widehat{\eta}^{m,\chi}, \eta^{m,\chi}) &= -\, i\sqrt{m}2^{2m}e^{2im\chi}\xi^{\chi,\frac{\pi}{2}}, \\ \Psi_1(\eta^{m,\chi}, \eta^{m,\chi}) &= \Psi_1(\widehat{\eta}^{m,\chi}, \widehat{\eta}^{m,\chi}) = 0. \end{aligned}$$

Proof. We give a proof for only the first one. Others are proved similarly. By definition we have $\Psi_1(\eta^{m,\chi}, \widehat{\eta}^{m,\chi}) = \sum_{r=0}^{2m-1}(C^1_{m,m})_r \eta^{m,\chi}_{m-r-1}\widehat{\eta}^{m,\chi}_{-m+r}$ where $(C^1_{m,m})_r$ is equal to $(-1)^r\sqrt{2m}^{-1}\sqrt{(r+1)(2m-r)}$. Define complex numbers $c_{\pm 1}$ and c_0 by $\Psi_1(\eta^{m,\chi}, \widehat{\eta}^{m,\chi}) = c_{-1}\boldsymbol{w}^1_{-1} + c_0\boldsymbol{w}^1_0 + c_1\boldsymbol{w}^1_1$. In order to obtain their values we use the restriction homomorphism $\pi_\mathbb{T} : C(SU_{-1}(2)) \longrightarrow C(\mathbb{T})$. In fact we get $\pi_\mathbb{T}(\Psi_1(\eta^{m,\chi}, \widehat{\eta}^{m,\chi})_s) = z^{2s}c_s$ for all $z \in \mathbb{T}$ and $1 \le s \le 1$. First we have

(8.2)
$$\Psi_1(\eta^{m,\chi}, \widehat{\eta}^{m,\chi})_{-1} = \sum_{r=0}^{2m-1} \sqrt{2m}^{-1}(-1)^r\sqrt{(r+1)(2m-r)}(\eta^{m,\chi})_{m-1-r}\widehat{\eta}^{m,\chi}_{-m+r}.$$

Applying $\pi_\mathbb{T}$ to the both side, we have

$$\begin{aligned} c_{-1} = &\sum_{r=0}^{2m-1} \sqrt{2m}^{-1}(-1)^r\sqrt{(r+1)(2m-r)}\begin{bmatrix} 2m \\ r+1 \end{bmatrix}^{\frac{1}{2}}\begin{bmatrix} 2m \\ 2m-r \end{bmatrix}^{\frac{1}{2}} \\ &\quad \cdot (-1)^{2m-r}e^{i(r+1)\chi}e^{i(2m-r)\chi} \\ = &(-1)^{2m}\sqrt{2m}^{-1}e^{i(2m+1)\chi}\sum_{r=0}^{2m-1}\frac{2m!}{r!(2m-1-r)!} \\ = &-2^{2m-1}\sqrt{2m}e^{i(2m+1)\chi}. \end{aligned}$$

If we apply the lowering operator f to (8.2), then we can easily obtain

$$\Psi_1(\eta^{m,\chi}, \widehat{\eta}^{m,\chi})_0 = -\sqrt{m}^{-1}\sum_{r=0}^{2m}(m-r)(\eta^{m,\chi})_{m-r}\widehat{\eta}^{m,\chi}_{-m+r}, \tag{8.3}$$

where we use the fact of $f \cdot \xi_p = i^{-2\nu+1}\sqrt{(\nu-p)(\nu+p+1)}\xi_{p+1}$ for a π_ν-eigenvector ξ. Applying $\pi_{\mathbb{T}}$ to (8.3), we obtain $c_0 = -\sqrt{m}^{-1}\sum_{r=0}^{2m}(m-r)\left[\begin{smallmatrix}2m\\ r\end{smallmatrix}\right](-1)^{2m-r} = 0$. Since we also have

$$\Psi_1(\eta^{m,\chi}, \widehat{\eta}^{m,\chi})_1 = \sum_{r=0}^{2m-1} \sqrt{2m}^{-1}(-1)^r\sqrt{(r+1)(2m-r)}(\eta^{m,\chi})_{m-r}\widehat{\eta}^{m,\chi}_{-m+r+1},$$

similarly we can derive $c_1 = 2^{2m-1}\sqrt{2m}e^{i(2m-1)\chi}$. □

Now we start to prove that B is a right coideal of type $\mathbb{T}_n$. Let n' be a smallest odd integer such that the $\pi_{\frac{n'}{2}}$- eigenvector space of B is not trivial. Put $m' = \frac{n'}{2}$. We claim $m' \geq \frac{3}{2}$. This is because there exist no non-zero complex numbers $c_{-\frac{1}{2}}$ and $c_{\frac{1}{2}}$ such that $c_{-\frac{1}{2}}x + c_{\frac{1}{2}}v$ is an element of $C^{m,\chi}$. From the previous lemma, B contains π_1-part and it shows B is not of type D_ℓ for some odd $\ell \geq 3$ but of $\mathbb{T}_{n'}$. Again by Lemma 8.3 for m', the $\pi_{m'}$-eigenvector space is spanned by $\eta^{m',\chi}$ and $\widehat{\eta}^{m',\chi}$. In particular, $\eta^{m',\chi}_{-m'}$ is an element of $C^{n,\chi}$. Since we have for $0 \leq k \leq n-1$

$$\rho_{g_{\frac{2\pi}{n}k,\chi}}(\eta^{m',\chi}_{-m'}) = \sum_{r\in I_{m'}}(\cos\frac{m'}{m}\pi k\,\sigma_1 - (-1)^{m'-r}i\sin\frac{m'}{m}\pi k\,\sigma_2)$$
$$\otimes e^{i(m'-r)\chi}\begin{bmatrix}2m'\\ m'-r\end{bmatrix}^{\frac{1}{2}} w(\pi_{m'})_{r,-m'}$$

by Lemma 8.4, $\frac{m'}{m}$ must be equal to 1. Hence the π_ℓ-eigenvector space is trivial for $\ell \in \{\frac{1}{2}, \ldots, m-1\}$. This case is only admitted as type $\mathbb{T}_n$. As a result we also see that B coincides with $C^{n,\chi}$.

We have proved that a right coideal of type $\mathbb{T}_n$ (odd $n \geq 3$) arises from the cyclic group in the maximal torus $\mathbb{T}$ or the cyclic group $\mathbb{T}_n^{\frac{\pi}{2},\chi}$. We have to check two right coideals $C(D_n \setminus SU_{-1}(2))$ and $C^{n,\chi}$ are not $SU_{-1}(2)$-isomorphic. It suffices to prove it in the case of $\chi = 0$ by considering β^L. Assume that there exists an $SU_{-1}(2)$-isomorphism $\vartheta : C(D_n \setminus SU_{-1}(2)) \longrightarrow C^{n,0}$. It induces the map between eigenvector spaces defined by $\vartheta(\xi) = (\vartheta(\xi_r))_{r\in I_\mu}$ for π_μ-eigenvector ξ. On the eigenvector spaces, ϑ preserves the inner product, the conjugation operation T and moreover the product map Ψ. Take complex numbers λ and μ which satisfy $\vartheta(\boldsymbol{w}^m_{-m}) = \lambda\eta^m + \mu\widehat{\eta}^m$. By Lemma 8.5 and $\Psi_1(\boldsymbol{w}^m_{-m}, \boldsymbol{w}^m_{-m}) = 0$, we have $-i\lambda\mu\sqrt{m}2^{2m+1}\xi^{0,\frac{\pi}{2}} = 0$. Hence $\lambda = 0$ or $\mu = 0$. By using β^L we may assume $\mu = 0$. Now we have $\vartheta(\boldsymbol{w}^m_{-m}) = \lambda\eta^m$. If we apply the conjugation T to the both side, we obtain $\vartheta(\boldsymbol{w}^m_m) = \overline{\lambda}\eta^m$. This shows the range of ϑ onto π_m-eigenvector space is one-dimensional, however, this is contradiction. Therefore $C(\mathbb{T}_n \setminus SU_{-1}(2))$ and $C^{n,0}$ are not $SU_{-1}(2)$-isomorphic. Finally we will make a discussion on characterizing a right coideal $C^{n,\chi}$ by a quantum subgroup of $SU_{-1}(2)$. Let us recall a closed subgroup of $SU_{-1}(2)$ which corresponds to $\mathbb{T}_n^{\frac{\pi}{2},0}$, that is, we consider the compact set $Z = \pi_1^{-1}(\mathbb{T}_n^{\frac{\pi}{2},0})$ and define $C(G_Z) = \pi_Z(C(SU_{-1}(2)))$. We denote a subgroup G_Z by $(\mathbb{T}_n^{\frac{\pi}{2},0})_\#$. The set Z

is actually a binary subgroup $(\mathbb{T}_n^{\frac{\pi}{2},0})^*$ and consists of $\{g^{\frac{2\pi}{n}k} \mid 0 \leq k \leq 2n-1\}$. Hence $C((\mathbb{T}_n^{\frac{\pi}{2},0})_\#) \subset \oplus_{0\leq k\leq 2n-1}\mathbb{M}_{s(k)}(\mathbb{C})$ where $s(k)=1$ if $k=0,\ n$ and $s(k)=2$ otherwise. We denote the coproduct of $C((\mathbb{T}_n^{\frac{\pi}{2},0})_\#)$ by δ_Z. Now we look at the equality $\rho_{g^{\frac{2\pi}{n}k}}(\eta_s^m) = (-1)^k\sigma_1 \otimes \eta_s^m$ for $0 \leq k \leq 2n-1$ and $s \in I_m$. Define a self-adjoint unitary in $C((\mathbb{T}_n^{\frac{\pi}{2},0})_\#)$ by $w = \oplus_{k\neq 0,n}(-1)^k\sigma_1 \oplus 1 \oplus -1$. Then the above equality is equivalent to $(\pi_Z \otimes \mathrm{id})\circ\delta(\eta_s^m) = w \otimes \eta_s^m$. Moreover, we have

$$\begin{aligned}
\delta_Z(w)\otimes\eta_s^m &= (\delta_Z\circ\pi_Z\otimes\mathrm{id})\circ\delta(\eta_s^m)\\
&= (\pi_Z\otimes\pi_Z\otimes\mathrm{id})\circ(\delta\otimes\mathrm{id})\delta(\eta_s^m)\\
&= (\pi_Z\otimes\pi_Z\otimes\mathrm{id})\circ(\mathrm{id}\otimes\delta)\delta(\eta_s^m)\\
&= w\otimes w\otimes\eta_s^m.
\end{aligned}$$

Hence we have proved the following proposition.

Proposition 8.6. *A self-adjoint unitary w is a group-like element in $C((\mathbb{T}_n^{\frac{\pi}{2},0})_\#)$.*

A C^*-algebra generated by w becomes a Hopf $*$-algebra which is isomorphic to $C(\mathbb{Z}_2)$. With this identification we have $C^{n,0} = \{a \in C(SU_{-1}(2)) \mid (\pi_Z\otimes\mathrm{id})\circ\delta(a) \in C(\mathbb{Z}_2)\otimes\mathbb{C}a\}$. For general $C^{n,\chi}$ we also get similar results by using β_z^L. We study all the irreducible representations of $(\mathbb{T}_n^{\frac{\pi}{2},0})_\#$ as follows. Let us denote the restriction of the fundamental representation to $(\mathbb{T}_n^{\frac{\pi}{2},0})_\#$ simply by $w(\pi_{\frac{1}{2}})|$. Then we have $w(\pi_{\frac{1}{2}})| = \left(\begin{smallmatrix}\pi_Z(x) & \pi_Z(u)\\ \pi_Z(v) & \pi_Z(y)\end{smallmatrix}\right)$ where we have

$$\begin{aligned}
\pi_Z(x) &= \oplus_{k\neq 0,n}\cos\frac{\pi}{n}k\,\sigma_1 \oplus 1 \oplus -1,\\
\pi_Z(u) &= \oplus_{k\neq 0,n}\, i\sin\frac{\pi}{n}k\,\sigma_2 \oplus 0 \oplus 0,\\
\pi_Z(v) &= \oplus_{k\neq 0,n} -i\sin\frac{\pi}{n}k\,\sigma_2 \oplus 0 \oplus 0,\\
\pi_Z(y) &= \oplus_{k\neq 0,n}\cos\frac{\pi}{n}k\,\sigma_1 \oplus 1 \oplus -1.
\end{aligned}$$

The unitary representation $w(\pi_{\frac{1}{2}})|$ invariantly acts on two lines $\mathbb{C}\xi_\pm = \mathbb{C}\left(\begin{smallmatrix}1\\ \pm i\end{smallmatrix}\right)$ and their one dimensional actions are given by w_+ and w_- with

$$w_\pm = \oplus_{k\neq 0,n}\begin{pmatrix}0 & e^{\pm i\frac{\pi}{n}k}\\ e^{\mp i\frac{\pi}{n}k} & 0\end{pmatrix}\oplus 1 \oplus -1.$$

Hence we have $w(\pi_{\frac{1}{2}})| = w_+ \oplus w_-$. Notice that $w_\pm$ are the self-conjugate unitary representations. This shows the difference from the fusion rules of the ordinary cyclic group $\mathbb{T}_{2n}$. And define the tensor product representations $w_{+-} = w_+w_-$ and $w_{-+} = w_-w_+$. Then they are self-conjugate and non-equivalent to each other, where non-equivalence comes from the non-commutativity of the underlying space $(\mathbb{T}_n^{\frac{\pi}{2},0})_\#$. If we want to write the McKay diagram, we compute $w(\pi_{\frac{1}{2}})|\cdot w_+ = 1 \oplus w_{-+}$ and bond the vertices $+$ and $-+$ by a single line. We



FIGURE 1. The McKay diagram of $(\mathbb{T}_n^{0,\frac{\pi}{2}})_\#$, $2n$ nodes

continue these procedures to get all the irreducible representations. As a result, they are represented by the reduced words ($+-+-$ etc.) whose lengths are less than or equal to n (Figure 1). For length n word, we can easily check $w = + - \cdots - + = - + \cdots + -$. Hence w sits at the bottom vertex. The fusion rules are given by this equality, $+^2 = 1$, $-^2 = 1$ and $\overline{(\pm)} = \pm$. In the above proposition, we get a self-adjoint unitary w. It is the n-th irreducible representation $+ - \cdots - +$. Let us denote $g = +-$ and $s = +$. Then we have $s^2 = 1$, $g^n = 1$ and $sgs = g^{-1}$. This shows that the dual group $\widehat{(\mathbb{T}_n^{\frac{\pi}{2},0})_\#}$ is isomorphic to D_n, and hence $C((\mathbb{T}_n^{\frac{\pi}{2},0})_\#) \cong C_r^*(D_n)$ as the Hopf-algebras.

(IV) D_n (odd $n \geq 3$) case. A has the spectral pattern

$$\bigoplus_{k\in\mathbb{Z}_{\geq 0}} \left(\left[\frac{k}{n}\right] + \frac{1+(-1)^k}{2}\right)\pi_k \oplus \bigoplus_{k\in\mathbb{Z}_{\geq 0}} \left(\left[\frac{2k+1}{n}\right] - \left[\frac{k}{n}\right]\right)\pi_{k+\frac{1}{2}}\,.$$

Let B be a right coideal of the integer parts of A. Since B is a right coideal of $C(SO_{-1}(3))$, it must be a quotient by subgroup of $SO_{-1}(3)$ for some angles $0 \leq \phi < \frac{2\pi}{n}$, $0 \leq \chi < 2\pi$ and $-\pi \leq \psi \leq \pi$. Look at the spectral pattern for B and we see that $B = C(D_n^{\phi,\chi,\psi} \setminus SO_{-1}(3))$ We have to clarify its self-conjugate π_2-eigenvector.

Lemma 8.7. *The following vector is a π_2-eigenvector for $C(D_n^{\phi,\chi,\psi} \setminus SU_{-1}(2))$.*

$$\zeta^{\phi,\chi,\psi} = \Big(e^{i2\chi}\frac{\sqrt{6}}{4}\sin^2\psi, ie^{i\chi}\frac{\sqrt{6}}{2}\sin\psi\cos\psi, 1 - \frac{6}{4}\sin^2\psi,$$
$$- ie^{-i\chi}\frac{\sqrt{6}}{2}\sin\psi\cos\psi, e^{-i2\chi}\frac{\sqrt{6}}{4}\sin^2\psi\Big)w(\pi_2)\,.$$

Proof. The proof is done as Lemma 8.2 with considering $w(\pi_2)$ and $D_{n,\phi,\chi,\psi} = \mathrm{Ad}(\pi_1(r^{12}(\chi)r^{13}(\psi)r^{12}(\phi)))(D_n^{0,0,0})$. □

Next we use the following lemma in order to get a $\pi_{\frac{n}{2}}$-eigenvector of A, which is proved by $q_0 \to 1$ in Lemma 7.5 and an elementary calculation for $\boldsymbol{w}_{\pm 1}^2$.

Lemma 8.8. *Let m be a half integer with $m \in \frac{1}{2} + \mathbb{Z}_{\geq 0}$. Then we have the following equalities for all $t \in I_m$.*

$$\Psi_{m-1}(\boldsymbol{w}^2_{-2}, \boldsymbol{w}^m_t) = (-1)^{m-t}\frac{1}{2m-2}\sqrt{\frac{4(m-t+1)(m+t)(m+t-1)(m+t-2)}{2m(2m-1)}}\,\boldsymbol{w}^{m-1}_{t-2},$$

$$\Psi_{m-1}(\boldsymbol{w}^2_{-1}, \boldsymbol{w}^m_t) = \frac{2(m-2t+2)}{2m-2}\sqrt{\frac{(m+t)(m+t-1)}{2m(2m-1)}}\boldsymbol{w}^{m-1}_{t-1},$$

$$\Psi_{m-1}(\boldsymbol{w}^2_{0}, \boldsymbol{w}^m_t) = -(-1)^{m-t}\frac{2t}{(2m-2)}\sqrt{\frac{3!(m+t)(m-t)}{2m(2m-1)}}\,\boldsymbol{w}^{m-1}_{t},$$

$$\Psi_{m-1}(\boldsymbol{w}^2_{1}, \boldsymbol{w}^m_t) = -\frac{2(m+2t+2)}{2m-2}\sqrt{\frac{(m-t)(m-t-1)}{2m(2m-1)}}\boldsymbol{w}^{m-1}_{t+1},$$

$$\Psi_{m-1}(\boldsymbol{w}^2_{2}, \boldsymbol{w}^m_t) = -(-1)^{m-t}\frac{1}{2m-2}\sqrt{\frac{4(m+t+1)(m-t)(m-t-1)(m-t-2)}{2m(2m-1)}}\,\boldsymbol{w}^{m-1}_{t+2}.$$

Let us define $\alpha_t, \beta_t, \gamma_t, \delta_t$ and ε_t for $t \in I_m$ by

$$\begin{aligned}
\alpha_t &= \frac{\sqrt{6}}{4}e^{i2\chi}\sin^2\psi\,(-1)^{m-t}\sqrt{4(m-t-1)(m+t+2)(m+t+1)(m+t)}\,,\\
\beta_t &= \sqrt{6}e^{i\chi}\sin\psi\cos\psi\,(m-2t)\sqrt{(m+t+1)(m+t)}\,,\\
\gamma_t &= 2(-1+\frac{6}{4}\sin^2\psi)(-1)^{m-t}t\sqrt{3!(m+t)(m-t)}\,,\\
\delta_t &= -\sqrt{6}e^{-i\chi}\sin\psi\cos\psi\,(m+2t)\sqrt{(m-t+1)(m-t)}\,,\\
\varepsilon_t &= -\frac{\sqrt{6}}{4}e^{-i2\chi}\sin^2\psi\,(-1)^{m-t}\sqrt{4(m+t-1)(m-t+2)(m-t+1)(m-t)}\,,
\end{aligned}$$

with $\alpha_m = 0 = \varepsilon_{-m}$. Now we put $m = \frac{n}{2}$. Take a self-conjugate π_m-eigenvector $\eta = \sum_{t\in I_m} d_t\boldsymbol{w}^m_t$ of A. The self-conjugacy means $d_{-t} = \overline{d_t}$. Since the π_{m-1}-part of A is zero, we obtain $\Psi_{m-1}(\zeta_\psi, \eta) = 0$ and this is equivalent to the following recurrence formula:

$$\alpha_t d_{t+2} + \beta_t d_{t+1} + \gamma_t d_t + \delta_t d_{t-1} + \varepsilon_t d_{t-2} = 0 \tag{8.4}$$

for $t \in I_m$, where we define $d_t = 0$ for $|t| \geq m+1$ as usual. Making use of $d_{-t} = \overline{d_t}$, $\alpha_{-t} = \overline{\varepsilon_t}$, $\beta_{-t} = -\overline{\delta_t}$ and $\gamma_{-t} = \gamma_t$, where we use $(-1)^{2t} = -1$, we also have

$$\alpha_t d_{t+2} - \beta_t d_{t+1} + \gamma_t d_t - \delta_t d_{t-1} + \varepsilon_t d_{t-2} = 0\,.$$

From (8.4) and this we get

$$\alpha_t d_{t+2} + \gamma_t d_t + \varepsilon_t d_{t-2} = 0\,, \tag{8.5}$$

$$\beta_t d_{t+1} + \delta_t d_{t-1} = 0\,. \tag{8.6}$$

We analyze them as follows.

(1) $0 < \psi < \frac{\pi}{2}$ case. We want to derive a contradiction to non-triviality of η. If we give a number d_{-m}, then by (8.5) or (8.6) we can inductively determine $d_{-m+2}, \cdots, d_{m-1}$. By the self-conjugacy, we obtain the whole numbers d_t. Hence η is uniquely determined by d_{-m} or d_m. Because of non-triviality of solutions $\{d_t\}_{t \in I_m}$, the determinant of the following matrix must be zero for all $t \in I_m$:

$$\begin{pmatrix} \alpha_t & \gamma_t & \varepsilon_t \\ \beta_{t+1} & \delta_{t+1} & 0 \\ 0 & \beta_{t-1} & \delta_{t-1} \end{pmatrix}.$$

Hence we have the equation for t, m and ψ

$$\alpha_t \delta_{t-1} \delta_{t+1} - \beta_{t+1} \gamma_t \delta_{t-1} + \beta_{t-1} \beta_{t+1} \varepsilon_t = 0 \tag{8.7}$$

for $t \in I_m$. If we put $t = m-1$, then we easily get $\sin^2 \psi = \frac{3m-4}{5m-8}$.

(1a) $n = 3$ case. Then m is equal to $\frac{3}{2}$ and we have $\sin^2 \psi = -1 < 0$. This is a contradiction.

(1b) $n \geq 5$ case. If we put $t = m-2 (\geq -m+1)$, then we easily get $\sin^2 \psi = \frac{12(m-2)^2}{5m^2-24m+24}$. This is contradiction to $\sin^2 \psi = \frac{3m-4}{5m-8}$.

Therefore ψ must be equal to 0 or $\frac{\pi}{2}$.

(2) $\psi = 0$ case. We have $\eta = \boldsymbol{w}_0^m$ and $\gamma_t = -2(-1)^{m-t} t \sqrt{3!(m+t)(m-t)}$ and $\alpha_t = \beta_t = \delta_t = \varepsilon_t = 0$. Then we get $d_t = 0$ for $|t| \leq m-1$. This case A is $C(D_n \setminus SU_{-1}(2))$.

(3) $\psi = \frac{\pi}{2}$ case. We have $\zeta^{\phi,\chi,\frac{\pi}{2}} = \frac{\sqrt{6}}{4} e^{i2\chi} \boldsymbol{w}_{-2}^2 - \frac{1}{2} \boldsymbol{w}_0^2 + \frac{\sqrt{6}}{4} e^{-i2\chi} \boldsymbol{w}_2^2$ and $\beta_t = \delta_t = 0$ and others are

$$\begin{aligned} \alpha_t =& \frac{\sqrt{6}}{4} e^{i2\chi} (-1)^{m-t} \sqrt{4(m-t-1)(m+t+2)(m+t+1)(m+t)}, \\ \gamma_t =& (-1)^{m-t} t \sqrt{3!(m+t)(m-t)}, \\ \varepsilon_t =& -\frac{\sqrt{6}}{4} e^{-i2\chi} (-1)^{m-t} \sqrt{4(m+t-1)(m-t+2)(m-t+1)(m-t)}. \end{aligned}$$

Recall vectors $\eta^{m,\chi}$ and $\widehat{\eta}^{m,\chi}$ which are introduced in the previous case $\mathbb{T}_n$. We easily confirm that they are independent solutions of $\alpha_t d_{t+2} + \gamma_t d_t + \varepsilon_t d_{t-2} = 0$. Let $\eta = \lambda_0 \eta^{m,\chi} + \lambda_1 \widehat{\eta}^{m,\chi}$ be a self-conjugate π_m-eigenvector of A where λ_0 and λ_1 are complex numbers. Since the conjugation operation T satisfies $T\eta^{m,\chi} = e^{-i2m\chi} \eta^{m,\chi}$ and $T\widehat{\eta}^{m,\chi} = -e^{-i2m\chi} \widehat{\eta}^{m,\chi}$, there exist real numbers μ_0 and μ_1 with $\lambda_0 = \mu_0 e^{-im\chi}$ and $\lambda_1 = i\mu_1 e^{-im\chi}$. The π_1-eigenvector $\Psi_1(\eta, \eta)$ must be zero because of the absence of π_1-part in A. By Lemma 8.5 we have $\Psi_1(\eta, \eta) = 2\mu_0 \mu_1 \sqrt{2m} 2^{2m-1} \xi^{\chi, \frac{\pi}{2}}$. It shows μ_0 or μ_1 is equal to zero. Hence the π_m-eigenvector space of B consists of scalar multiples of $\eta^{m,\chi}$ or $\widehat{\eta}^{m,\chi}$. We can easily check $\beta_i^L(\eta_r^{m,\chi}) = i^{-2m} \widehat{\eta}_r^{m,\chi}$ for all $r \in I_m$ and hence may assume that the π_m-eigenvector space of A is spanned by $\eta^{m,\chi}$ if necessary by applying β_i^L. Moreover we may also assume $\chi = 0$ by applying β_z^L where $z = e^{i\frac{\chi}{2}}$. Let B be a right coideal generated by $\eta_r^m = \eta_r^{m,0}$ with all $r \in I_m$. Notice that B is contained in $C^{n,0}$ which is defined in the discussion on the type $\mathbb{T}_n$. Hence B is of type $\mathbb{T}_n$ or D_n. We want to show B is actually of type D_n. This is proved by Proposition 8.11 or the following direct calculation which takes us the similar characterization on the right coideal as Proposition 8.6. In order to study it we have to clarify the type of the right coideal C which is the integer part of B. Whether C is of type $\mathbb{T}_n$ or D_n, its π_2-multiplicity is one. Hence the π_2-eigenvector of C is a scalar multiple of $\zeta^{\phi,0,\frac{\pi}{2}}$. We show that there exists an angle $0 \leq \phi < \frac{2\pi}{n}$ such that $C = C(D_n^{\phi,0,\frac{\pi}{2}} \setminus SO_{-1}(3))$. We already know all η_r^m for $r \in I_m$ are fixed by the rotation of $\mathbb{T}_n^{0,\frac{\pi}{2}}$. It suffices to show $\eta_r^m \eta_s^m$ is fixed by the rotation of angle π around the axis $\pi_1(r^{23}(-\phi)) \cdot \sigma_3$ for any $r, s \in I_m$. This rotation is obtained by the matrix $k^\phi \in SU(2)$ $k^\phi = \left(\begin{smallmatrix} -i\cos\phi & -\sin\phi \\ \sin\phi & i\cos\phi \end{smallmatrix}\right)$. Define the $*$-homomorphism $\rho_{k^\phi} = (\upsilon_{k^\phi} \otimes \mathrm{id}) \circ \delta$.

Lemma 8.9. *For any $m \in \frac{1}{2} + \mathbb{Z}_{\geq 0}$ and $s \in I_m$ the following equalities holds.*

$$\rho_{k^\phi}(\eta_s^m) = i^{2m} \sum_{r \in I_m} (\cos 2m\phi\sigma_1 - (-1)^{m-r} i \sin 2m\phi\sigma_2) \otimes (-1)^{m-r} \begin{bmatrix} 2m \\ m-r \end{bmatrix}^{\frac{1}{2}} w(\pi_m)_{r,s},$$

$$\rho_{k^\phi}(\widehat{\eta}_s^m) = i^{2m} \sum_{r \in I_m} (\cos 2m\phi\sigma_1 + (-1)^{m-r} i \sin 2m\phi\sigma_2) \otimes \begin{bmatrix} 2m \\ m-r \end{bmatrix}^{\frac{1}{2}} w(\pi_m)_{r,s}.$$

Proof. k^ϕ is equal to $r^{12}(\pi) r^{23}(2\phi) = r^{12}(\pi) g^{2\phi,0}$ where $g^{\theta,\chi}$ is a matrix defined in the study of $\mathbb{T}_n$ case. Since we have $\rho_{k^\phi}(x) = -i\rho_{g^{-2\phi,0}}(x)$ and $\rho_{k^\phi}(v) = i\rho_{g^{-2\phi,0}}(v)$, the equalities $\rho_{k^\phi}(\eta_{-m}^m) = i^{2m} \rho_{g^{2\phi,0}}(\widehat{\eta}_{-m}^m)$ and $\rho_{k^\phi}(\widehat{\eta}_{-m}^m) = i^{2m} \rho_{g^{2\phi,0}}(\eta_{-m}^m)$ hold. By Lemma 8.4 we obtain the desired equalities. □

Now we solve the equation for ϕ; $\rho_{k^\phi}(\eta_s^m \eta_t^m) = 1 \otimes \eta_s^m \eta_t^m$ for all $s, t \in I_m$. First we consider $s = t = -m$. We focus on the term of x^{4m} in the both sides. In the left hand side, we have $i^{4m}(\cos 2m\phi\sigma_1 - i \sin 2m\phi\sigma_2)^2 \otimes x^{4m}$. In the right one, we have $1 \otimes x^4$. Hence we have $i^{4m}(\cos 2m\phi\sigma_1 - i \sin 2m\phi\sigma_2)^2 = 1$. It yields $\cos 4m\phi = -1$ and hence ϕ must be equal to $\frac{\pi}{2n}$ or $\frac{3\pi}{2n}$. Then we obtain $\rho_{k^\phi}(\eta_s^m) = \pm i^{2m+1} \sigma_2 \otimes \eta_s^m$ for all $s \in I_m$ with respect to $\phi = \frac{\pi}{2n}$ or $\frac{3\pi}{2n}$. It shows the desired fixed element property of $\eta_s^m \eta_t^m$ by ρ_{k^ϕ} for all $s, t \in I_m$ because of $i^{2m+1} \in \{-1, 1\}$. Therefore we have proved B is a right coideal of type D_n. Note that B does not depend on the choice of $\phi = \frac{\pi}{2n}$ or $\frac{3\pi}{2n}$. In summary, a right coideal of type D_n is made from the quotient $D_n \setminus SU_{-1}(2)$ or generated by a π_m-eigenvector η^m.

As in the previous case of $\mathbb{T}_n$ (odd $n \geq 3$) we can also observe the similar results about a group-like unitary. Here we treat only the case $\chi = 0, \phi = \frac{\pi}{2n}$, that is, Z is $\pi_1^{-1}(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})$ in order to avoid similar arguments. Denote the subgroup G_Z by $(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_\#$. Now Z consists of $\{g^{\frac{2\pi}{n}k}\}_{0 \leq k \leq 2n-1} \cup \{k^\phi g^{\frac{2\pi}{n}\ell}\}_{0 \leq \ell \leq 2n-1}$. Hence the subgroup $(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_\#$ contains $(\mathbb{T}_n^{0,\frac{\pi}{2}})_\#$. We call the subsets $\{g^{\frac{2\pi}{n}k}\}_{0 \leq k \leq 2n-1}$



FIGURE 2. The McKay diagram for $(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_{\#}$, $n+3$ nodes

and $\{k^{\phi}g^{\frac{2\pi}{n}\ell}\}_{0\leq\ell\leq 2n-1}$ the cyclic component and the reflective component, respectively. By definition of the restriction map, we have $C((D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_{\#}) \subset \oplus_{0\leq k\leq 2n-1}\mathbb{M}_{s(k)}(\mathbb{C}) \oplus \oplus_{0\leq\ell\leq 2n-1}\mathbb{M}_{t(\ell)}(\mathbb{C})$ where $s(k)=2$ except for $k=0,2$ and $t(\ell)=2$ except for $\ell=\frac{n-1}{2},\frac{3n-1}{2}$. First note the equality $k^{\phi}g^{\theta}=k^{\phi+\frac{\theta}{2}}$. Then we get

$$\rho_{k^{\phi}g^{\theta}}(\eta_s^m) = i^{2m}\sum_{r\in I_m}(-\sin m\theta\sigma_1-(-1)^{m-r}i\cos m\theta\sigma_2)\otimes(-1)^{m-r}\begin{bmatrix}2m\\m-r\end{bmatrix}^{\frac{1}{2}}w(\pi_m)_{r,s}.$$

Hence it immediately derives $\rho_{k^{\phi}g^{\frac{2\pi}{n}k}}(\eta_s^m) = i^{2m+1}(-1)^{k+1}\sigma_2\otimes\eta_s^m$ for $0\leq k\leq n-1$. Define a self-adjoint unitary $\widetilde{w} = w\oplus\oplus_{\ell\neq\frac{n-1}{2},\frac{3n-1}{2}}(-1)^{\ell+1}i^{2m+1}\sigma_2\oplus 1\oplus -1$ where w is a self-adjoint group-like unitary defined in the previous $\mathbb{T}_n$ case.

Proposition 8.10. *A self-adjoint unitary $\widetilde{w}$ is a group-like element in $C((D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_{\#})$.*

Let $C(\mathbb{Z}_2)$ be a Hopf $*$-subalgebra $\mathbb{C}+\mathbb{C}\widetilde{w}$ in $C((D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_{\#})$. Then we get $B=\{a\in C(SU_{-1}(2)) \mid (\pi_Z\otimes\mathrm{id})\circ\delta(a)\in C(\mathbb{Z}_2)\otimes\mathbb{C}a\}$. We study all the irreducible representations of $(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_{\#}$. The McKay diagram of $(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_{\#}\subset SU_{-1}(2)$ about $\pi_{\frac{1}{2}}$ is shown in Figure 2. In Figure 2, τ_0^+ is the trivial representation and τ_1 is the restriction of $w(\pi_{\frac{1}{2}})$ to $D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}}$. We study where the above group-like unitary $\widetilde{w}$ sits. The restriction of $w(\pi_{\frac{1}{2}})$ to $(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_{\#}$ is denoted simply by $w(\pi_{\frac{1}{2}})|$. For the $\pi_{\frac{1}{2}}$-module W_1 we use an orthonormal basis $\xi_{\pm}=\mathbb{C}\binom{1}{\pm i}$. Then we obtain $w(\pi_{\frac{1}{2}})|\cdot\xi_{\pm}=\xi_{\pm}\otimes w_{\pm}$ for the cyclic component and $w(\pi_{\frac{1}{2}})|\cdot\xi_{\pm}=\xi_{\mp}\otimes v_{\pm}$ for the the reflective component, where $w_{\pm}$ have been already defined in the previous $\mathbb{T}_n$ case and

$$v_{\pm}=\oplus_{\ell\neq\frac{n-1}{2},\frac{3n-1}{2}}\Big(-i\cos\Big(\phi+\frac{\pi}{n}\ell\Big)\sigma_1\pm i\sin\Big(\phi+\frac{\pi}{n}\ell\Big)\sigma_2\Big)\oplus\pm i\oplus\mp i.$$

Hence $v_{\pm}^*=-v_{\pm}$ and $v_{\pm}^2=-1$. Let us consider the tensor product $W_1\otimes W_1$. Its irreducible submodules are $W_0^{\pm}=\mathbb{C}(\xi_+\otimes\xi_+\mp\xi_-\otimes\xi_-)$ and $W_2=\mathbb{C}\xi_+\otimes\xi_-+\mathbb{C}\xi_-\otimes\xi_+$ which give $\tau_0^{\pm}$ and τ_2, respectively. Hence for τ_0^-, the cyclic component acts trivially and the reflective component acts as the multiplication of -1. Proceeding tensor products and decompositions, we get the τ_j-module $W_j = \mathbb{C}\xi_+ \otimes \xi_- \otimes \cdots \xi_- \otimes \xi_+ + \mathbb{C}\xi_- \otimes \xi_+ \otimes \cdots \xi_+ \otimes \xi_-$ for $1 \leq j \leq n-1$, where the length of the words $\xi_\pm$ is j. The cyclic component acts on W_j as the direct sum module of $w_+w_- \cdots w_-w_+$ and $w_-w_+ \cdots w_+w_-$. The reflective component acts there $\upsilon_+\upsilon_- \cdots \upsilon_-\upsilon_+$ and $\upsilon_-\upsilon_+ \cdots \upsilon_+\upsilon_-$. When j is odd, we have

$$\upsilon_\pm\upsilon_\mp \cdots \upsilon_\mp\upsilon_\pm = \oplus_{\ell \neq \frac{n-1}{2}, \frac{3n-1}{2}} i^j \left(-\cos j\left(\phi + \frac{\pi}{n}\ell\right)\sigma_1 \pm \sin j\left(\phi + \frac{\pi}{n}\ell\right)\sigma_2 \right) \oplus \pm i \oplus \mp i.$$

Finally considering the tensor products module $W_1 \otimes W_{n-1}$, we get its one-dimensional submodules $W_n^\pm = \mathbb{C}(\xi_\pm \otimes \xi_\mp \otimes \cdots \xi_\mp \otimes \xi_\pm \pm i\xi_\mp \otimes \xi_\pm \otimes \cdots \xi_\pm \otimes \xi_\mp)$. We investigate the action of $(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_\#$ on them. The cyclic component acts by $w_+w_- \cdots w_-w_+ = w$. Since we have the equality $\upsilon_\pm\upsilon_\mp \cdots \upsilon_\mp\upsilon_\pm = \oplus_{k \neq \frac{n-1}{2}, \frac{3n-1}{2}} i^n (-1)^k (-1)^{\pm 1+1}\sigma_2 \oplus \pm i \oplus \mp i$, the reflection component acts on $W_n^\pm$ by $\oplus_{k \neq \frac{n-1}{2}, \frac{3n-1}{2}} i^{n+1} (-1)^k (-1)^{\frac{\pm 1+1}{2}}\sigma_2 \oplus \pm 1 \oplus \mp 1$. Hence the self-adjoint group-like unitary $\widetilde{w}$ is the unitary representation on W_n^+. Especially we get $\overline{\tau_n^\pm} = \tau_n^\pm$. Recall the classical $q = 1$ case $D_n^* \subset SU(2)$. Then it has the same McKay diagram, however, we know $\overline{\tau_n^\pm} = \tau_n^\mp$. This shows the difference of the representation theory of $(D_n^{\frac{\pi}{2n},0,\frac{\pi}{2}})_\# \subset SU_{-1}(2)$ and $D_n^* \subset SU(2)$.

We have classified all the right coideals of type D_n (odd $n \geq 3$). They are $C(D_n \setminus SU_{-1}(2))$ or $C^*(\eta^m)$ up to conjugacy by the maximal torus action β^L, where $C^*(\eta^m)$ is a right coideal generated by η_r^m for all $r \in I_m$. Finally we end this section with the following result.

Proposition 8.11. *As $SU_{-1}(2)$-covariant systems, $C(D_n \setminus SU_{-1}(2))$ and $C^*(\eta^m)$ are isomorphic.*

Proof. Let g be a matrix in $SU(2)$ $r^{13}(\frac{\pi}{2}) = \begin{pmatrix} \sqrt{2}^{-1} & \sqrt{2}^{-1} \\ -\sqrt{2}^{-1} & \sqrt{2}^{-1} \end{pmatrix}$. Recall the $SU_{-1}(2)$-homomorphism $\rho_g = (\upsilon_g \otimes \mathrm{id}) \circ \delta : C(SU_{-1}(2)) \longrightarrow \mathbb{B}(\mathbb{C}^2) \otimes C(SU_{-1}(2))$. υ_g satisfies $\begin{pmatrix} \upsilon_g(x) & \upsilon_g(u) \\ \upsilon_g(v) & \upsilon_g(y) \end{pmatrix} = \begin{pmatrix} \sqrt{2}^{-1}\sigma_1 & -\sqrt{2}^{-1}\sigma_2 \\ -\sqrt{2}^{-1}\sigma_2 & \sqrt{2}^{-1}\sigma_1 \end{pmatrix}$. Then we obtain

$$\upsilon_g(w(\pi_m)_{-m,r}) = \sqrt{2}^{-2m}(-1)^{m+r}\begin{bmatrix} 2m \\ m-r \end{bmatrix}^{\frac{1}{2}} \sigma_1^{m-r}\sigma_2^{m+r},$$

$$\upsilon_g(w(\pi_m)_{m,r}) = \sqrt{2}^{-2m}(-1)^{m-r}\begin{bmatrix} 2m \\ m-r \end{bmatrix}^{\frac{1}{2}} \sigma_2^{m-r}\sigma_1^{m+r}$$

for all $r \in I_m$. We also have the equality

$$(-1)^{m+r}\sigma_1^{m-r}\sigma_2^{m+r} + (-1)^{m-r}\sigma_2^{m-r}\sigma_1^{m+r} = (-1)^{2m}(\sigma_2 - \sigma_1)$$

for all $r \in I_m$. Then we have

$$\begin{aligned}
\rho_g(x^n + \upsilon^n) &= \rho_g(w(\pi_m)_{-m,-m} + w(\pi_m)_{m,-m}) \\
&= \sum_{r \in I_m} (\upsilon_g(w(\pi_m)_{-m,r}) + \upsilon_g(w(\pi_m)_{m,r})) \otimes w(\pi_m)_{r,-m} \\
&= \sqrt{2}^{-2m} \sum_{r \in I_m} \begin{bmatrix} 2m \\ m-r \end{bmatrix}^{\frac{1}{2}} ((-1)^{m+r} \sigma_1^{m-r} \sigma_2^{m+r} + (-1)^{m-r} \sigma_2^{m-r} \sigma_1^{m+r}) \\
&\qquad\qquad \otimes w(\pi_m)_{r,-m} \\
&= -\sqrt{2}^{-2m} \sum_{r \in I_m} \begin{bmatrix} 2m \\ m-r \end{bmatrix}^{\frac{1}{2}} (\sigma_2 - \sigma_1) \otimes w(\pi_m)_{r,-m} \\
&= -\sqrt{2}^{-2m} (\sigma_2 - \sigma_1) \otimes \eta^m_{-m} .
\end{aligned}$$

Let us define the self-adjoint unitary $\nu = \sqrt{2}^{-1}(\sigma_2 - \sigma_1)$. Then we have $\rho_g(w(\pi_m)_{-m,s} + w(\pi_m)_{m,s}) = -\sqrt{2}^{1-2m} \nu \otimes \eta^m_s$ for all $s \in I_m$. Hence we get $\rho_g(C(D_n \backslash SU_{-1}(2))) \subset C^*(\nu) \otimes C^*(\eta^m)$. Take a $*$-homomorphism $\omega : C^*(\nu) \longrightarrow \mathbb{C}$ and we have an $SU_{-1}(2)$-homomorphism $(\omega \otimes \mathrm{id}) \circ \rho_g : C(D_n \setminus SU_{-1}(2)) \longrightarrow C^*(\eta^m)$. This is clearly surjective and the injectivity follows from Remark 4.23.

$\square$

9. APPENDIX

We list all the connected graphs of norm 2 from Figure 3 to Figure 15 and the spectral patterns of ergodic systems which sit at the vertex of 1 (the entry of the Perron-Frobenius vector). Because we need not to specify the detailed pattern of type A'_m in our classification program, we do not list it in the A'_m case.



FIGURE 3. 1



FIGURE 4. $\mathbb{T}_m$, m nodes



FIGURE 5. $\mathbb{T}$



FIGURE 6. $SU(2)$



FIGURE 7. D_n^* $(n \geq 2)$, $n+3$ nodes



FIGURE 8. D_∞^*



FIGURE 9. A_4^*



FIGURE 10. S_4^*



FIGURE 11. A_5^*



FIGURE 12. D_1



FIGURE 13. D_n (odd $n \geq 3$), $(n+3)/2$ nodes



FIGURE 14. A'_m, $(m \geq 3)$ nodes



FIGURE 15. A'_∞

$$\begin{array}{ll}
1 & \oplus_{\nu\in\frac{1}{2}}(2\nu+1)\pi_\nu, \\
\mathbb{T}_n \ \ (\text{even } n \geq 2) & \oplus_{k\in\mathbb{Z}_{\geq 0}}(1+2\left[\frac{2k}{n}\right])\pi_k, \\
\mathbb{T}_n \ \ (\text{odd } n \geq 3) & \oplus_{k\in\mathbb{Z}_{\geq 0}}\big(1+2\left[\frac{k}{n}\right]\big)\pi_k \oplus \oplus_{k\in\mathbb{Z}_{\geq 0}} 2\left[\frac{2k+n+1}{2n}\right]\pi_{k+\frac{1}{2}}, \\
\mathbb{T} & \oplus_{k\in\mathbb{Z}_{\geq 0}}\pi_k, \\
SU(2) & \pi_0, \\
D^*_m \ \ (m \geq 2) & \oplus_{k\in\mathbb{Z}_{\geq 0}}\big(\frac{1+(-1)^k}{2}+\left[\frac{k}{m}\right]\big)\pi_k, \\
D^*_\infty & \oplus_{k\in\mathbb{Z}_{\geq 0}}\pi_{2k}, \\
A^*_4 & \pi_0 \oplus \pi_3 \oplus \pi_4 \oplus 2\pi_6 \oplus \pi_7 \oplus \cdots, \\
S^*_4 & \pi_0 \oplus \pi_4 \oplus \pi_6 \oplus \pi_8 \oplus \pi_9 \oplus \pi_{10} \oplus \cdots, \\
A^*_5 & \pi_0 \oplus \pi_6 \oplus \pi_{10} \oplus \pi_{12} \oplus \cdots, \\
D_n \ \ (\text{odd } n \geq 1) & \oplus_{k\in\mathbb{Z}_{\geq 0}}\big(\frac{1+(-1)^k}{2}+\left[\frac{k}{n}\right]\big)\pi_k \oplus \oplus_{k\in\mathbb{Z}_{\geq 0}}\big(\left[\frac{2k+1}{n}\right]-\left[\frac{k}{n}\right]\big)\pi_{k+\frac{1}{2}}.
\end{array}$$

REFERENCES

[BS1] S. Baaj, G. Skandalis, *C^*-algèbres de Hopf et théorie de Kasparov équivariante* (French) [Hopf C^*-algebras and equivariant Kasparov theory], K-Theory **2** (1989), no. 6, 683–721.

[BS2] S. Baaj, G. Skandalis, *Unitaires multiplicatifs et dualité pour les produits croisés de C^*-algèbres*(French) [Multiplicative unitary operators and duality for crossed products of C^*-algebras], Ann. Sci. École Norm. Sup. (4) **26** (1993), no. 4, 425–488.

[B] F.P. Boca, *Ergodic actions of compact matrix pseudogroups on C^*-algebras*, Recent advances in operator algebras (Orléans, 1992), Astérisque No. **232** (1995), 93–109.

[DK] M. S. Dijkhuizen, T. H. Koornwinder, *Quantum homogeneous spaces, duality and quantum 2-spheres*, Geom. Dedicata **52** (1994), no. 3, 291–315.

[D] V. G. Drinfel'd, *Hopf algebras and the quantum Yang-Baxter equation*(Russian), Dokl. Akad. Nauk SSSR **283** (1985), no. 5, 1060–1064.

[E] M. Enock, *Sous-facteurs intermédiaires et groupes quantiques mesurés*, J. Operator Theory **42** (1999), 305–330.

[ES] M. Enock, J.-M. Schwartz, *Kac algebras and duality of locally compact groups*, Springer-Verlag, Berlin, 1992.

[GHJ] F. M. Goodman, P. de la Harpe, V. F. R. Jones, *Coxeter graphs and towers of algebras*, Mathematical Sciences Research Institute Publications 14, Springer-Verlag, New York, 1989.

[H-L-S] R. Høegh-Krohn, M.B. Landstad, E. Størmer, *Compact ergodic groups of automorphisms*, Ann. of Math. (2) **114** (1981), no. 1, 75–86.

[J] M. Jimbo, *A q-difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985), no. 1, 63–69.

[Ka] G. G. Kasparov, *Equivariant KK-theory and the Novikov conjecture*, Invent. Math. **91**, (1988), pp. 147-201.

[Ko] T. H. Koornwinder *Askey-Wilson polynomials as zonal spherical functions on the* SU(2) *quantum group*, SIAM J. Math. Anal. **24** (1993), no. 3, 795–813.

[MMNNU] T. Masuda, K. Mimachi, Y. Nakagami, M. Noumi, K. Ueno, *Representations of the quantum group* $\mathrm{SU}_q(2)$ *and the little q-Jacobi polynomials*, J. Funct. Anal. **99** (1991), no. 2, 357–386.

[MNW] T. Masuda, Y. Nakagami, J. Watanabe, *Noncommutative differential geometry on the quantum two sphere of Podleś. I. An algebraic viewpoint*, K-Theory **5** (1991), no. 2, 151–175.

[NM] M. Noumi, K. Mimachi, *Quantum* 2*-spheres and big q-Jacobi polynomials*, Comm. Math. Phys. **128** (1990), no. 3, 521–531.

[Po1] P. Podleś, *Quantum spheres*, Lett. Math. Phys. **14** (1987), no. 3, 193–202.

[Po2] P. Podleś, *Symmetries of quantum spaces. Subgroups and quotient spaces of quantum* SU(2) *and* SO(3) *groups*, Comm. Math. Phys. **170** (1995), no. 1, 1–20.

[S] K. Shiga, *Representations of a compact group on a Banach space*, J. Math. Soc. Japan **7** (1955), 224–248.

[V] R. Vergioux, *KK-théorie équivariante et opérateur de Julg-Valette pour les groupes quantiques*, thesis, universite Paris 7, 2002.

[Wa1] A. Wassermann, *Ergodic actions of compact groups on operator algebras. I. General theory*, Ann. of Math. (2) **130** (1989), no. 2, 273–319.

[Wa2] A. Wassermann, *Ergodic actions of compact groups on operator algebras. III. Classification for* SU(2), Invent. Math. **93** (1988), no. 2, 309–354.

[Wo1] S. L. Woronowicz, *Twisted* SU(2) *group An example of a noncommutative differential calculus*, Publ. Res. Inst. Math. Sci. **23** (1987), no. 1, 117–181.

[Wo2] S. L. Woronowicz, *Compact quantum groups*, Symétries quantiques (Les Houches, 1995), 845–884, North-Holland, Amsterdam, (1998).

[Wo3] S. L. Woronowicz, *Compact matrix pseudogroups*, Comm. Math. Phys. **111** (1987), no. 4, 613–665.

Department of Mathematical Sciences, University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8914, Japan

E-mail address: `tomatsu@ms.u-tokyo.ac.jp`